

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problems Mailbox.**

THIS PAGE BLANK (USPTO)



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets



09/921,397
Priority POC.

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

00402225.7

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk

DEN HAAG, DEN
THE HAGUE, 08/11/01
LA HAYE, LE



THIS PAGE BLANK (USPTO)



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

Blatt 2 der Bescheinigung
Sheet 2 of the certificate
Page 2 de l'attestation

Anmeldung Nr.:
Application no.:
Demande n°: 00402225.7

Anmeldetag:
Date of filing:
Date de dépôt: 03/08/00 ✓

Anmelder:
Applicant(s):
Demandeur(s):
Hybrigenics S.A.
75012 Paris
FRANCE

Bezeichnung der Erfindung:
Title of the invention:
Titre de l'invention:

Sid nucleic acids and polypeptides selected from a pathogenic strain of hepatitis C virus and applications thereof

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s) revendiquée(s)

Staat:
State:
Pays:

Tag:
Date:
Date:

Aktenzeichen:
File no.
Numéro de dépôt:

Internationale Patentklassifikation:
International Patent classification:
Classification internationale des brevets:

C12N15/62, C07K14/18, C12N5/10

Am Anmeldetag benannte Vertragsstaaten:
Contracting states designated at date of filing: AT/BE/CH/CY/DE/DK/ES/FI/FR/GB/GR/IE/IT/LI/LU/MC/NL/PT/SE/TR
Etats contractants désignés lors du dépôt:

Bemerkungen:
Remarks:
Remarques:

**See for original title of the application
page 1 of the description.**

THIS PAGE BLANK (USPTO)

FIELD OF THE INVENTION

The present invention relates to nucleic acids encoding SID® polypeptides which bind selectively to a polypeptide encoded by a pathogenic strain of the hepatitis C virus, as well as to the SID® polypeptides which are encoded by said nucleic acids.

The invention also concerns vectors comprising a nucleic acid encoding a SID® polypeptide as well as host cells transformed with such vectors.

The invention is also directed to two-hybrid methods which make use of the nucleic acids encoding a SID® polypeptide selected from a pathogenic strain of the hepatitis C virus as well as to methods for selecting molecules which inhibit the binding between a SID® polypeptide and a polypeptide which specifically binds thereto.

The invention also pertains to marker compounds containing a SID® polypeptide as well as nucleic acids encoding such marker compounds and methods and kits using the same.

BACKGROUND OF THE INVENTION

The hepatitis C virus (HCV) causes several liver diseases, including liver cancer. The HCV genome is a plus-stranded RNA that encodes the single polypeptide processed into at least 10 mature polypeptides.

The structural proteins are located in the amino terminal quarter of the polypeptide, and the non-structural (NS) polypeptides in the remainder (for a review, see HOUGHTON, 1996). The genome organisation resembles that of flaviviruses and pestiviruses and HCV is now considered to be a member of the *flaviviridae* family.

The gene products of HCV are, from the N-terminus to the C-terminus: core (p22), E1 (gp35), E2 (gp70), NS2(p21), NS3 (p70), NS4a (p4), NS4b(p27), NS5a (p58), NS5b (p66), as disclosed in figure 1. Core, E1 and E2 are the structural proteins of the virus processed by the host
5 signal peptidase(s). The core protein and the genomic RNA constitute the internal viral core and E1 and E2 together with lipid membrane constitute the viral envelop (DUBUISSON et al., 1994; GRAKOU I et al., 1993; HIGIKATA et al. , 1993.).

The NS proteins are processed by the viral protein NS3 which
10 has two functional domains: one (Cro-1), encompassing the NS2 region and the N-terminal portion of NS3, which cleaves autocatalytically between NS2 and NS3, and the other (Cro-2), located solely in the N-terminal portion of NS3, cleaves the other sites downstream NS3 (BARTENSCHLAGER et al; 1995; HIGIKATA et al; 1993).

15 Various HCV protein-protein interactions have already been identified, notably by two hybrid methods. Noticeably, FLAJOLET et al; (2000) have shown interactions between NS3 and NS4A proteins as well as between NS4A and NS2 proteins. These authors have also shown core-core, NS3-E2, NS5A-E1, NS4A-NS3 and NS4A-NS2 interactions.
20 Covalent as well as non-covalent interactions between E1 and E2 have been shown by PATEL et al; (1999). The protein interactions between NS3 and the HCV RNA helicase have also been described (MIN et al; 1999; GALLINARI et al., 1999) as well as interaction between NS3 and NS4A (URBANI et al. , 1999; DI MARCO et al., 2000; BUTKIEWICZ et
25 al. , 2000).

However, the prior art methods allow the determination of interactions between full length proteins or large domains of proteins encoded by the genome of the hepatitis C virus which may contain more than one region of interaction with one or several HCV proteins.
30 BUTKIEWICZ et al. (2000) discloses the interaction between the NS3 protease and a small peptide derived from NS4A. However,

DA.

Le August 2 2000

BUTKIEWICZ et al. (2000) discloses exclusively *in vitro* assays for interactions between the small peptides derived from NS4A and the NS3 protease from HCV which may not be of physiological relevance.

There is a need in the art for polypeptides that contain the
5 minimal aminoacid sequence that is able to bind specifically with a naturally-occurring HCV protein in physiological conditions in order to design new tools for therapeutic and detection purposes related to HCV.

SUMMARY OF THE INVENTION

10

This invention provides nucleic acids encoding polypeptides, which are termed SID® polypeptides, wherein these polypeptides are the final products of a double selection method involving a first step of selection of HCV-derived polynucleotides through a two-hybrid system
15 and a second selection step involving an alignment between the different polynucleotides selected at the first step.

The invention also pertains to the SID® polypeptides encoded by the SID® nucleic acids.

Another object of the invention are recombinant vectors
20 containing a SID® nucleic acid as defined above as well as host cells transformed with such vectors or nucleic acids.

A further object of the invention consists of two-hybrid methods which make use of these SID® nucleic acids as well as to methods for selecting molecules which inhibit the binding between a SID®
25 polypeptide and a polypeptide that binds specifically thereto, as well as kits for performing these methods.

It is still a further object of the invention to provide for marker compounds which comprise a SID® polypeptide or which are encoded by a polynucleotide containing a SID® nucleic acid as defined above, as

well as to methods and kits which make use of these marker compounds.

This invention also relates to pharmaceutical compositions as well as to methods for preventing or curing a HCV viral infection in a
5 human or an animal that use a SID® polypeptide or a SID® nucleic acid as disclosed herein.

Throughout this application, various publications, patents and published patent applications are cited. The disclosures of these publications, patents and published patent specifications, referenced in
10 this application are hereby incorporated by reference into the present disclosure to more fully describe the state of the art to which this invention pertains.

BRIEF DESCRIPTION OF THE FIGURES.

15

Figure 1 consists of a general overview of HCV genome and its encoded polyprotein. The RNA coding strand is represented with a line for untranslated regions (NCR) and boxes for coding regions.

Positions and enzymes responsible for cleavage are indicated
20 above. p7 is a secondary cleavage product of E2 (adapted from HOUGHTON, 1996).

Fig. 2 is a restriction map of the plasmid pAS2ΔΔ which may be used for producing a recombinant " Selected Interacting Domain (SID®) " polypeptide or a recombinant marker compound of the invention.

25 Fig. 3 is a restriction map of the plasmid pACTII which may be used for producing a recombinant " Selected Interacting Domain (SID®) ".

Fig. 4 is a restriction map of the plasmid pUT18 which may be used for producing a recombinant " Selected Interacting Domain
30 (SID®) ".

DA.
Le August 2 2000

Fig. 5 is a restriction map of the plasmid pUT18C which may be used for producing a recombinant "Selected Interacting Domain (SID®)".

Fig. 6 is a restriction map of the plasmid pT25 which may be used for producing a recombinant "Selected Interacting Domain (SID®)".

Fig. 7 is a restriction map of the plasmid pKT25 which may be used for producing a recombinant "Selected Interacting Domain (SID®)".

Fig.8 is an illustration of the first step of selecting a SID® nucleic acid of the invention, wherein it is performed a selection of different sets of overlapping nucleic acids primarily selected through a two-hybrid method, in order to define pre-SID nucleic acids. Three fragments frg1, frg2 and frg3 of lengths l1, l2 and l3 respectively. Fragment l1 and l2 are clustered together if the length of intersection, l, is greater than 30% of l1 and l2. Fragment frg3 is grouped with fragments frg1 and frg2 if the length of intersection between frg1 and frg3, l', is greater than 30% of l1 and l3 and if the length of intersection between frg 2 and frg 3, l », is greater than 30% of l2 and l3.

Fig.9 illustrates the selection of pre-SID® nucleic acid from a particular set of overlapping nucleic acids previously selected through a two-hybrid method. The pre-SID® is defined as the intersection of all the fragments (frg1-6) in a cluster.

Fig.10 illustrates the selection of a SID® nucleic acid from the overlapping regions between two pre-SID nucleic acids. A SID® is defined if the length of overlap between two pre-SID®, l, is greater than 30 bp. Further SID®s are defined by non-overlapping areas if their length (l') represents more than 30% of the length of one of the fragments which contributes to the corresponding pre-SID® (frg1-6).

Fig.11 illustrates a further step of determining SID® nucleic acids after alignment of two overlapping SID nucleic acids identified according to figure 10. Fragments frg1' and frg2' contribute to both SID®1 and SID®2 (top panel). For each SID®, the number of fragments are counted and fragments are assigned to the SID® with the most fragments. The remaining fragments are re-analysed and a new SID® is

defined as the region of intersection of these fragments (bottom panel, SID®2' - fragment 3' and fragment 4'.

Fig.12 illustrates a map of the vector pB5 which may be used in example 1.

5 Fig.13 illustrates a map of the vector pP6 which may be used in example 1.

DETAILED DESCRIPTION OF THE INVENTION

10 The present invention firstly provides for nucleic acids encoding SID® polypeptides.

As generally used herein, a « bait » nucleic acid encodes a « bait » polypeptide. A polypeptide is termed a « bait » polypeptide when
15 this polypeptide is used to select a formerly unknown « prey » nucleic acid encoding a « prey » polypeptide which binds selectively with said « bait » polypeptide. Indeed, a « prey » nucleic acid which has been selected for binding to a given bait polypeptide may be used in another selection method or in another round of the same selection method as a
20 « bait » nucleic acid encoding a « bait » polypeptide for the purpose of selection of new prey nucleic acids, encoding prey polypeptides which bind selectively with said bait polypeptide, it being understood that the nucleic acid encoding said bait polypeptide was formerly selected from a population of prey nucleic acids.

25

SELECTED INTERACTING DOMAIN (SID®) POLYPEPTIDES AND METHODS FOR THEIR PREPARATION.

30 A selected interacting domain polypeptide that binds specifically to a polypeptide of interest is the result of a two-step screening procedure, wherein :

DA.

Le August 2 2000

1) the first step consists of selecting and characterizing a collection of nucleic acids (prey nucleic acids) encoding polypeptides which bind specifically to a given bait polypeptide of interest;

and

5 2) the second step of the two-step procedure consists of determining the nucleic acid sequences which encode for SID® polypeptides after having generated sets of polynucleotides from the collection of nucleic acids selected at step 1).

10 As a result of the original two-step screening procedure disclosed hereunder, every nucleic acid finally selected encodes a « Selected Interacting Domain (SID®) » polypeptide which binds with a high specificity with the bait polypeptide of interest.

Step 1) Selecting prey nucleic acids

15

 The first step of selecting a collection of nucleic acids encoding polypeptides which binds specifically to the bait polypeptide is carried out through a yeast two-hybrid system. The yeast two-hybrid system is designed to study protein-protein interactions *in vivo*, and relies upon the fusion of a bait protein to the DNA binding domain of the yeast Gal4 protein.

20 According to the present invention, the first step of the procedure for selecting a Selected Interacting Domain (SID®) polynucleotide encoding a Selected Interacting Domain (SID®) polypeptide consists of the two-hybrid screening system described by Fromont-Racine et al. (1997) or the method described by FLAJOLET et al. (2000). The yeast two-hybrid system utilizes hybrid proteins to detect protein-protein interactions by means of direct activation of a reporter gene expression. In essence, the nucleic acids encoding the two putative protein partners, the bait polypeptide of interest and the prey polypeptide, are genetically fused to the DNA-binding domain of a transcription factor and to a transcriptional activation domain, respectively.

Construction of the prey HCV nucleic acids library.

Then, a genomic DNA library prepared from the genome of the
5 pathogenic H77 strain of HCV (Yanagi et al., 1997), is constructed in the
specially designed vector pP6 shown in figure 13 after ligation to suitable
linkers, such that every genomic DNA insert is fused to a nucleotide
sequence in the vector that encodes the transcription of domain of the
Gal4 protein.

10 The polypeptides encoded by the nucleotide inserts of the
genomic DNA library thus prepared are termed "prey" polypeptides in
the context of the presently described selection method of prey nucleic
acids.

Construction of the bait nucleic acids library

The DNA fragments obtained after nebulization of the HCV
genomic DNA are also inserted in plasmid pB5 shown in figure 12
wherein these DNA inserts are fused to a polynucleotide encoding the
20 DNA binding domain of the Gal4 protein and the recombinant vectors are
used to transform *E. coli* cells. The transformed *E. coli* cells are grown
and plasmid DNA is extracted and sequenced.

These plasmids which code in frame fusion proteins are used
as bait plasmids. Bait plasmids thus consist of a collection of
25 recombinant pB5 plasmids each containing inserted therein a DNA
fragment from the H77 strain HCV genome encoding a polypeptide
consisting of all or part of a HCV protein or alternatively a polypeptide
consisting of all or part of two HCV proteins encoded by contiguous
nucleic acid sequences of the HCV genome.

30 The selected HCV bait nucleic acids of the invention are
referred to as the nucleotide sequences SEQ ID N°114 to 150.

The selected HCV bait polypeptides encoded by the nucleic
sequences SEQ ID N°114 to 150 consist respectively of the aminoacid
sequences SEQ ID N°77 to 113.

DA.
Le August 2 2000

Detectable marker genes are already present within the chromosomal yeast DNA and consist respectively of the His3 and LacZ genes, such as described by FROMONT-RACINE et al. (1997) or FLAJOLET et al. (2000).

5 Then, the collection of nucleic acid inserts contained in the collection of *E. Coli* cell clones containing the genomic DNA or HCV DNA library previously prepared are used to transform a first yeast strain, namely the Y187 *Saccharomyces cerevisiae* strain (phenotype:MAT α , Gal4 Δ , gal80 Δ , ade2-101, His3, Leu2-3, -112 Trp1-901, Ura3-52, 10 URA3::UASGAL1-LacZ Met).

 The nucleic acid encoding the bait polypeptide of interest is inserted in the appropriate vector, said vector being used to transform a second yeast strain which may be the CG1945 (MAT α Gal4-542 Gal180-538, Ade2-101, His3*200, Leu2-3, -112 Trp1-901 Ura3-52, Lys2-801, 15 URA3::GAL4 17Mers (X3)-CyC1TATA-LacZ LYS2::GAL1 UAS-GAL1TATA-His3 CYH^R).

 Then, the two yeast strains are mated to obtain a collection of mated cells.

 The clones derived from the collection of mated cells above 20 which are positive in an X-Gal overlay assay are those for which an interaction between the recombinant bait polypeptide and a polypeptide encoded by a nucleic acid insert originating from the HCV genomic library has occurred.

 The clones derived from the collection of mated cells above 25 may also be selected in the presence of histidine, and the positive clones are those for which an interaction between the recombinant bait polypeptide and a polypeptide encoded by a nucleic acid insert originating from the HCV genomic library has occurred.

 In a further step, the prey nucleic acid inserts contained in the 30 positively selected clones are amplified and sequenced.

Step 2:determination of the nucleic acid sequences encoding a Selected Interacting Domain (SID®) polypeptide which binds specifically to a bait polypeptide of interest.

35

This is the second step of the two step procedure defined above, which allows the precise selection of nucleic acids encoding the SID® nucleic acids of the present invention which are derived from the H77 strain HCV genome.

5 The SID® nucleic acid selection procedure, which is disclosed hereunder, has been specifically designed for the HCV genome which encodes for a single polyprotein and which thus comprises contiguous Open Reading Frames, said polyprotein being further processed to produce at least 10 mature structural and non-structural viral proteins.

10 Thus, the second selection step of the two-step procedure consists of a method for determining a polynucleotide encoding a Selected Interacting Domain (SID®) of a prey polypeptide of interest derived from HCV, which prey polypeptide interacts with a bait polypeptide, wherein said method comprises the steps of :

15 a) selecting, from the collection of prey polynucleotides obtained at the end of the first step of the two-step procedure described herein, all prey polynucleotides encoding a prey polypeptide capable of interacting with said bait polypeptide and containing a common nucleic acid fragment;

20 b) aligning the nucleotide sequences of the prey polynucleotides selected at step a) and gathering in one set or in a plurality of sets of sequences those nucleotide sequences which have sequences that overlap for more than 30% of their respective nucleic acid length, wherein each common overlapping nucleotide sequence in
25 one set of sequences defines a sequence encoding a pre-SID® polypeptide (see Figures 8 and 9); and

 c) aligning two sequences encoding two respective pre-SID® polypeptides (see Figure 10), and :

30 i) defining an overlapping nucleic acid sequence between the sequences encoding the two respective pre-SID® polypeptides as a

DA.

Le August 2 2000

sequence encoding a SID® polypeptide, provided that the overlapping sequence is of at least 30 nucleotides in length;

ii) defining a non-overlapping nucleic acid sequence between the sequences encoding the two respective pre-SID® polypeptides as a sequence encoding a SID® polypeptide, provided that (1) said non-overlapping sequence has more than 30 nucleotides in length and (2) said non-overlapping sequence represents at least 30% in length of any one of the polynucleotides contained in the set of prey polynucleotides used for defining the sequence encoding each pre-SID® polypeptide.

10 This method may further comprise the steps of :

d) counting the number of overlapping prey polynucleotides contained in a first set of polynucleotides defining a sequence encoding a first SID® polypeptide;

e) counting the number of overlapping prey polynucleotides contained in a second set of polynucleotides defining a sequence encoding a second SID® polypeptide which overlaps with the sequence encoding the first SID® polypeptide;

f) determining which sequence among those encoding respectively the first SID® polypeptide and the second SID® polypeptide has been defined with the largest number of prey polynucleotides and selecting this set of prey sequences.

g) adding to the set of prey sequences selected at step f) those sequences that were contained in the set of prey sequences used for defining the sequence encoding the SID® polypeptide with the smallest number of prey sequences and which overlap with the sequence encoding the SID® polypeptide with the largest number of prey sequences.;

h) aligning the prey sequences added at step g) with the sequences already contained in the set of prey sequences which defined the sequence encoding the SID® polypeptide with the largest number of prey sequences;

i) defining an overlapping sequence between the whole sequences which were aligned in step h), wherein said overlapping sequence consists of a sequence encoding a SID® polypeptide. (See Figure 11).

5 The method for selecting a SID® nucleic acid encoding a SID® polypeptide is an object of the present invention, as well as any SID® nucleic acid or any SID® polypeptide which may be obtained by this selection method.

10 **SID® nucleic acids of the invention**

The SID® nucleic acids selected as described above starting from the genome of the H77 strain of HCV are the nucleic acid sequences of SEQ ID N°39 to 76 which encode the SID® polypeptides of SEQ ID N°1 to 38.

15 A first object of the invention consists of a nucleic acid which encodes a polypeptide selected from the group consisting of the aminoacid sequences SEQ ID N°1 to 38 or a variant thereof, and a sequence complementary thereto.

For the purposes of the present invention, a first polynucleotide
20 is considered as being « complementary » to a second polynucleotide when each base of the first polynucleotide is paired with the complementary base of the second polynucleotide whose orientation is reversed. The complementary bases are A and T(or A and U), or C and G.

25 Preferably, any one of the nucleic acid or the polypeptides encompassed by the invention is under a purified or an isolated form.

The term "isolated" for the purposes of the present invention designates a biological material (nucleic acid or protein) which has been removed from its original environment (the environment in which it is
30 naturally present).

DA.

Le August 2 2000

For example, a polynucleotide present in the natural state in a plant or an animal is not isolated. The same polynucleotide separated from the adjacent nucleic acids in which it is naturally inserted in the genome of the plant or animal is considered as being "isolated".

5 Such a polynucleotide may be included in a vector and/or such a polynucleotide may be included in a composition and remains nevertheless in the isolated state because of the fact that the vector or the composition does not constitute its natural environment.

10 The term "purified" does not require the material to be present in a form exhibiting absolute purity, exclusive of the presence of other compounds. It is rather a relative definition.

A polynucleotide is in the "purified" state after purification of the starting material or of the natural material by at least one order of magnitude, preferably 2 or 3 and preferably 4 or 5 orders of magnitude.

15 "Isolated polypeptide" or "isolated protein" is a polypeptide or protein which is substantially free of those compounds that are normally associated therewith in its natural state (e.g., other proteins or polypeptides, nucleic acids, carbohydrates, lipids). "Isolated" is not meant to exclude artificial or synthetic mixtures with other compounds, or
20 the presence of impurities which do not interfere with biological activity, and which may be present, for example, due to incomplete purification, addition of stabilisers, or compounding into a pharmaceutically acceptable preparation.

25 ***Variants of a selected interacting domain (SID®) polypeptide and nucleic acids encoding them.***

As intended herein, a variant of a Selected Interacting Domain (SID®) polypeptide may be either a variant polypeptide of the Selected
30 Interacting Domain (SID®) polypeptide or a polypeptide which is encoded by a nucleic acid variant of the polynucleotide encoding said Selected Interacting Domain (SID®) polypeptide.

Polynucleotides which encode a polypeptide variant of a Selected Interacting Domain (SID®) polypeptide, as the term is used
35 herein, are polynucleotides that differ from the reference polynucleotide

encoding the parent SID® polypeptide. A variant of a polynucleotide may be a naturally occurring variant such as a naturally occurring allelic variant, or it may be a variant that is not known to occur naturally. Such non-naturally occurring variants of the reference polynucleotide may be generated by mutagenesis techniques, including those applied to polynucleotides, cells or organisms well known to one skilled in the art.

Generally, differences are limited so that the nucleotide sequences of the reference and the variant are closely similar overall and, in many regions, identical.

Variants of polynucleotides according to the invention include, without being limited to, nucleotide sequences which are at least 95% identical after optimal alignment to the reference polynucleotide of SEQ ID N°39 to 76 encoding the reference Selected Interacting Domain (SID®) polypeptide, preferably at least 96%, 97%, 98% and most preferably at least 99% identical to the reference polynucleotide. Similarly, a variant of a SID® polypeptide of the invention consists of a polypeptide having at least 95% aminoacid identity with a polypeptide selected from the aminoacid sequences SEQ ID N°1 to 38, and preferably at least 96%, 97%, 98% and most preferably at least 99% aminoacid identity with one of SEQ ID N°1 to 38.

Identity refers to sequence identity between two peptides or between two nucleic acid molecules. Identity between sequences can be determined by comparing a position in each of the sequences which may be aligned for purposes of comparison. When a position in the compared sequences is occupied by the same base or amino acid, then the sequences are identical at that position. A degree of identity between nucleic acid sequences is a function of the number of identical nucleotides at positions shared by these sequences. A degree of identity between amino acid sequences is a function of the number of identical aminoacids at positions shared by these sequences. Since two polynucleotides may each (1) comprise a sequence (i.e., a portion of the complete polynucleotide sequence) that is similar between the two

DA.

Le August 2 2000

polynucleotides, and (2) may further comprise a sequence that is divergent between the two polynucleotides, sequence comparisons between two (or more) polynucleotides are typically performed by comparing sequences of the two polynucleotides over a "comparison window" to identify and compare local regions of sequence similarity. A "comparison window", as used herein, refers to a conceptual segment of at least 20 contiguous nucleotide positions wherein a polynucleotide sequence may be compared to a reference sequence of at least 20 contiguous nucleotides and wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) of 20 percent or less as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. Optimal alignment of sequences for determining a comparison window may be conducted by the local homology algorithm of Smith and Waterman (1981), by the homology alignment algorithm of Needleman and Wunsch (1972), by the search for similarity method of Pearson and Lipman (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package Release 7.0, Genetics Computer Group, 575, Science Dr. Madison, WI), or by inspection. The best alignment (i.e., resulting in the highest percentage of identity over the comparison window) generated by the various methods is selected. The term "sequence identity" means that two polynucleotide sequences are identical (i.e., on a nucleotide-by-nucleotide basis) over the window of comparison. The term "percentage of sequence identity" is calculated by comparing two optimally aligned sequences over the window of comparison, determining the number of positions at which the identical nucleic acid base (e.g. A, T, C, G, U or I) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison

(i.e., the window size), and multiplying the result by 100 to yield the percentage of sequence identity.

Most preferably, the percentage of nucleic acid or aminoacid identity between two nucleic acid or aminoacid sequences is calculated
5 using the BLAST software (Version 2.06 of September 1998) with the default parameters.

Nucleotide changes present in a variant polynucleotide may be silent, which means that they do not alter the aminoacid encoded by the reference polynucleotide.

10 However, nucleotide changes may also result in aminoacid substitutions, additions, deletions, fusions and truncations in the Selected Interacting Domain (SD®) polypeptide encoded by the reference sequence.

The substitutions, deletions or additions may involve one or
15 more nucleotides. Alterations may produce conservative or non-conservative aminoacid substitutions, deletions or additions.

Most preferably, the variant of a Selected Interacting Domain (SID®) polypeptide encoded by a variant polynucleotide possesses at least the same affinity of binding to its protein or polypeptide counterpart,
20 against which it has been initially selected as described above.

The affinity of a given SID® polypeptide of the invention for a polypeptide into which it specifically binds is defined as the affinity constant K_a , wherein

25

$$K_a = \frac{[\text{SID®/polypeptide complex}]}{[\text{free SID®}] [\text{free polypeptide}]}$$

30 with [free SID®], [free polypeptide] and [SID®/polypeptide complex] consist of the concentrations at equilibrium respectively of the free SID® polypeptide, of the free polypeptide onto which the SID® polypeptide specifically binds and of the complex formed between the SID®

DA.

Le August 2 2000 .

polypeptide and the polypeptide onto which said SID® polypeptide specifically binds.

Most preferably, the affinity of a SID® polypeptide of the invention or a variant thereof for its polypeptide counterpart (polypeptide partner) is assessed on a Biacore™ apparatus marketed by Amercham Pharmacia Biotech Company such as described by SZABO et al. (1995) and by Edwards and Leartherbarrow (1997).

As used herein, the expression « at least the same affinity» with reference to the affinity of binding between a SID® polypeptide of the invention to another polypeptide means that the K_a is identical or is of at least two-fold, preferably at least three-fold and most preferably at least five-fold greater than the K_a value of reference.

In another preferred embodiment, the variant of a Selected Interacting Domain (SID®) polypeptide which is encoded by a variant polynucleotide of the invention possesses a higher specificity of binding to its counterpart polypeptide or protein than the reference Selected Interacting Domain (SID®) polypeptide.

A variant of a Selected Interacting Domain (SID®) polypeptide according to the invention may be (1) one in which one or more, most preferably from one to three, of the aminoacid residues are substituted with a conserved or a non-conserved aminoacid residue and such substituted aminoacid residue may or may not be one encoded by the genetic code, or (2) one in which one or more of the aminoacid residues includes a substituent group.

In the case of an aminoacid substitution in the aminoacid sequence of a Selected Interacting Domain (SID®) polypeptide according to the invention, one or several-consecutive or non-consecutive - aminoacids are replaced by "equivalent" aminoacids. The expression "equivalent" aminoacid is used herein to designate any aminoacid that may be substituted for one of the aminoacids belonging to the native Selected Interacting Domain (SID®) polypeptide structure without decreasing the binding properties of the corresponding peptides to their counterpart polypeptide or protein, as regards the reference Selected Interacting Domain (SID®) polypeptide.

These equivalent aminoacids may be determined either by their structural homology with the initial aminoacids to be replaced, by the similarity of their net charge or of their hydrophobicity.

By an equivalent aminoacid according to the present invention
 5 is also meant the replacement of a residue in the L-form by a residue in the D-form or the replacement of a glutamic acid residue by a pyro-glutamic acid compound. The synthesis of peptides containing at least one residue in the D-form is, for example, described by KOCH (1977). A specific embodiment of a variant of a Selected Interacting Domain
 10 (SID®) polypeptide according to the invention includes, but is not limited to, a peptide molecule which is resistant to proteolysis, such as a peptide in which the -CONH- peptide bond is modified and replaced by a (-CH₂NH-) reduced bond, a (-NHCO-) retroinverso bond, a (-CH₂-O-) methylene-oxy bond, a (-CH₂-S-) thiomethylene bond, a (-CH₂CH₂-)
 15 carba bond, a (-CO-CH₂) hydroxyethylene bond, a (-N-N-) bond or also a -CH=CH bond.

As used herein, a variant of a SID® polypeptide of the invention also encompasses a polypeptide having an aminoacid sequence consisting of at least:

- 20 - 45 consecutive aminoacids of SEQ ID N°1;
- 30 consecutive aminoacidss of SEQ ID N°2;
- 65 consecutive aminoacids of SEQ ID N°3;
- 30 consecutive aminoacids of SEQ ID N°4;
- 130 consecutive aminoacids of SEQ ID N°5;
- 25 - 25 consecutive aminoacids of SEQ ID N°6;
- 23 consecutive aminoacids of SEQ ID N°7.
- 48 consecutive aminoacids of SEQ ID N°8;
- 36 consecutive aminoacids of SEQ ID N°9;
- 25 consecutive aminoacids of SEQ ID N°10;
- 30 - 24 consecutive aminoacids of SEQ ID N°11;
- 37 consecutive aminoacids of SEQ ID N°12;
- 25 consecutive aminoacids of SEQ ID N°13;
- 30 consecutive aminoacids of SEQ ID N°14;
- 27 consecutive aminoacids of SEQ ID N°15;
- 35 - 69 consecutive aminoacids of SEQ ID N°16;

DA.

Le August 2 2000

- 130 consecutive aminoacids of SEQ ID N°17;
- 33 consecutive aminoacids of SEQ ID N°18;
- 25 consecutive aminoacids of SEQ ID N°19;
- 40 consecutive aminoacids of SEQ ID N°20;
- 5 - 78 consecutive aminoacids of SEQ ID N°21;
- 39 consecutive aminoacids of SEQ ID N°22;
- 57 consecutive aminoacids of SEQ ID N°23;
- 26 consecutive aminoacids of SEQ ID N°24;
- 68 consecutive aminoacids of SEQ ID N°25;
- 10 - 34 consecutive aminoacids of SEQ ID N°26;
- 42 consecutive aminoacids of SEQ ID N°27;
- 48 consecutive aminoacids of SEQ ID N°28.
- 102 consecutive aminoacids of SEQ ID N°29;
- 49 consecutive aminoacids of SEQ ID N°30;
- 15 - 92 consecutive aminoacids of SEQ ID N° 31;
- 49 consecutive aminoacids of SEQ ID N°30;
- 92 consecutive aminoacids of SEQ ID N°31;
- 71 consecutive aminoacids of SEQ ID N°32;
- 55 consecutive aminoacids of SEQ ID N°33;
- 20 - 69 consecutive aminoacids of SEQ ID N°34;
- 23 consecutive aminoacids of SEQ ID N°35;
- 33 consecutive aminoacids of SEQ ID N°36;
- 32 consecutive aminoacids of SEQ ID N°37;
- and
- 25 - 22 consecutive aminoacids of SEQ ID N°38.

Without wishing to be bound by any particular theory, the inventors believe that polypeptides having an aminoacid length of about 10% lesser than the aminoacid length of anyone of the SID®

30 polypeptides of SEQ ID N°1 to 39 of the invention have a high probability to retain the binding properties to a given (bait) polypeptide of the parent SID® polypeptide.

The invention also pertains to a nucleic acid encoding a SID® polypeptide which is selected from the group consisting of the sequences

35 SEQ ID N°39 to 76, and a sequence complementary thereto.

The invention is also directed to a nucleic acid encoding a variant of SID® polypeptide selected from the group consisting of the sequences SEQ ID N°39 to 76, in reference to the definition of the SID® polypeptide variants above.

5 For example, a nucleic acid encoding a polypeptide having an aminoacid sequence consisting of at least 45 consecutive aminoacids of SEQ ID N°1 comprise at least 135 (45 x 3) consecutive nucleotides of the polynucleotide of SEQ ID N°39.

 The same definition also apply for nucleic acids encoding
10 variants of the SID® polypeptides of SEQ ID N°2 to 38, which are part of the invention.

 The invention further relates to a nucleic acid encoding a polypeptide having an aminoacid sequence comprising from 1 to 3 substitutions, additions or deletions of one aminoacid as regards a
15 polypeptide selected from the group consisting of the aminoacid sequences SEQ ID N°1 to 38 or a sequence complementary thereto.

 Another object of the invention consists of a polypeptide selected from the group consisting of the aminoacid sequences SEQ ID
... N°39 to 76 or a variant thereof.

20 Are encompassed in the family of variants of a SID® polypeptide of the invention those polypeptides having an aminoacid sequence comprising from 1 to 3 substitutions, additions or deletions of one aminoacid as regards a polypeptide selected from the group consisting of the aminoacid sequences SEQ ID N°1 to 38.

25 The invention is also directed to an antibody directed against a SID® polypeptide as defined above, or to a variant thereof.

 The antibodies directed specifically against the Selected Interacting Domain (SID®) polypeptide or a variant thereof may be indifferently radioactively or non-radioactively labelled.

30 Monoclonal antibodies directed against a SID® polypeptide may be prepared from hybridomas according to the technique described

DA.
Le August 2 2000

by Kohler and Milstein in 1975. Polyclonal antibodies may be prepared by immunization of a mammal, especially a mouse or a rabbit, with the SID® polypeptide that is combined with an adjuvant of immunity, and then by purifying the specific antibodies contained in the serum of the immunized animal on a affinity chromatography column on which has previously been immobilized the polypeptide that has been used as the antigen.

Antibodies directed against a SID® polypeptide may also be produced by the trioma technique and by the human B-cell hybridoma technique (Kozbor et al., 1983).

Antibodies directed to a SID® polypeptide include chimeric single chain Fv antibody fragments (US Patent N° US 4,946,778; Martineau et al., 1998), antibody fragments obtained through phage display libraries (Ridder et al., 1995) and humanized antibodies (Reinmann et al., 1997; Leger et al., 1997). Also, transgenic mice, or other organisms such as other mammals, may be used to express antibodies, including for example, humanized antibodies directed against a SID® polypeptide of the invention, or a variant thereof.

VECTORS OF THE INVENTION

The nucleic acids coding for a Selected Interacting Domain (SID®) polypeptide or a variant thereof, which are defined in the section above, can be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for the transcription and translation of the inserted protein-coding sequence. Such transcription elements include a regulatory region and a promoter as defined previously. Thus, the nucleic acid encoding a marker compound of the invention is operably linked with a promoter in a expression vector, wherein said expression vector may include a replication origin.

The necessary transcriptional and translation of signals is most preferably provided by the recombinant expression vector.

Structure of the vectors encompassed by the invention

A wide variety of host/expression vector combinations may be employed in expressing the nucleic acids of this invention. Useful expression vectors, for example, may consist of segments of chromosomal, non-chromosomal and synthetic DNA sequences. Suitable vectors include derivatives of SV40 and known bacterial plasmids, e.g., *Escherichia coli* plasmids col EI, pCR1, pBR322, pMal-C2, pET, pGEX (Smith *et al.*, 1988), pMB9 and their derivatives, plasmids such as RP4; phage DNAs, e.g., the numerous derivatives of phage I, e.g., NM989, and other phage DNA, e.g., M13 and filamentous single stranded phage DNA; yeast plasmids such as the 2m plasmid or derivatives thereof; vectors useful in eukaryotic cells, such as vectors useful in insect or mammalian cells; vectors derived from combinations of plasmids and phage DNAs, such as plasmids that have been modified to employ phage DNA or other expression control sequences; and the like.

For example, in a baculovirus expression system, both non-fusion transfer vectors, such as but not limited to pVL941 (*Bam*H1 cloning site; Summers), pVL1393 (*Bam*H1, *Sma*I, *Xba*I, *Eco*R1, *Not*I, *Xma*III, *Bgl*II, and *Pst*I cloning site; Invitrogen), pVL1392 (*Bgl*II, *Pst*I, *Not*I, *Xma*III, *Eco*R1, *Xba*I, *Sma*I, and *Bam*H1 cloning site; Summers and Invitrogen), and pBlueBacIII (*Bam*H1, *Bgl*II, *Pst*I, *Nco*I, and *Hind*III cloning site, with blue/white recombinant screening possible; Invitrogen), and fusion transfer vectors, such as but not limited to pAc700 (*Bam*H1 and *Kpn*I cloning site, in which the *Bam*H1 recognition site begins with the initiation codon; Summers), pAc701 and pAc702 (same as pAc700, with different reading frames), pAc360 (*Bam*H1 cloning site 36 base pairs downstream of a polyhedrin initiation codon; Invitrogen(195)), and pBlueBacHisA, B, C (three different reading frames, with *Bam*H1, *Bgl*II, *Pst*I, *Nco*I, and *Hind*III cloning site, an N-terminal peptide for ProBond purification, and blue/white recombinant screening of plaques; Invitrogen (220) can be used.

Mammalian expression vectors contemplated for use in the invention include vectors with inducible promoters, such as the dihydrofolate reductase (DHFR) promoter, e.g., any expression vector with a *DHFR* expression vector, or a *DHFR*/methotrexate co-

DA.

Le August 2 2000

amplification vector, such as pED (*Pst*I, *Sal*I, *Sba*I, *Sma*I, and *Eco*RI cloning site, with the vector expressing both the cloned gene and *DHFR*; Kaufman,1991). Alternatively, a glutamine synthetase/methionine sulfoximine co-amplification vector, such as pEE14 (*Hind*III, *Xba*I, *Sma*I, *Sba*I, *Eco*RI, and *Bcl*I cloning site, in which the vector expresses glutamine synthase and the cloned gene; Celltech). In another embodiment, a vector that directs episomal expression under control of Epstein Barr Virus (EBV) can be used, such as pREP4 (*Bam*H1, *Sfi*I, *Xho*I, *Not*I, *Nhe*I, *Hind*III, *Nhe*I, *Pvu*II, and *Kpn*I cloning site, constitutive RSV-LTR promoter, hygromycin selectable marker; Invitrogen), pCEP4 (*Bam*H1, *Sfi*I, *Xho*I, *Not*I, *Nhe*I, *Hind*III, *Nhe*I, *Pvu*II, and *Kpn*I cloning site, constitutive hCMV immediate early gene, hygromycin selectable marker; Invitrogen), pMEP4 (*Kpn*I, *Pvu*I, *Nhe*I, *Hind*III, *Not*I, *Xho*I, *Sfi*I, *Bam*H1 cloning site, inducible methallothionein Ila gene promoter, hygromycin selectable marker: Invitrogen), pREP8 (*Bam*H1, *Xho*I, *Not*I, *Hind*III, *Nhe*I, and *Kpn*I cloning site, RSV-LTR promoter, histidinol selectable marker; Invitrogen), pREP9 (*Kpn*I, *Nhe*I, *Hind*III, *Not*I, *Xho*I, *Sfi*I, and *Bam*HI cloning site, RSV-LTR promoter, G418 selectable marker; Invitrogen), and pEBVHis (RSV-LTR promoter, hygromycin selectable marker, N-terminal peptide purifiable via ProBond resin and cleaved by enterokinase; Invitrogen). Selectable mammalian expression vectors for use in the invention include pRc/CMV (*Hind*III, *Bst*XI, *Not*I, *Sba*I, and *Apa*I cloning site, G418 selection; Invitrogen), pRc/RSV (*Hind*III, *Spe*I, *Bst*XI, *Not*I, *Xba*I cloning site, G418 selection; Invitrogen), and others. Vaccinia virus mammalian expression vectors (see, Kaufman, 1991, *supra*) for use according to the invention include but are not limited to pSC11 (*Sma*I cloning site, TK- and b-gal selection), pMJ601 (*Sal*I, *Sma*I, *Afl*I, *Nar*I, *Bsp*MII, *Bam*HI, *Apa*I, *Nhe*I, *Sac*II, *Kpn*I, and *Hind*III cloning site; TK- and b-gal selection), and pTKgptF1S (*Eco*RI, *Pst*I, *Sal*I, *Acc*I, *Hind*II, *Sba*I, *Bam*HI, and *Hpa*I cloning site, TK or XPRT selection).

Yeast expression systems can also be used according to the invention to express a Selected Interacting Domain (SID®) polypeptide or a variant thereof and also a marker compound as defined herein. For example, the non-fusion pYES2 vector (*Xba*I, *Sph*I, *Sho*I, *Not*I, *Gst*XI,

EcoRI, *BstXI*, *BamHI*, *SacI*, *KpnI*, and *HindIII* cloning sit; Invitrogen) or the fusion pYESHisA, B, C (*XbaI*, *SphI*, *ShoI*, *NotI*, *BstXI*, *EcoRI*, *BamHI*, *SacI*, *KpnI*, and *HindIII* cloning site, N-terminal peptide purified with ProBond resin and cleaved with enterokinase; Invitrogen), to
5 mention just two, can be employed according to the invention.

Once a suitable host system and growth conditions are established, recombinant expression vectors can be propagated and prepared in quantity. As previously explained, the expression vectors which can be used include, but are not limited to, the following vectors or
10 their derivatives: human or animal viruses such as vaccinia virus or adenovirus; insect viruses such as baculovirus; yeast vectors; bacteriophage vectors (e.g., lambda), and plasmid and cosmid DNA vectors, to name but a few.

Vectors are introduced into the desired host cells by methods
15 known in the art, e.g., transfection, electroporation, microinjection, transduction, cell fusion, DEAE dextran, calcium phosphate precipitation, lipofection (lysosome fusion), use of a gene gun, or a DNA vector transporter (see, e.g., Wu et al., 1992; Wu and Wu, 1988; Canadian Patent Application No. 2,012,311, filed March 15, 1990).

20 A cell has been "transfected" by exogenous or heterologous DNA when such DNA has been introduced inside the cell. A cell has been "transformed" by exogenous or heterologous DNA when the transfected DNA effects a phenotypic change.

For introducing a vector in a cell host, explicit reference is made
25 to research carried out by the group of E. Wagner, relating to gene delivery by means of plasmid-polylysine complexes (Curiel et al., 1991; and Curiel et al., 1992). The plasmid-polylysine complex investigated upon exposition to certain cell lines showed at least some expression of the gene. Further, it was found that the expression efficiency increased
30 considerably due to the binding of transferrin to the plasmid-polylysine complex. Transferrin gives rise to close cell-complex contact with cells comprising transferrin receptors; it binds the entire complex to the transferrin receptor of cells. Subsequently, at least part of the entire complex was found to be incorporated in the cells investigated.

35

DA.
Le August 2 2000

Several different approaches have been developed for gene transfer. These include the use of viral based vectors (e.g., retroviruses, adenoviruses, and adeno-associated viruses) (Drumm, M. L. et al., Rosenfeld, M. A. et al., 1992; and Muzyczka, 1992), charge associating
5 the DNA with an asialorosomucoid/poly L-lysine complex (Wilson, J. M. et al. 1992), charge associating the DNA with cationic liposomes (Brigham, K. L. et al., 1993) and the use of cationic liposomes in association with a poly-L-lysine antibody complex (Trubetskoy, V. S. et al., 1993).

10

Compositions comprising vectors of the invention.

Although non-viral based transfection systems have not exhibited the efficiency of viral vectors, they have received significant
15 attention, in both in vitro and in vivo research, because of their theoretical safety when compared to viral vectors. Synthetic cationic molecules, have been reported which reportedly "coat" the nucleic acid through the interaction of the cationic sites on the transfection agent and the anionic sites on the nucleic acid. The positively charged coating
20 reportedly interacts with the negatively charged cell membrane to facilitate the passage of the nucleic acid through the cell membrane by non-specific endocytosis. (Schofield, 1995) These compounds have, however, exhibited considerable sensitivity to natural serum inhibition, which has probably limited their efficiency in vivo as gene transfection
25 agents. (Behr 1994)

A number of attempts have been made to improve the efficiency of lipid-like cationic transfection agents, some involving the use of polycationic molecules. For example, several transfection agents have been developed that contain the polycationic compound spermine
30 covalently attached to a lipid carrier. (Behr, 1994), discloses a lipopolyamine and shows it to be more efficient at transfecting cells than single charge molecules (albeit still less efficient than viral vectors). The agent reported by Behr was, however, toxic, and caused cell death.

A few such lipid delivery systems for transporting DNA, proteins, and other chemical materials across membrane boundaries have been synthesized by research groups and business entities. Most of the synthesis schemes are relatively complex and generate lipid based delivery systems having only limited transfection abilities. A need exists in the field of gene therapy for cationic lipid species that have a high biopolymer transport efficiency. It has been known for some time that a very limited number of certain quaternary ammonium derivatized (cationic) liposomes spontaneously associate with DNA, fuse with cell membranes, and deliver the DNA into the cytoplasm (as noted above, these species have been termed "cytofectins"). LIPOFECTIN™ represents a first generation of cationic liposome formulation development. LIPOFECTIN™ is composed of a 1:1 formulation of the quaternary ammonium containing compound DOTMA and dioleoylphosphatidylethanolamine sonicated into small unilamellar vesicles in water. Problems associated with LIPOFECTIN™ include non-metabolizable ether bonds, inhibition of protein kinase C activity, and direct cytotoxicity. In response to these problems, a number of other related compounds have been developed. The monoammonium compounds of the subject invention improve upon the capabilities of existing cationic liposomes and serve as a very efficient delivery system for biologically active chemicals.

Most preferred vectors of the invention.

25

Most preferred recombinant vectors according to the invention include pASΔΔ (figure 2), pACT11st (figure 3), pT18 (figure 4), pUT18C (figure 5), pT25 (figure 6), pKT25 (figure 7), pB5 (Figure 12) and pP6 (Figure 13) containing inserted therein a nucleic acid encoding a Selected Interacting Domain (SID®) polypeptide or a variant thereof as defined above.

The present invention is also directed to a vector usable in a two-hybrid method which consists of the vector pP6 which is shown in figure 13. As disclosed in example 1, the vector pP6 has been successfully used for preparing a collection of recombinant plasmids

35

DA.
Le August 2 2000

consisting of a genomic DNA library from the pathogenic strain H77 of the hepatitis C virus.

The invention also pertains to a vector usable in two-hybrid method which consists of the vector pB5. As disclosed in example 1, the
5 vector pB5 has been successfully used in a yeast two hybrid method as a bait plasmid.

RECOMBINANT CELL HOSTS

10 In one embodiment, a Selected Interacting Domain (SID®) polypeptide of the invention or a variant thereof is recombinantly produced in a desired host cell which has been transfected or transformed with a nucleic acid encoding said Selected Interacting Domain (SID®) polypeptide or with a recombinant vector as defined
15 above within which a nucleic acid encoding a Selected Interacting Domain (SID®) polypeptide of the invention is inserted.

Recombinant cell hosts are another aspect of the present invention.

20 Such cell hosts generally comprise at least one copy of a nucleic acid encoding a Selected Interacting Domain (SID®) polypeptide of the invention or a variant thereof

Preferred cells for expression purposes will be selected in function of the objective which is sought. For example, in the
25 embodiment wherein the production of a Selected Interacting Domain (SID®) polypeptide according to the invention in large quantities is sought, the nature of the host cell used for its production is relatively indifferent, provided that large amounts of Selected Interacting Domain (SID®) polypeptides of the invention are produced and that optional
30 further purification steps may be carried out easily.

However, in the embodiment wherein the Selected Interacting Domain (SID®) polypeptide is recombinantly produced within a host organism for the purpose of interfering with a specific protein-protein interaction, then the host organism is selected among the host

organisms which are suspected to produce naturally said polypeptide of interest.

Consequently, mammalian and typically human cells, as well as bacterial, yeast, fungal, insect, nematode and plant cells are cell hosts encompassed by the invention and which may be transfected either by a nucleic acid or a recombinant vector as defined above.

Examples of suitable recombinant host cells include VERO cells, HELA cells (e.g. ATCC N°CCL2), CHO cell-lines (e.g. ATCC N°CCL61) COS cells (e.g. COS-7 cells; COS cell referred to ATCC N°CRL1650), W138, BHK, HepG2, 3T3 (e.g. ATCC N°CRL6361), A549, PC12, K562 cells, 293 cells, Sf9 cells (e.g. ATCC N°CRL1711) and Cv1 cells (e.g. ATCC N°CCL70).

Other suitable host cells are usable according to the invention include prokaryotic host cells strains of *Escherichia coli* (e.g. strain DH5- α), of *Bacillus subtilis*, of *Salmonella typhimurium*, or strains of genera such as *Pseudomonas*, *Streptomyces* and *Staphylococcus*.

Further suitable host cells usable according to the invention include yeast cells such as those of *Saccharomyces*, typically *Saccharomyces cerevisiae*.

The invention also relates to a method for producing a SID® polypeptide as defined above, wherein said method comprises the steps of:

a) cultivating a cell host which has been transformed with a SID® nucleic acid of the invention or with a vector containing a SID® nucleic acid in an appropriate culture medium;

b) recovering the SID® recombinant polypeptide from the culture supernatant or from the cell lysate.

The SID® polypeptides or variant thereof thus recombinantly obtained may be purified, for example by high performance liquid chromatography, such as reverse phase and/or cationic exchange HPLC, as described by ROUGEOT et al. (1994). The reason to prefer this kind of peptide or protein purification is the lack of by-products found in the elution samples which renders the resultant purified protein more suitable for a therapeutic use.

35

DA.

Le August 2 2000

TWO-HYBRID METHODS OF THE INVENTION

a) Yeast two-hybrid methods

5 The invention also pertains to a yeast two-hybrid method for selecting a recombinant cell clone containing a vector comprising a nucleic acid insert encoding a prey polypeptide which binds with a SID® polypeptide of SEQ ID N°1 to 38 or a variant thereof, wherein said method comprises the steps of :

10 a) mating at least one first recombinant yeast cell clone of a collection of recombinant yeast cell clones transformed with a plasmid containing the prey polynucleotide to be assayed with a second aploid recombinant *Saccharomyces cerevisiae* cell clone transformed with a plasmid containing a bait polynucleotide encoding a SID® polypeptide of
15 the invention or a variant thereof;

 b) cultivating diploid cells obtained in step a) on a selective medium; and

 c) selecting recombinant cell clones which grow on said selective medium.

20 The yeast two-hybrid method above may further comprise the step of :

 d) characterizing the prey polynucleotide contained in each recombinant cell clone selected in step c).

25 Most preferably, such a yeast two-hybrid method may be performed by the one skilled in the art as it is disclosed in example 2 hereafter.

 According to the yeast two-hybrid method above, a SID® polypeptide of the invention or a variant thereof is used as a bait polypeptide.

30 In a preferred embodiment of the yeast two-hybrid method described above, the prey polynucleotide is a DNA fragment from the genome of a pathogenic strain of the hepatitis C virus (HCV) ranging from about 150 to about 600 nucleotides in length and which is inserted in a vector which is contained in one recombinant clone of a collection of
35 recombinant cell clones.

b) Bacterial two-hybrid method

5 A bacterial two-hybrid method of the invention may be performed by the one skilled in the art according to the teachings of KARIMOVA et al. (1998).

The first step of selecting a collection of nucleic acids encoding polypeptides which binds specifically to the bait polypeptide may also be carried out through a bacterial two-hybrid system.

10 According to such bacterial two-hybrid system, bacterial cell clones, preferably *Escherichia coli* cells, are transformed with a plasmid containing a bait polynucleotide encoding a bait polypeptide.

Then, plasmids containing a DNA insert are provided by rescuing the plasmids obtained from the collection of yeast clones containing the genomic DNA or cDNA library which are described in the previous section entitled "Yeast two-hybrid system". For example, the plasmid rescue may be carried out according to the following steps:

(i) extracting plasmid DNA contained in the collection of yeast clones obtained as disclosed in the previous section, by using a conventional DNA extraction buffer and a phenol: chloroform: isoamyl alcohol (25:24:1) before centrifuging;

(ii) transferring a desired volume of the supernatant obtained at the end of step (i) to a sterile Eppendorf tube and add a precipitation buffer (ethanol/NH₄Ac) before centrifuging and resuspending the pellet after washing in ethanol;

(iii) transforming *Escherichia coli* cells (e.g. *Escherichia coli* cells of strain NC 1066) which have been rendered electrocompetent with a desired volume (e.g. 1 µl) of the yeast plasmid DNA extract obtained at step (ii) by electroporation;

30 (iv) collecting the transformed *Escherichia coli* cells.

Alternatively, a collection of *Escherichia coli* cell clones containing a collection of HCV genomic DNA inserts may be obtained by constructing the DNA library directly in the bacterial cell, such as disclosed in Flajolet et al. (2000).

DA.

Le August 2 2000

Then, the bacterial recombinant cells which have been transformed both with a plasmid containing a bait polynucleotide encoding a bait polypeptide and a plasmid containing a prey polynucleotide encoding a prey polypeptide is cultivated on a selective medium.

Then, recombinant cell clones capable of growing on said selective medium are selected and the DNA inserts of the plasmids containing therein are sequenced.

By bacterial two-hybrid system is generally intended a method that usually makes use of at least one reporter gene, the transcription of which is activated when a prey polypeptide and a bait polypeptide produced by the recombinant cell due to the triggering of the transcription of said at least one reporter gene when both the specific domain contained in one prey polypeptide and the complementary domain contained in the bait polypeptide are binding one to the other.

The invention further pertains to a bacterial two-hybrid method for identifying a recombinant cell clone containing a prey polynucleotide encoding a prey polypeptide which binds with a SID® polypeptide of SEQ ID N°1 to 38 or a variant thereof, wherein said method comprises the steps of :

- a) transforming bacterial cell clones with a plasmid containing a SID® polynucleotide encoding a SID® polypeptide of the invention or a variant thereof;
 - b) rescuing prey plasmids containing prey polynucleotides wherein each prey polynucleotide is a DNA fragment from the genome of a desired organism and wherein each prey plasmid is contained in one recombinant yeast cell clone of a collection of recombinant yeast cell clones;
 - c) transforming the recombinant bacterial cell clones obtained in step a) with the plasmids rescued in step b);
 - d) cultivating bacterial recombinant cells obtained in step c) on a selective medium;
- and

e) selecting recombinant cell clones which grow on said selective medium.

The bacterial two-hybrid system described above may further comprise the step of f) characterizing the prey polynucleotide contained
5 in each recombinant cell clone selected at step e).

In one preferred embodiment of the yeast or bacterial two-hybrid methods described above, the prey polypeptide is a human polypeptide expressed by a mammal which is infected by the Hepatitis C virus, like human and monkeys, typically chimpanzees.

10 Generally, the yeast two-hybrid method or the bacterial two-hybrid method as disclosed herein may be performed with prey polypeptides of any origin, either of viral, fungal, bacterial or mammal origin, i.e. either of prokaryotic or eukaryotic origin.

15

In a second preferred embodiment of the two-hybrid methods above, the prey polypeptide is an HCV polypeptide.

Most preferably, the prey polypeptide is encoded by a strain of
20 the hepatitis C virus which is pathogenic for human, such as strain H77.

SETS OF NUCLEIC ACIDS AND SETS OF POLYPEPTIDES OF THE INVENTION

25 In yet another aspect, the present invention relates to a set of two nucleic acids consisting of:

- i) a first nucleic acid encoding a SID® polypeptide of SEQ ID N°1 to 39 of the invention or a variant thereof; and
- 30 ii) a second nucleic acid encoding a prey polypeptide which binds specifically with a SID® polypeptide defined in i).

In still a further aspect, the invention is also directed to a set of two polypeptides consisting of :

- 35 i) a first polypeptide consisting of a SID® polypeptide of SEQ ID N°1 to 39 of the invention or a variant thereof; and

DA.

Le August 2 2000

ii) a second polypeptide which binds specifically with the first polypeptide.

The invention further relates to a complex formed between :

- 5 i) a first polypeptide consisting of a SID® polypeptide of SEQ ID N°1 to N°38 of the invention; and
ii) a second polypeptide which binds specifically with the first polypeptide.

10 The invention also relates to a protein-protein interaction wherein the two interacting proteins consist of a set of two polypeptides as defined above.

In a preferred embodiment, the invention relates to the protein-protein interactions wherein the sets of two polypeptides consist of a SID® polypeptide of SEQ ID N°1 to 38 and an HCV polypeptide.

15 When several reiterations of the two-hybrid method are performed and thus common SID® polypeptide and prey polypeptides are selected, a map of all the interactions between these polypeptides may be designed, that take into account of the known and/or suspected biological function of each of the interacting polypeptides.

20 Table 1 illustrates protein-protein interaction between the SID® polypeptides of SEQ ID N°1 to 38 and polypeptides of SEQ ID N°77 to 113 which are encoded by the genome of strain H77 of the hepatitis C virus which is pathogenic for a mammal, like human or chimpanzee.

25 Thus, the data presented in table 1 disclose particular sets of nucleic acids as well as particular sets of polypeptides which are encompassed by the present invention.

30 For example, table 1 discloses that the nucleic acid of SEQ ID N°39 encodes the SID® polypeptide of SEQ ID N°1 which contains exclusively (100 %) an aminoacid sequence from the Core protein of HCV strain H77.

The nucleic acid of SEQ ID N°39 starts at the nucleotide in position 446 and ends at the nucleotide in position 600 of the HCV genome which is described by YANAGI et al. (1997).

35 Table 1 also discloses that the SID® polypeptide of SEQ ID N°1 is part of a set of polypeptides of the invention, wherein the second

polypeptide of said set of polypeptides consists of the polypeptide of SEQ ID N°77 which is encoded by the nucleic acid sequence of SEQ ID N°114, which nucleic acid sequence has 87% of its sequence which is derived from the region of the H77 strain HCV DNA encoding the Core protein.

Thus , a particular set of polypeptides according to the invention consists of:

- i) the polypeptide of SEQ ID N°1; and
- ii) the polypeptide of SEQ ID N°77.

The same reasoning apply for every set of polypeptides disclosed in table 1, which are expressly part of the present invention.

Similarly, a particular set of nucleic acids according to the invention consists of :

- (i) the nucleic acid of SEQ ID N°39; and
- (ii) the nucleic acid of SEQ ID N°114.

The same reasoning apply for every set of nucleic acids disclosed in table 1, which are expressly part of the present invention.

Thus, particular sets of two polypeptides of the invention are respectively SEQ ID N°77/SEQ ID N°1; SEQ ID N°78/SEQ ID N°2; SEQ ID N°78/SEQ ID N°3; SEQ ID N°79/SEQ ID N°4; SEQ ID N°80/SEQ ID N°5; SEQ ID N°81/SEQ ID N°6; SEQ ID N°82/SEQ ID N°7; SEQ ID N°83/SEQ ID N°8; SEQ ID N°84/SEQ ID N°9; SEQ ID N°85/SEQ ID N°10; SEQ ID N°86/SEQ ID N°11; SEQ ID N°87/SEQ ID N°12; SEQ ID N°88/SEQ ID N°13; SEQ ID N°89/SEQ ID N°14; SEQ ID N°90/SEQ ID N°15; SEQ ID N°91/SEQ ID N°16; SEQ ID N°92/SEQ ID N°17; SEQ ID N°93/SEQ ID N°18; SEQ ID N°94/SEQ ID N°19; SEQ ID N°95/SEQ ID N°20; SEQ ID N°96/SEQ ID N°21; SEQ ID N°97/SEQ ID N°22; SEQ ID N°98/SEQ ID N°23; SEQ ID N°99/SEQ ID N°24; SEQ ID N°100/SEQ ID N°25. SEQ ID N°101/SEQ ID N°26. SEQ ID N°102/SEQ ID N°27; SEQ ID N°103/SEQ ID N°28. SEQ ID N°104/SEQ ID N°29; SEQ ID N°105/SEQ ID N°30; SEQ ID N°106/SEQ ID N°31; SEQ ID N°107/SEQ ID N°32; SEQ ID N°108/SEQ ID N°33; SEQ ID N°109/SEQ ID N°34; SEQ ID N°110/SEQ ID N°35; SEQ ID N°111/SEQ ID N°36; SEQ ID N°112/SEQ ID N°37; and SEQ ID N°113/SEQ ID N°38.

DA.

Le August 2 2000

Similarly, particular sets of two nucleic acids according to the invention are respectively: SEQ ID N°114/SEQ ID N°39; SEQ ID N°115/SEQ ID N°40; SEQ ID N°115/SEQ ID N°41; SEQ ID N°116/SEQ ID N°42; SEQ ID N°117/SEQ ID N°43; SEQ ID N°118/SEQ ID N°44; 5 SEQ ID N°119/SEQ ID N°45; SEQ ID N°120/SEQ ID N°46; SEQ ID N°121/SEQ ID N°47; SEQ ID N°122/SEQ ID N°48; SEQ ID N°123/SEQ ID N°49; SEQ ID N°124/SEQ ID N°50; SEQ ID N°125/SEQ ID N°51; SEQ ID N°126/SEQ ID N°52; SEQ ID N°127/SEQ ID N°53; SEQ ID N°128/SEQ ID N°54; SEQ ID N°129/SEQ ID N°55; SEQ ID N°130/SEQ ID N°56; SEQ ID N°131/SEQ ID N°57; SEQ ID N°132/SEQ ID N°58; 10 SEQ ID N°133/SEQ ID N°59; SEQ ID N°134/SEQ ID N°60; SEQ ID N°135/SEQ ID N°61; SEQ ID N°136/SEQ ID N°62; SEQ ID N°137/SEQ ID N°63; SEQ ID N°138/SEQ ID N°64; SEQ ID N°139/SEQ ID N°65; SEQ ID N°140/SEQ ID N°66; SEQ ID N°141/SEQ ID N°67; SEQ ID N°142/SEQ ID N°68; SEQ ID N°143/SEQ ID N°69; SEQ ID N°144/SEQ ID N°70. SEQ ID N°145/SEQ ID N°71; SEQ ID N°146/SEQ ID N°72. 15 SEQ ID N°147/SEQ ID N°73; SEQ ID N°148/SEQ ID N°74; SEQ ID N°149/SEQ ID N°75 and SEQ ID N°150/SEQ ID N°76.

20 The protein-protein interactions disclosed in table 1 allows the design of a map of interactions between various polypeptides encoded by the genome of the H77 strain of HCV.

In such a Protein Interaction Map (PIM®) wherein each SID® polypeptide is linked to the bait polypeptide onto which it specifically 25 binds, for example by an arrow.

Such a Protein Interaction Map (PIM®) may help the one skilled in the art to decipher a whole metabolical and/or physiological pathway that is functionally active within a pathogenic strain of HCV. Protein Interaction Map and computable version of PIM® are part of the present 30 invention.

Therefore, in still another aspect, the present invention is directed to a computable readable medium (such as floppy disk, CD-ROM and all electronic or magnetic format which can be read by a computer) having stored thereon protein-protein interactions according to

the invention, preferably stored in a form of a Protein Interaction MAP, as shown, for example, in FROMONT-RACINE et al. (1997).

In a preferred embodiment, the invention comprises a computable readable medium as defined above, wherein the protein-protein interactions stored thereon are linked to annotated data base, for example through Internet.

In another preferred embodiment, the invention comprises a data bank containing the protein-protein interactions stored thereon, said data bank being available on a world-wide web site.

10

METHODS FOR SELECTING INHIBITORS OF PROTEIN-PROTEIN INTERACTIONS OF THE INVENTION

The transformed host cells as described above can also be used as models so as to study the interactions between a SID® polypeptide of the invention and its binding partner polypeptide, or between a SID® polypeptide of the invention and chemical or protein compounds which inhibit the binding between said SID® polypeptide and its binding partner polypeptide.

Example of a SID® polypeptide and its binding partner polypeptides are typically the sets of polypeptides of the invention which are described above.

In particular, the transformed host cells of the invention may be used for the selection of molecules which interact with a SID® polypeptide as described herein, as cofactor or as inhibitor, in particular a competitive inhibitor, or alternatively having an agonist or antagonist activity on the protein-protein interaction wherein said SID® polypeptide is involved. Preferably, the said transformed host cells will be used as a model allowing, in particular, the selection of products which make it possible to prevent and/or to treat pathologies induced by the hepatitis C virus.

Consequently, the invention also consists of a method for selecting a molecule which inhibits the protein-protein interaction of a set of two polypeptides as defined above, wherein said method comprises the steps of :

DA.

Le August 2 2000

a) cultivating a recombinant host cell containing a reporter gene the expression of which is toxic for said recombinant host cell, said host cell being transformed with two vectors wherein:

5 i) the first vector contains a nucleic acid comprising a polynucleotide encoding a first hybrid polypeptide containing one of said two-polypeptides and a DNA binding domain;

10 ii) the second vector contains a nucleic acid comprising a polynucleotide encoding a second hybrid polypeptide containing the second of said two polypeptides and an activating domain capable of activating said toxic reporter gene when the first and the second hybrid polypeptides are interacting;

on a selective medium containing the molecule to be tested and allowing the growth of said recombinant host cell when the toxic reporter gene is not activated; and

15 b) selecting the molecule which inhibits the growth of the recombinant host cell defined in step a).

The invention is also directed to a method for selecting a molecule which inhibits the protein-protein interaction of a set of two polypeptides as defined above, wherein said method comprises the steps of :

20 a) cultivating a recombinant host cell containing a reporter gene the expression of which is toxic for said recombinant host cell, said host cell being transformed with two vectors wherein:

25 i) the first vector contains a nucleic acid comprising a polynucleotide encoding a first hybrid polypeptide containing one of said two polypeptides and the first domain of an enzyme;

30 ii) the second vector contains a nucleic acid comprising a polynucleotide encoding a second hybrid polypeptide containing the second of said two polypeptides and the second part of said enzyme capable of activating said toxic reporter gene when the first and the second hybrid polypeptides are interacting, said interaction recovering the catalytic activity of the enzyme;

35 on a selective medium containing the molecule to be tested and allowing the growth of said recombinant host cell when the toxic gene is not activated; and

b) selecting the molecule which inhibits the growth of the recombinant host cell defined in step a).

In a preferred embodiment, said toxic reporter gene that can be used for negative selection is URA3, CYH1 or CYH2 gene.

5 For example, a method for the screening of a molecule which inhibits the interaction between a SID® polypeptide of the invention with its binding protein counterpart may comprise the following steps:

- transform a permeabilized yeast cell with two vectors, respectively a first vector containing a SID® nucleic acid of the invention and a second vector containing a prey nucleic acid as defined in the present specification;

10 - plate on top agar the transformed permeabilized yeast cells above on square boxes;

- apply by spotting the candidate inhibitor molecules to test on top agar as soon as it is solidified;

15 - incubates, for example, overnight at 30°C, and

- select the inhibitor compounds that allow the growth of the transformed yeast cells.

The invention also provides for a kit for the screening of a molecule which inhibits the protein-protein interaction of a set of two polypeptides as defined above, wherein said kit comprises a recombinant host cell containing a reporter gene the expression of which is toxic for said recombinant host cell, said host cell being transformed with two vectors wherein:

25 i) the first vector contains a nucleic acid comprising a polynucleotide encoding a first hybrid polypeptide containing one of said two polypeptides and a DNA binding domain;

ii) the second vector contains a nucleic acid comprising a polynucleotide encoding a second hybrid polypeptide containing the second of said two polypeptides and an activating domain capable of activating said toxic reporter gene when the first and the second hybrid polypeptides are interacting.

30 Another object of the invention consists of a kit for the screening of a molecule which inhibits the protein-protein interaction of a set of two polypeptides as defined above, wherein said kit comprises a

DA.

Le August 2 2000

recombinant host cell containing a reporter gene the expression of which is toxic for said recombinant host cell, said host cell being transformed with two plasmids wherein:

5 i) the first vector contains a nucleic acid comprising a polynucleotide encoding a first hybrid polypeptide containing one of said two polypeptides and the first domain of a protein;

10 ii) the second vector contains a nucleic acid comprising a polynucleotide encoding a second hybrid polypeptide containing the second of said two polypeptides and the second part of said protein capable of activating said toxic reporter gene when the first and the second hybrid polypeptides are interacting, said interaction recovering the activity of the protein. In the selection methods above, the transcription or activating domain and the DNA-binding domain may be derived from Gal4 and LexA respectively.

15 In the embodiment wherein the first domain is a first part of an enzyme and a complementary domain is a second part of the same enzyme, and wherein the proximity of the two parts of the enzyme restores the enzyme activity and activates a reporter gene, the two parts of the enzymes are most preferably the T25 and T18 polypeptides that
20 form the catalytic domain of the *Bordetella pertussis adenylate cyclase*.

As an illustrative embodiment, the reporter gene is chosen among the group consisting of a nutritional gene or also a gene the expression of which is visualised by colorimetry such as His3, LacZ or both LacZ and His3.

25

MARKER COMPOUNDS OF THE INVENTION

30 The Selected Interacting Domain (SID®) polypeptides of SEQ ID N°1 to 38 of the invention and variants thereof defined in the present specification, and which bind specifically to a polypeptide of interest (e.g. a bait polypeptide), are useful as reagents for detecting, labelling, targeting or purifying specifically a polypeptide of interest, typically a polypeptide encoded by HCV, within a sample, since the SID® polypeptides possess properties that have never been reached using

conventional detection compounds, such as those of an antibody or an antibody fragment.

Firstly, the SID® polypeptides of the invention possess a high specificity of binding to the polypeptide of interest, since a SID® polypeptide consists of a portion of a larger polypeptide which binds in a highly specific manner to the polypeptide of interest in the natural environment within the eukaryotic cell infected by the Hepatitis C virus.

Secondly, the SID® polypeptide generally has a low molecular weight, generally from 3 kDa, and are thus easy to produce, on the one hand, and, on the other hand, can be easily introduced within a cell when the detection of the localisation or of the expression of the polypeptide of interest is sought. Moreover, the small size of a SID® polypeptide allows its passage through inner cell barriers such as the nucleus membrane, or the membranes surrounding the different cell organites.

Thus, a first object of the invention consists of a marker compound wherein said compound comprises :

- a) a Selected Interacting Domain (SID®) polypeptide of the invention or a variant thereof that binds specifically to the polypeptide of interest; and
- b) a detectable molecule bound thereto.

Such a marker compound is primarily useful for detecting, labelling or targeting a polypeptide of interest, for example a polypeptide of interest contained in a sample.

A detectable molecule according to the invention comprises, or alternatively consists of, any molecule which produces or can be induced to produce a signal. The detectable molecule can be a member of the signal producing system that includes the signal producing means .

The detectable molecule may be isotopic or non-isotopic. By way of example and not limitation, the detectable molecule can be part of a catalytic reaction system such as enzymes, enzyme fragments, enzyme substrates, enzyme inhibitors, co-enzymes, or catalysts. Part of a chromogen system such as fluorophores, dyes, chemilumescers, luminescers, or sensitizers. A dispersible particle that can be non-magnetic or magnetic, a solid support, a liposome, a ligand, a receptor, a hapten radioactive isotope, and so forth.

DA.

Le August 2 2000

It must be generally understood that the whole embodiments disclosed in the present specification involving a Selected Interacting Domain (SID®) polypeptide is straightfully applied also to any variant thereof.

5

Fluorescent detectable molecules

In one aspect of the marker compound according to the invention, the detectable molecule consists of a fluorescent molecule. Fluorescent moieties which are frequently used as labels are for example
10 those described by Ichinose et al. (1991). Other fluorescent detectable molecules are fluorescing isothiocyanate (FITC) such as described by Shattil et al. (1987) or by Goding et al. (1986). The fluorescent detectable molecule may also comprise a phycoerythrin as taught by Goding et al. (1986), and Shattil et al. (1985). Other examples of
15 fluorescent detectable molecules suitable for use as labels of a marker compound according to the invention are rhodamine isothiocyanate, dansyl chloride and XRITC.

Another fluorescent detectable molecule consists of the green
20 fluorescent protein (GFP) of the jelly fish *Aequorea victoria*, and their numerous fluorescent protein derivatives.

The one skilled in the art may advantageously refer to the articles of CHALFIE et al. (1994) and of HEIM et al. (1994) which discloses the uses of GFP for the study of gene expression and protein
25 localisation. The one skilled in the art may also refer to the article of Rizzuto et al. (1995) , which discusses the use of wild-type GFP as a tool for visualising subcellular organelles in cells, to the article of KAETHER and GERDES (1995), which reports the visualisation of protein transport along the secretary passway using wild-type GFP, the article of HU and
30 CHENG (1995), which relates to the expression of GFP in plant cells and also to the article of Davis et al. (1995) which discloses the GFP expression in drosophilia embryos. For the use of several fluorescent variants of GFP, the one skilled in the art may refer to the article of Delagrave et al. (1995), as well as to the article of Heim et al. (1995).
35 DNA encoding GFP is available commercially, for example from

Clontech in Palo Alto, California, USA. The one skilled in the art may use also humanized GFP genes such as those described in the US Patent N°6,020,192 and also the GFP protein disclosed in the US Patent N°5,941,084.

5 Another fluorescent protein that may be used in a marker compound according to the invention consists of the yellow fluorescent protein (YFP).

 A further suitable luminescent protein consists of the luciferase protein.

10

Detectable molecules exhibiting a catalytic activity

 In another embodiment of a detectable molecule included in a marker compound according to the invention, said detectable molecule is
15 endowed with a catalytic activity and may thus consists of enzymes and catalytically active enzyme fragments. Some enzymatic labels are described in US Patent N°3,654,090. Such enzymes may be for example horse radish peroxidase (HRP), alkaline phosphatase or glutathione peroxidase which are well known from the one skilled in the art.

20

 Enzymes, enzyme fragments, enzyme inhibitors, enzyme substrates, and other components of enzyme reaction systems can be used as detectable molecules. Where any of these components is used as a detectable molecule, a chemical reaction involving one of the
25 components is part of the signal producing system.

 Coupled catalysts can also involve an enzyme with a non-enzymatic catalyst. The enzyme can produce a reactant, which undergoes a reaction catalysed by the non-enzymatic catalyst or the non-enzymatic catalyst may produce a substrate (including co-enzymes)
30 for the enzyme. The one skilled in the art may advantageously refer to the US Patent N°4,160 645 which disclose a white variety of non enzymatic catalysts, which may be employed, the appropriate portions of which are incorporated therein by reference.

 The enzyme or co-enzyme employed provides the desired
35 amplification by producing a product, which absorbs light, e.g., a dye, or

DA.

Le August 2 2000

emits lights upon irradiation, e.g., a fluorescer. Alternatively, the catalytic reaction can lead to direct light emission, e.g., chemiluminescence. A large number of enzymes and co-enzymes for providing such products are described in the US Patents N°4,275,149, columns 19 to 23 and
5 N°4,318,980, columns 10 to 14 which disclosures are incorporated herein by reference.

A number of enzyme combinations are set forth in US Patent N°4,275,149, columns 23 to 28 which disclosures are incorporated herein by reference.

10 When a single enzyme is used as the detectable molecule, or alternatively as comprised in the detectable molecule, such enzymes may find use are hydrolases, transferases, lyases, isomerases, ligases or synthetases and oxydoreductases.

Alternatively, luciferases may be used such as firefly luciferase
15 and bacterial luciferase.

Primarily, the enzymes of choice, based on the I.U.B. classification are: (i) class 1. Oxydoreductases and (ii) class 3. Hydrolases. Most preferred oxydoreductases are (i) dehydrogenases of class 1.1, more particularly 1.1.1, 1.1.3. and 1.1.99 and (ii) peroxydases
20 in class 1.11. of the hydrolases, particularly class 3.1., more particularly 3.1.3 and class 3.2, more particularly 3.2.1. are preferred.

Illustrative dehydrogenases include malate dehydrogenase, glucose-6-phosphate dehydrogenase and lactate dehydrogenase. Of the oxydases, glucose oxydases is exemplary. Of the peroxydases, horse
25 radish peroxydase is illustrative. Of the hydrolases, alkaline phosphatases, β -glucosydase and lysozyme are illustrative.

Chemiluminescent detectable molecules

30 The detectable molecule comprised within the marker compound according to the invention may also consist in a chemiluminescent moiety. The chemiluminescent source involves a compound, which becomes electronically excited by a chemical reaction and may emit light which serves at as the detectable signal or donates
35 energy to a fluorescent acceptor.

A diverse number of families of compounds have been found to provide chemiluminescent under a variety of conditions. When family of compounds is 2,3-dihydro-1,4-phtalazinedinone. The most utilised compound is luminol, which is the 5-amino analogue of the compound
5 above. Other members of the family include the 5-amino-6,7,8-trimethoxy-and the dimethylamine-[ca]benzo analogue. These compounds can be made to luminance with alkaline hydrogen peroxyde or calcium hypochlorite and base.

Another family of compounds is the 2,4,5-triphenylimidazoles,
10 with lophine as the common name for the parent product. Chemiluminescent analogues include para-dimethylamino- and para-methoxy-substituents. Chemiluminescents may also be obtained with geridinium esters, dioxetanes and oxalates, usually oxalyl active esters, e.g., p-nitrophenyl and a peroxide, e.g., hydrogen peroxide, under basic
15 conditions. Alternatively, luciferins may be used in conjunction with luciferase or lucigenins.

Radioactive detectable molecules

20 In a further embodiment of a detectable molecule comprised in a marker compound according to the invention, said detectable molecule is radio-actively labelled such as with [^3H], [^{32}P], [and [^{125}I]].

Colloïdal metal detectable molecules

25 In still a further embodiment, the detectable molecule comprised in a marker compound according to the invention may include a colloïdal metal particle. Colloïdal metals have been employed in immuno assays previously. Mostly, they consisted of either colloïdal iron
30 or gold. The one skilled in the art may advantageously refer to the articles of Horisberger (1981) and Martin et al. (1990). In other case, the metals are chosen for their colour, i.e., their presence is determined by their colour or electron density under an electron microscope. Both the colour and electron density are directly proportional to the mass of the
35 metal colloïd.

DA.

Le August 2 2000

STRUCTURE OF THE MARKER COMPOUNDS OF THE INVENTION

5 In a first preferred embodiment of a marker compound of the invention, the detectable molecule is covalently bound to the Selected Interacting Domain (SID®) polypeptide of SEQ ID N°1 to SEQ ID N°38 or a variant thereof.

10 According to this specific embodiment, detectable molecules comprising fluorescent proteins such as GFP and YFP, enzymes or enzyme fragments such as alkaline phosphatase, glutathione peroxidase and horse radish peroxidase, chemiluminescent molecules, radioactive labels or colloidal metal particles will be preferred.

15 General methods that may be used by the one skilled in the art for covalently binding the detectable molecules to the Selected Interacting Domain (SID®) polypeptide are described in the numerous bibliographic references related to the preparation of the antibody conjugates used for carrying out immunoassays.

20 In a second preferred embodiment of a marker compound according to the invention, the detectable molecule is non-covalently bound to the Selected Interacting Domain (SID®) polypeptide or a variant thereof.

25 In a first preferred aspect of this second preferred embodiment, the detectable molecule consists of an antibody directed specifically against the Selected Interacting Domain (SID®) polypeptide or a variant thereof.

The antibodies directed specifically against the Selected Interacting Domain (SID®) polypeptide or a variant thereof may be indifferently radioactivity or non radioactivity labelled.

30 **NUCLEIC ACIDS ENCODING A MARKER COMPOUND OF THE INVENTION.**

The present invention also relates to a nucleic acid encoding a marker compound as defined above.

Most preferred nucleic acids encompassed by the invention include polynucleotides that encode a marker compound wherein the Selected Interacting Domain (SID®) polypeptide of SEQ ID N°1 to 38 or a variant thereof is covalently bound to the detectable molecule and wherein the detectable molecule consists itself of a polypeptide.

Most preferred nucleic acids are those of SEQ ID N°39 to 76.

In a first preferred embodiment of a nucleic acid according to the invention, said nucleic acid encodes for a Selected Interacting Domain (SID®) polypeptide which is fused to a fluorescent protein, such as GFP and YFP.

In a second preferred embodiment of a nucleic acid according to the invention, said nucleic acid encodes for a Selected Interacting Domain (SID®) polypeptide which is fused to a polypeptide endowed with a catalytic activity, such as an enzyme or an enzymatically active enzyme fragment, like alkaline phosphatase, glutathione peroxydase and horse radish peroxydase.

In a preferred embodiment, a nucleic acid encoding a marker compound of the invention comprises a DNA coding sequence which is transcribed and translated into said marker compound in a cell *in vitro* or *in vivo* when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a start codon and a translation stop codon. A coding sequence can include, but is not limited to:

- prokaryotic sequences, for example when the Selected Interacting Domain (SID®) nucleic acid and the nucleic acid fused thereto which encodes the detectable molecule are of prokaryotic origin;
- prokaryotic and eukaryotic sequences, for example the nucleic acid encoding the detectable molecule originates from an eukaryotic host organism.

If the coding sequence is intended for expression in an eukaryotic cell, a polyadenylation signal and transcription termination sequence will usually be located 3' to the coding sequence.

In a most preferred embodiment of a nucleic acid sequence according to the invention, said nucleic acid sequence include a

DA.

Le August 2 2000

regulatory region which is functional in the host organism within which the expression of said nucleic acid sequence is sought, wherein said regulatory region comprises a promoter sequence.

5 "Regulatory region" means a nucleic acid sequence which regulates the expression of a nucleic acid. A regulatory region may include sequences which are naturally responsible for expressing a particular nucleic acid (a homologous region), or may include sequences of a different origin (responsible for expressing different proteins or even
10 synthetic proteins). In particular, the sequences can be sequences of eukaryotic or viral genes or derived sequences which stimulate or repress transcription of a gene in a specific or non-specific manner and in an inducible or non-inducible manner. Regulatory regions include origins of replication, RNA splice sites, enhancers, transcriptional
15 termination sequences, signal sequences which direct the polypeptide into the secretory pathways of the target cell, and promoters.

A "promoter sequence" is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. For purposes of defining the
20 present invention, the promoter sequence is bounded at its 3' terminus by the transcription initiation site and extends upstream (5' direction) to include the minimum number of bases or elements necessary to initiate transcription at levels detectable above background. Within the promoter sequence will be found a transcription initiation site (conveniently defined
25 for example, by mapping with nuclease S1), as well as protein binding domains (consensus sequences) responsible for the binding of RNA polymerase.

A coding sequence is "under the control" of transcriptional and translational control sequences in a cell when RNA polymerase
30 transcribes the coding sequence into mRNA, which is then trans-RNA spliced and translated into the protein encoded by the coding sequence.

Most preferred vectors for the expression of a marker compound of the invention.

Most preferred recombinant vectors for expressing a marker compound of the invention include pASΔΔ (figure 2), pACT11st (figure 3), pT18 (figure 4), pUT18C (figure 5), pT25 (figure 6), pKT25 (figure 7), pB5 (Figure 12) and pP6 (Figure 13) containing inserted therein a nucleic acid encoding a Selected Interacting Domain (SID®) polypeptide as defined above or a variant thereof.

The invention also pertains to recombinant host cells transformed with a vector expressing a marker compound as defined above, more particularly a vector comprising inserted therein a nucleic acid encoding said marker compound, which is operably linked to suitable regulation signals which are functional in the host cell wherein its expression is sought.

Preferred cells for expression purposes will be selected in function of the objective which is sought. For example, in the embodiment wherein the production of a marker compound according to the invention in large quantities is sought, the nature of the cell host used for its production is relatively indifferent, provided that large amounts of Selected Interacting Domain (SID®) polypeptides or marker compounds of the invention are produced and that optional further purification steps may be carried out easily.

However, in the embodiment wherein the marker compound is recombinantly produced within a host organism for the purpose of qualitative or quantitative analysis of the polypeptide of interest onto which said marker compound specifically binds, then the host organism is selected among the host organisms which are suspected to produce naturally said polypeptide of interest.

Consequently, mammalian and human cells, as well as bacterial, yeast, fungal, insect, nematode and plant cells are cell host encompassed by the invention and which may be transfected either by a nucleic acid or a recombinant vector as defined above.

DETECTION METHODS OF THE INVENTION

DA.
Le August 2 2000

The present invention further relates to the use of a Selected Interacting (SID®) polypeptide of SEQ ID N°1 to 38 or a variant thereof as well as a nucleic acid encoding it for detection purposes such as nucleic acids of SEQ ID N°39 to 76. It is herein reminded that a Selected
5 Interacting Domain (SID®) polypeptide is determined according to the ability of such a (SID®) polypeptide to bind in a highly specific manner to a given (e.g. bait) polypeptide of interest, since the aminoacid sequence of a SID® polypeptide is encoded by a nucleic acid, the nucleotide
10 sequence of which consists of the polynucleotide sequence which is common to a collection of nucleic acid sequences encoding prey polypeptides that have been selected for their specific binding properties to a (bait) polypeptide of interest, such as explained above in the section entitled "SELECTED INTERACTING DOMAIN (SID®)
POLYPEPTIDES".

15 The specific properties of a Selected Interacting Domain (SID®) polypeptide for binding to a given polypeptide of interest, either a viral, yeast, fungal, bacterial, insect, plant or mammal polypeptide, including a polypeptide of human origin, allow its use as a specific ligand for said polypeptide of interest of which the detection is sought.

20 Therefore, the use of a Selected Interacting Domain (SID®) in any detection method known in the art and which makes use of the ability of a detection ligand to bind specifically to a molecule of interest, most preferably a polypeptide of interest, fall under the scope of the present invention.

25 Detection methods that make use of the recognition of a molecule of interest, most preferably a polypeptide of interest, by a detection ligand are well known in the art and are primarily illustrated by the abundant literature that relate to immunoassays, which is incorporated herein by reference in its entirety.

30 The one skilled in the art may particularly refer to the book of Maggio (1980) (Heterogeneous assays), the US Patent N°3,817,837 (homogeneous Immunoassays), US Patent N° 3,993,345 (Immunofluorescence methods), US Patent N°4,233,402 (enzyme channelling techniques), US Patent N°3,817,837 (Enzyme multiplied
35 immunoassay technique), US Patent N°4,366,241 and European Patent

Application N°EP-A 0 143 574 (Migration type assays), US Patent N°5,202,006, US Patent N°5,120,413 and US Patent N°5,145,567 (Immunofixation electrophoresis, mmunoelectrophoresis), the article of Aguzzi et al. (1977), the article of White et al. (1986), the article of Merlini
5 et al. (1983), the US Patent n°5,228,960 (Immunosubstraction electrophoresis), the articles of Chen et al. (1991), Nielsen et al. (1991) and the US Patent n° 5,120,413 (Capillary electrophoresis).

Acellular detection method of the invention.

10

A first detection method of the invention consists of a method for detecting a polypeptide of interest within a sample, wherein said method comprises the steps of:

a) contacting a marker compound or a plurality of marker
15 compounds according to the invention with the sample which is suspected to contain the polypeptide of interest the detection of which is sought;

b) detecting the complexes formed between said marker compound or said plurality of marker compounds and said polypeptide of
20 interest.

The sample which is assayed for the presence of the polypeptide of interest the detection of which is sought may be of any nature , including every sample that may be used for carrying out an immunoassay.

25 In a first aspect, the sample may be any biological fluid, such as blood or blood separation products (e.g. serum, plasma, buffy coat), urine, saliva, tears.

In a second aspect, the sample may be any isolated biological tissue sample, including tissue sections previously fixed for purposes of
30 histological studies.

In a third aspect, the sample may be a culture supernatant of a cell culture and a cell lysate of cultured cells.

In a first preferred embodiment of the first detection method of the invention described above, the detection step b) consists of the
35 measure of the fluorescence signal intrinsically emitted by the detectable

DA.

Le August 2 2000

molecule. It may for example be taken the advantage of SID® polypeptides or variants thereof having in their amino acid sequence one or several tryptophan amino acid residues.

5 In a second preferred embodiment of the first detection method of the invention detailed above, the detection step b) consists of submitting the detectable molecule to a source of energy at the excitation wavelength of said detectable molecule, and measuring the light emitted at the emission wavelength of said detectable molecule.

10 An illustrative example of this second embodiment above is when the marker compound used consists of a Selected Interacting Domain (SID®) which is bound to a fluorescent molecule, such as the fluorescent proteins GFP or YFP.

15 For example, in the embodiment wherein the detectable molecule of the marker compound of the invention which is used according to the first detection method above comprises, or alternatively consists of, a GFP protein, the detection step c) includes illuminating the sample tested at an emission wavelength substantially equal to 490 nm, and measuring the light emitted by the marker compound which is bound
20 to the polypeptide of interest within the sample at an emission wavelength substantially equal to 510 nm.

 Preferably, the marker compounds which are not bound to the polypeptide of interest the detection of which is sought within the sample are removed before carrying out the detection step.

25 In a third preferred embodiment, the detection step c) of the first detection method of the invention consists of measuring the catalytic activity of the detectable molecule. In this specific embodiment, the marker compound used in the detection method comprises a detectable molecule which comprises, or alternatively which consists of, an enzyme
30 or a catalytically active enzyme fragment, such as already detailed in the section entitled " Marker compounds of the invention ".

 In a fourth preferred embodiment, the detection step b) consists of measuring the radioactivity emitted by the detectable molecule.

The present invention further relates to a kit for detecting a polypeptide of interest within a sample, wherein said kit comprises a marker compound according to the invention.

Optionally, said detection kit further comprises the reagents
5 necessary for carrying out the detection step b), such as a suitable substrate for the particular enzyme or a catalytically active enzyme fragment used, as well as suitable buffer solutions, which may be identical to those conventionally used for performing immunoassays.

10 **Cellular detection assay using a recombinantly produced marker compound of the invention.**

As already described above, any marker compound according to the invention may be produced according to genetic engineering
15 techniques. Particularly, nucleic acid encoding a particular marker compound which binds specifically to a polypeptide of interest the detection of which is sought may be inserted in a vector, wherein said vector may be used to transfect or transform a host organism, either a prokaryotic or an eukaryotic cell host such as defined above.

20 In this specific embodiment, the production of a recombinant marker compound of the invention is allowed within such a transfected or transformed host cell. Once the host cell of interest is transfected or transformed with such a recombinant vector and once the recombinant marker compound is produced within the cell host of interest, then the
25 Selected Interacting Domain (SID®) polypeptide portion of said marker compound will be able to bind specifically to its specific target polypeptide within the cell host. In this situation, the recombinantly produced marker compound of the invention will predominantly be localised at cell sites wherein the targeted polypeptide of interest is
30 present.

This is the purpose of the second detection method of the invention which is detailed below.

A further object of the invention consists of a method for detecting a polypeptide of interest within a prokaryotic or an eukaryotic
35 cell host, wherein said method comprises the steps of :

DA.

Le August 2 2000

- a) providing a cell host to be assayed;
- b) transfecting said cell host with a nucleic acid encoding a marker compound of the invention, or with a recombinant vector
5 encoding a marker compound of the invention;
- c) detecting the complexes formed between the marker compound expressed by the transfected cell host and the polypeptide of interest.

Because the Selected Interacting Domain (SID®) polypeptide
10 which is part of a marker compound of the invention specifically binds to a polypeptide which is suspected to be naturally produced by the targeted cell host, the second detection method of the invention defined above allows a qualitative as well as a quantitative detection of this targeted polypeptide which is suspected to be naturally produced by the
15 transfected target cell host under assay.

For example, in the embodiment within which the procedure for selecting the Selected Interacting Domain (SID®) polypeptide which is part of a marker compound of the invention includes a first step wherein a collection of clones containing nucleic acid inserts derived from a H77
20 strain HCV genomic DNA library is prepared, the transfection of a mammalian cell, preferably a human cell, with a vector encoding such a marker compound of the invention will allow to detect the expression of a human polypeptide naturally expressed within said mammalian host cell and which naturally interacts with the HCV viral protein from which is
25 derived the Selected Interacting Domain (SID®) polypeptide.

The second detection method of the invention defined above firstly allows the qualitative detection of the targeted polypeptide of interest which binds specifically with the recombinantly produced marker compound of the invention, and thus permits to know in which
30 environmental conditions or at which differentiation stage the targeted polypeptide of interest is naturally produced within the cell host transfected with a vector expressing a marker compound of the invention.

Secondly, this second detection method of the invention allows
35 the localisation of the targeted polypeptide of interest within the interior

of the cell, including localisation in the plasma membrane, cytosol, nucleus and any organelle such as ribosomes, Golgi apparatus, lysosomes, phagosomes, endoplasmic reticulum and chloroplasts.

5 The localisation of a targeted polypeptide of interest which is expressed within the cell host under assay according to the second detection method of the invention may be carried out by any means well known in the art, including using a confocal microscope.

10 Thirdly, the second detection method of the invention allows also a quantitative analysis of the expression of the targeted polypeptide of interest within the cell host under assay, since the level of the detection signal produced by the detectable molecule which is part of the marker compound will be proportional to the number of complexes formed between the cell host under assay between the targeted polypeptide of interest and the recombinantly produced marker
15 compound of the invention.

Essentially, the one skilled in the art may refer to the section entitled "Acellular detection method of the invention" above to find the teachings necessary for performing the detection step c) of the second detection method described herein.

20 In a first embodiment of said second detection method of the invention, the detection step c) consists of the measure of the fluorescence signal intrinsically emitted by the detectable molecule comprised in the recombinantly expressed marker compound of the invention.

25 In a second preferred embodiment of the second detection method above, the detection step c) consists of submitting the detectable molecule to a source of energy at the excitation wavelength of said detectable molecule and measuring the light emitted at the emission wavelength of said detectable molecule.

30 In still a further embodiment of the second detection method of the invention, the detection step c) consists of measuring the catalytic activity of the detectable molecule.

In another embodiment, the detection step c) consists of measuring the radioactivity emitted by the detectable molecule.

DA.

Le August 2 2000

In yet a further embodiment of the second detection method of the invention, the detection step c) allows the location of the complexes formed between the recombinantly produced marker compound and the targeted polypeptide of interest within the transfected cell host.

5 A further object of the invention consists of a kit for detecting a polypeptide of interest within a prokaryotic or an eukaryotic cell host, wherein said kit comprises a nucleic acid encoding a marker compound as defined herein, or a recombinant vector containing inserted therein a nucleic acid encoding a marker compound of the invention.

10 Optionally, the detection kit above may further comprise the reagents necessary to carry out the detection step c).

Cellular detection method of the invention using a marker compound which is introduced within a cell host.

5 There is a third detection method according to the invention wherein the marker compound comprising a Selected Interacting Domain (SID®) polypeptide OF SEQ ID N°1 to 38 or a variant thereof is previously produced by any means and subsequently introduced into a target cell host for the purpose of detecting a targeted polypeptide of
10 interest which binds specifically with said Selected Interacting Domain (SID®) polypeptide.

 Thus, the invention further relates to a method for detecting a polypeptide of interest within a prokaryotic or an eukaryotic cell host, wherein said method comprises the step of :

- 15 a) providing a cell host to be assayed;
 b) introducing a marker compound as defined herein within said cell host;
 and
 c) detecting the complexes formed between the marker
20 compound and the polypeptide of interest within the cell host.

 Taking into account the low molecular weight of the Selected Interacting Domain (SID®) polypeptide selected from SEQ ID N°1 to 38 which is part of a marker compound of the invention, when compared with conventional specific detection molecules such as antibodies or
25 antibody fragments, it results that the introduction of a marker compound of the invention into the interior of a target cell host will be much more easier to perform, as compared with the introduction within a cell host of a conventional marker like a labelled antibody or a labelled antibody fragment.

30 According to the third detection method of the invention defined above, step b) of introducing the marker compound within the target cell host may be performed by any technique well known in the art, including electroporation, and the use of molecules that will facilitate the passage of the marker compound of the invention through the cell membranes,
35 and typically the plasma membrane.

DA.

Le August 2 2000

Such molecules that facilitate the passage of a marker compound of the invention through cell membranes include, but are not limited to, penetratin, like penetratin 1.RTM (Encor, Gaithersburg, Md), Antenna Pediae protein, cationic lipids and cationic polyacrylates.

5 Permeation enhancers which may be employed include bile salts such as sodium glycocholate and other molecules such as β -cyclodextrin. Bile salts are known to increase the absorption of macromolecules across membranes (Pontiroli et al., 1987).

As already detailed for the second detection method of the
10 invention described in the previous section, the third detection method of the invention allows also the localisation of the targeted polypeptide of interest which is expressed by the cell host under assay, as well as the qualitative and quantitative analysis of the expression of said target polypeptide of interest.

15 The detection step c) according to the third detection method of the invention described above may be carried out in the same way than the detection step c) of anyone of the first detection method and the second detection method detailed in the previous sections herein.

In a first embodiment of the third detection method above, the
20 detection step c) consists of the measure of the fluorescence signal intrinsically emitted by the detectable molecule.

In a second embodiment, the detection step c) consists of submitting the detectable molecule to a source of energy at the excitation wavelength of said detectable molecule and measuring the
25 light emitted at the emission wavelength of said detectable molecule.

In a third embodiment, the detection step c) consists of measuring the catalytic activity of the detectable molecule.

In a fourth embodiment, the detection step c) consists of measuring the radioactivity emitted by the detectable molecule.

30 In a fifth embodiment of the third detection method of the invention, the detection step c) allows the location of the complexes formed between the marker compound and the polypeptide of interest within the target cell host under assay.

A further object of the invention consists of a kit for detecting a polypeptide of interest within a prokaryotic or an eukaryotic cell host, wherein said kit comprises a marker compound as defined herein.

The detection kit above may further comprise the reagents
5 necessary to carry out the detection step c).

The detection kit above may also further comprise the reagents necessary to facilitate the introduction of the marker compound within the target cell host under assay.

10 **SOLID PHASE DETECTION METHOD USING A SELECTED INTERACTING DOMAIN (SID®) POLYPEPTIDE.**

In a further aspect of the invention, the use of a Selected Interacting Domain (SID®) polypeptide of SEQ ID N°1 to 38 or a variant
15 thereof for detection purpose include a step wherein said Selected Interacting Domain (SID®) polypeptide is immobilised on a suitable substrate before bringing a sample to be assayed in contact with the substrate onto which said Selected Interacting Domain (SID®) polypeptide has been previously immobilised.

20 A subsequent step will consist in detecting the complexes formed between the Selected Interacting Domain (SID®) polypeptide immobilised on the substrate and the targeted polypeptide of interest the presence of which is suspected in the sample assayed.

Thus, the invention also pertains to a fourth detection method
25 which consists of a method for detecting a polypeptide or a plurality of polypeptides of interest within a sample, wherein said method comprises the steps of :

a) providing a substrate onto which a Selected Interacting Domain (SID®) polypeptide or a plurality of Selected Interacting Domain
30 (SID®) polypeptides is (are) immobilised;

b) bringing into contact the substrate defined in a) with the sample to be assayed;

c) detecting the complexes formed between the Selected Interacting Domain (SID®) polypeptide or the plurality of Selected

DA.
Le August 2 2000

Interacting Domain (SID®) polypeptides and the target polypeptide or the plurality of target polypeptides contained in the sample.

Substrates, supports or surfaces for immobilising protein molecules are well known in the art, and a lot of them have been described for performing solid phase immunoassays.

Preferably, a plurality of Selected Interacting Domain (SID®) polypeptides of different aminoacid sequences chosen among the sequences SEQ ID N°1 to 38 are immobilised on the substrate used according to the fourth detection method of the invention.

For example, a complete collection of Selected Interacting Domain (SID®) polypeptides which have been determined according to the methods described in the section entitled "Selected Interacting Domain (SID®) polypeptides" above, using nucleic acids derived from the H77 strain HCV genomic DNA as starting material, may be used for being immobilised on a suitable substrate.

According to this embodiment, the collection of Selected Interacting Domain (SID®) polypeptides of SEQ ID N°1 to 38 are immobilised on the substrate in another manner, thus forming an ordered area of SID® polypeptides immobilised at known locations of the surface of said substrate.

The substrate, support or surface may be a porous or a non-porous water insoluble material. The support can be hydrophilic or capable of being rendered hydrophilic and includes inorganic powders such as silica, magnesium sulphate, and alumina; natural polymeric materials, particularly cellulosic materials and materials derived from cellulose, such as fiber containing papers; synthetic or modified naturally occurring polymers, such as nitro-cellulose, cellulose acetate, poly(vinyl chloride), polyacrylamide, cross-linked dextran, agarose, polyacrylate, polyethylene, polypropylene, poly(4-methylbutene), polystyrene, polymethacrylate, poly(ethylene terephthalate), nylon, poly(vinyl butyrate), said materials being used by themselves or in conjunction with other materials; glass available as Bioglass, ceramic metals and the like.

An ordered area onto which a plurality of Selected Interacting Domain (SID®) polypeptides are immobilised may be manufactured

according to the techniques disclosed in the US Patent N°5,143,854 or the PCT Application n°WO 92/10092, incorporated herein by reference for all purposes. The combination of photolithographic and fabrication techniques may, for example, enable each Selected Interacting Domain (SID®) polypeptide to occupy a very small area (" site ") on the support.
5 In some embodiments, the site may be as small as few microns or even a single Selected Interacting Domain (SID®) polypeptide.

In a first embodiment of the fourth detection method detailed
10 above, the plurality of Selected Interacting Domain (SID®) polypeptides are immobilized on the substrate in an order manner.

In a second embodiment of Selected Interacting Domain (SID®), the Selected Interacting Domain (SID®) polypeptide or the plurality of Selected Interacting Domain (SID®) polypeptides are
15 covalently bound to the substrate.

In a third embodiment of said method, the Selected Interacting Domain (SID®) polypeptide or the plurality of Selected Interacting Domain (SID®) polypeptides are non-covalently bound to the substrate. According to this specific embodiment, the Selected Interacting Domain (SID®) polypeptide or the plurality of Selected Interacting Domain (SID®) polypeptides are covalently bound to a first ligand molecule and the substrate is coated with a second ligand molecule, wherein said second ligand molecule specifically binds to the first ligand molecule. According to such a specific embodiment, the first ligand may be biotin in
20 which case the second ligand is most preferably streptavidin.
25

In still a further embodiment of the fourth detection method according to the invention, the Selected Interacting Domain (SID®) polypeptide or the plurality of Selected Interacting Domain (SID®) polypeptides are covalently linked to a spacer, which spacer is itself also
30 covalently bound to the substrate in order to immobilise the Selected Interacting Domain (SID®) polypeptide or the plurality of Selected Interacting Domain (SID®) polypeptides onto said substrate. Such a spacer may be a peptide polymer such as a poly-alanine or a poly-lysine peptide of 10 to 15 amino acids in length.

DA.

Le August 2 2000

In still a further embodiment of the fourth detection method above, the detection step c) consists of detecting changes in the optical characteristics of the substrate onto which the Selected Interacting Domain (SID®) polypeptide or the plurality of Selected Interacting Domain (SID®) polypeptides are bound.

In yet a further embodiment of the fourth detection method of the invention, the detection step c) consists of bringing into contact the substrate wherein complexes are formed between the targeted polypeptide molecule contained in the sample assayed and the Selected Interacting Domain (SID®) polypeptide or the plurality of Selected Interacting Domain (SID®) polypeptides bound to said support, with a detectable molecule having the ability to bind to such complexes.

A further object of the invention consists of a device or an apparatus for the detection of a polypeptide or a plurality of polypeptides of interest within a sample, wherein said device or apparatus comprises a substrate onto which a Selected Interacting Domain (SID®) polypeptide (or a plurality of Selected Interacting Domain (SID®) polypeptides) is (are) immobilised.

Such a device or apparatus of the invention above may comprise or consist of a suitable substrate onto which the plurality of Selected Interacting Domain (SID®) polypeptides are arranged in an ordered manner, thus forming an area such as described above.

PHARMACEUTICAL COMPOSITIONS CONTAINING A SELECTED INTERACTING DOMAIN (SID®) POLYPEPTIDE.

It results from the method according to which a Selected Interacting Domain (SID®) polypeptide of SEQ ID N°1 to 38 has been selected and characterized that such a Selected Interacting Domain (SID®) polypeptide or a variant thereof is both:

(i) endowed with highly specific binding properties to a (bait) polypeptide of interest;
and

(ii) devoided of the biological activity of the naturally occurring protein from which this Selected Interacting Domain (SID®) polypeptide or a variant thereof is derived.

These original properties of a Selected Interacting Domain (SID®) polypeptide of SEQ ID N°1 to 38 or a variant thereof allow its use for interfering with a naturally occurring interaction between a first protein and a second protein within the cell of an organism by the binding of said Selected Interacting Domain (SID®) polypeptide specifically either to said first polypeptide or said second polypeptide.

The (SID®) polypeptides of the invention or variants thereof are capable of interfering with the *in vivo* protein-protein interactions between HCV proteins or between a HCV protein and a protein from the organism which has been infected with the Hepatitis C virus.

For example the SID® polypeptide of SEQ ID N°2 interferes with the naturally occurring interaction between the core and the NS3 protein HCV. Similarly, the SID® polypeptide of SEQ ID N°17 interferes with the interaction between the NS4A and the NS4B proteins (see table 1).

Thus, another object of the invention consists of a pharmaceutical composition comprising a pharmaceutically effective amount of a Selected Interacting Domain (SID®) polypeptide or a variant thereof.

The invention also relates to a pharmaceutical composition comprising a pharmaceutically effective amount of a nucleic acid comprising a polynucleotide encoding a Selected Interacting Domain (SID®) polypeptide of SEQ ID N°1 to 38 or a variant thereof which polynucleotide is placed under the control of an appropriate regulatory sequence.

Preferred nucleic acids are the nucleotide sequences SEQ ID N°39 to 76.

The invention also pertains to a pharmaceutical composition comprising a pharmaceutically effective amount of a recombinant expression vector comprising a polynucleotide encoding the Selected Interacting Domain (SID®) polypeptide or a variant thereof.

DA.
Le August 2 2000

The invention also pertains to a method for preventing or curing a viral infection by a hepatitis C virus in a human or an animal, wherein said method comprises a step of administering to the human or animal body a pharmaceutically effective amount of a Selected Interacting Domain (SID®) polypeptide of SEQ ID N°1 to 38 or a variant thereof
5 which binds to a targeted viral or mammal, typically- human protein.

A pharmaceutical composition as described above, wherein said composition is administered by any route, such as intravenous route, intramuscular route, oral route, or mucosal route with an
10 acceptable physiological carrier and/or adjuvant, also forms part of the invention.

The Selected Interacting Domain (SID®) polypeptide or a variant thereof as a medicament for the prevention and/or treatment of pathologies induced by HCV are the most preferred.

15 The Selected Interacting Domain (SID®) polypeptides of SEQ ID N°1 to 38 as active ingredients of a pharmaceutical composition will be preferably in a soluble form combined with a pharmaceutically acceptable vehicle.

Such compounds which can be used in a pharmaceutical
20 composition offer a new approach for preventing and/or treating pathologies linked to infection by HCV. Preferably, these compounds will be administered by the systemic route, in particular by the intravenous route, by the intramuscular or intradermal route or by the oral route.

Their modes of administration, optimum dosages and galenic
25 forms can be determined according to the criteria generally taken into account in establishing a treatment suited to a patient, such as for example the age or body weight of the patient, the seriousness of his general condition, the tolerance to treatment and the side effects observed, and the like.

30 The identified compound can be administered to a mammal, including a human patient, alone or in pharmaceutical compositions where they are mixed with suitable carriers or excipients at therapeutically effective doses to treat disorders associated with prokaryotic micro-organism infection. Techniques for formulation and
35 administration of the compounds of the invention may be found in

" Remington's Pharmaceutical Sciences " Mack Publication Co., Easton, PA, latest edition.

For any Selected Interacting Domain (SID®) polypeptide or any variant thereof used according to the invention, the therapeutically effective dose can be estimated initially from cell culture assays. For example, a dose can be formulated in animal models to achieve a circulating concentration range that includes or encompasses a concentration point or range shown the desired effect in an *in vitro* system. Such information can be used to more accurately determine useful doses in humans.

A therapeutically effective dose refers to that amount of the compound that results in amelioration of symptoms in a patient. Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g. for determining the LD50, (the dose lethal to 50% of the test population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio between LD50 and ED50. Compounds which exhibit high therapeutic indices are preferred.

The data obtained from these cell culture assays and animal studies can be used in formulating a range of dosage for use in human. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED50, with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilised. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. (See, e.g. Fingl et al. 1975, in " The Pharmacological Basis of Therapeutics ", CH.I).

Dosage amount and interval may be adjusted individually to provide plasma levels of the active compound which are sufficient to maintain the modulating effects. Dosages necessary to achieve the modulating effect will depend on individual characteristics and route of administration.

DA.

Le August 2 2000

The amount of composition administered will, of course, be dependent on the subject being treated, on the subject's weight, the severity of the affliction, the manner of administration and the judgement of the prescribing physician.

5 The invention also pertains to a method for preventing or curing a viral in a human or an animal, wherein said method comprises the step of administering to the human or animal body a pharmaceutically effective amount of a nucleic acid comprising a polynucleotide encoding a Selected Interacting Domain (SD®) polypeptide of SEQ ID N°1 to 38,
10 or a variant thereof, and wherein said polynucleotide is placed under the control of a regulatory sequence which is functional in said human or said animal.

Preferred polynucleotides are the nucleic acids of SEQ ID N°39 to 76.

15 The invention also relates to a method for preventing or curing a viral or in a human or an animal, wherein said method comprises the step of administering to the human or animal body a pharmaceutically effective amount of a recombinant expression vector comprising a polynucleotide encoding a Selected Interacting Domain (SD®)
20 polypeptide which binds to a viral or bacterial protein.

Other characteristics and advantages of the invention appear in the remainder of the description with the examples below, without linking the invention in any manner.

25 **EXAMPLES:**

Preparation of a HCV genomic collection.

1.A. Collection preparation and transformation in *Escherichia coli*

30 **1.A.1 Fragmentation of genomic DNA preparation.**

The genomic DNA of the infectious HCV strain H77 (Yanagi et al., P.N.A.S. 1997, 94, 8738-43) is fragmented in a nebulizer (GATC) for 2 minutes at a pressure of 2 bars, precipitated and resuspended in water.

The obtained nubilized genomic DNA is successively treated with Mung Bean Nuclease (Biolabs) (30 minutes at 30°C), T4 DNA polymerase (Biolabs) (10 minutes at 37°C) and Klenow enzyme (Pharmacia) (10 minutes at room temperature and 1 hour at 16°C).

5 DNA is then extracted, precipitated and resuspended in water.

1.A.2. Ligation of linkers to blunt-ended genomic DNA

Oligonucleotide HGX931 (5' end phosphorylated) 1 µg/µl and HGX932
10 1µg/µl.

Sequence of the oligo HGX931: 5'-GGGCCACGAA-3' (SEQ ID N°151).

Sequence of the oligo HGX932: 5'-TTCGTGGCCCCTG-3'(SEQ ID N°152).

Linkers were preincubated (5 minutes at 95°C, 10 minutes at
15 68°C, 15 minutes at 42°C) then cooled down at room temperature and ligated with genomic DNA inserts at 16°C overnight.

Linkers were further removed on a separation column (Chromaspin TE 400, Clontech), according to the manufacturer's protocol.

20

1.A.3. Vector preparation

Plasmid pP6 (see figure 13) was prepared by replacing the Spe1/Xho1 fragment of pGAD3S2X with the double-stranded oligonucleotide:

25

5'CTAGCCATGGCCGCAGGGGCCGCGGCCGCACTAGTGGGGATCCTTAATTAAAG
GGCCACTGGGGCCCCCGTACCGGCGTCCCCGGCGCCGGCGTGATCACCCCTA
GGAATTAATTTCCCGGTGACCCCGGGGAGCT 3' (SEQ ID N°153).

30 The pP6 vector is successively digested with Sfi1 and BamHI restriction enzymes (Biolabs) for 1 hour at 37°C, extracted, precipitated

DA.
Le August 2 2000

and resuspended in water. Digested plasmid vector backbones are purified on a separation column (Chromaspin TE 400, Clontech), according to the manufacturer's protocol.

5 **1.A.4 Ligation between vector and insert of genomic DNA**

The prepared vector is ligated overnight at 15°C with the genomic blunt-ended DNA described in section 2 using T4 DNA ligase (Biolabs). The DNA is then precipitated and resuspended in water.

10

1.A.5. Library transformation in *Escherichia coli*.

Transform DNA from section 1.A.4. into Electromax DH10B electrocompetent cells (Gibco BRL) with Cell Porator apparatus (Gibco
15 BRL). Add 1 ml SOC medium and incubate transformed cells at 37°C for 1 hour. Add 9 ml volume of SOC medium per tube and plate on LB+ampicillin medium. Scrape colonies with liquid LB medium. Aliquot and freeze at -80°C .

The obtained collection of recombinant cell clones is named
20 HGXBHCV1.

1.B. Collection transformation in *Saccharomyces cerevisiae*

The *Saccharomyces cerevisiae* strain (Y187 (MAT α Gal4 Δ
25 Ga180 Δ ade2-101 His3 Leu2-3, -112 Trp1-901 Ura3-52 URA3::UASGAL1-LacZ Met) transformed with the HGXBHCV1 HCV genomic DNA library.

The plasmid DNA contained in *E. coli* are extracted (Qiagen) from aliquoted *E. coli* frozen cells (1.A.5.).

30 Grow *Saccharomyces cerevisiae* yeast Y187 in YPGlu.

Yeast transformation is performed according to standard protocol (GUEST et al. Yeast, 11, 355-360, 1995) using yeast carrier DNA (Clontech). This experiment leads to 10^4 to 5.10^4 cells/ μ g DNA. Spread 2.10^4 cells on DO-Leu medium per plates. Aliquot and freeze at -80°C .
5 The obtained collection of recombinant cell clones is named HGXYHCV1.

1.C. Construction of bait plasmids

10 Plasmid pB5 (see figure 12) is prepared by replacing the NcoI/SalI polylinker fragment with the double-stranded oligonucleotide.

5'CATGGCCGCAGGGGCCGCGGCCGCACTAGTGGGGATCCTTAATTAAAGGGCCA
CTGGGGCCCCCGGCGTCCCCGGCGCGCGGTGATCACCCCTAGGAATTAATTT
15 CCCGGTGACCCCGGGGGAGCT 3'. (SEQ ID N°154).

The linked genomic DNA described in section 2 is ligated into pB5 that has been digested with SfiI restriction enzyme and DNA transformed into competent *E. coli*. Cells are grown and plasmid DNA
20 extracted and sequenced. Those plasmids which code in-frame fusion proteins are used as bait plasmids.

EXAMPLE 2 : Screening the collection with the two-hybrid in yeast system.

25

2.A. The mating protocol.

We have chosen the mating two-hybrid in yeast system (firstly described by FROMONT-RACINE et al., Nature Genetics, 1997, vol. 16,
30 277-282, Toward a functional analysis of the yeast genome through exhaustive two-hybrid screens) for its advantages but we could also

DA.
Le August 2 2000

screen the HCV collection in classical two-hybrid system as described in Fields et al. or in a yeast reverse two-hybrid system.

The mating procedure allows a direct selection on selective plates because the two fusion proteins are already produced in the parental cells. No replica plating is required. This protocol is written for the use of the library transformed into the Y187 strain.

Before mating, transform *S. cerevisiae* (CG 1945 strain (MATa Ga14-542 Gal180-538 ade2-101 His3*200 Leu2-3, -112 Trp1-901 Ura3-52 Lys2-801 URA::GAL4 17 mers (X3)- CyC1TATA-LacZ LYS2::GAL1UAS-GAL1TATA-HIS3 CYH^R)) according to step 1.B. and spread on DO-Trp medium.

Day 1, morning: preculture

Preculture of Y187 cells carrying the bait plasmid obtained at step 1.C. in 20 ml DO-Trp medium. Grow at 30°C with vigorous agitation.

Day 1, late afternoon: culture

Measure OD_{600nm} of the DO-Trp pre-culture of Y187 cells carrying the bait plasmid preculture. The OD_{600nm} must lie between 0.1 and 0.5 in order to correspond to a linear measurement.

Inoculate 50 ml DO-Trp at OD_{600nm} 0.006/ml, grow overnight at 30°C with vigorous agitation.

Day 2 : mating

medium and plates

1 YPGlu 15 cm plate

50 ml tube with 13 ml DO-Leu-Trp-His

100 ml flask with 5 ml of YPGlu

8 DO-Leu-Trp-His plates

2 DO-Leu plates

2 DO-Trp plates

2 DO-Leu-Trp plates

Measure OD_{600nm} of the DO-Trp culture. It should be around 1.

- 5 For the mating, you must use twice as many bait cells as library cells. To get a good mating efficiency, you must collect the cells at 10^8 cells per cm².

Estimate the amount of bait culture (in ml) that makes up 30 OD_{600nm} units for the mating with the prey library.

- 10 Thaw a vial containing the HGXYHCV1 library slowly on ice. Add the 0.5 ml of the vial to 5 ml YPGlu. Let those cells recover at 30°C, under gentle agitation for 10 minutes.

Mating

- 15 Put the 30 OD_{600nm} units of bait culture into a 50 ml flacon tube. Add the HGXYHCV1 library culture to the bait culture. Centrifuge, discard the supernatant and resuspend in 0.8 ml YPGlu medium.

- 20 Distribute the cells onto a YPGlu plate with glass beads. Spread cells by shaking the plates.

Incubate the plate cells-up at 30°C for 4 h 30 min.

Collection of mated cells

- 25 Wash and rinse the plate with 6 ml and 7 ml consecutively of DO-Leu-Trp-His.

- Perform two parallel serial ten-fold dilutions in 500 µl DO-Leu-Trp-His up to 1/10,000. Spread out 50 µl of each 1/10000 dilution onto DO-Leu and DO-trp plates and 50 µl of each 1/1000 dilution onto DO-
- 30 Leu-Trp plates.

DA.

Le August 2 2000

Spread 3.2 ml of collected cells in 400 µl aliquots on DO-Leu-Trp-His+Tet plates.

DAY 4

5

Selection of clones able to grow on DO-Leu-Trp-His+Tetracyclin: this medium allows us to isolate diploid clones presenting an interaction.

Count the Trp+Leu+ colonies on control plates and the total
10 number of His+ colonies on the DO-Leu-Trp-His+Tetracyclin plates.

The number of His+ cell clones will define which protocol is to be processed:

Upon $2 \cdot 10^6$ Trp+Leu+ colonies:

- if number of His+ cell clones < 95: then process luminometry
15 protocol on all colonies;
- if number of His+ cell clones > 95 and <5000: then process luminometry protocol on 95 colonies;
- if number of His+ cell clones >500: repeat screen using DO-Leu-Trp-His+Tetracyclin plates containing 3-aminotriazol.

20

2.B The luminometry assay

Grow His+ colonies overnight at 30°C in microtiter plates containing DO-Leu-Trp-His-Tetracyclin medium with shaking. The day
25 after, dilute 15 times overnight culture into a new microtiter plate containing the same medium. Incubate 5 hours at 30°C with shaking. Dilute samples 5 times and read OD_{600nm}. Dilute again to obtain between 10 000 and 75 000 yeast cells/well in 100 µl final volume.

Per well, add 76 μ l of One Step Yeast Lysis Buffer (Tropix), 20 μ l SapphireII Enhancer (Tropix), 4 μ l Galacton Star (Tropix), incubate 40 minutes at 30°C.

Measure the β -Gal read-out (L) using a Luminometer (Trilux,
5 Wallach).

Calculate value of OD_{600nm}XL and selected interacting preys having highest values.

At this step of the protocol, we have isolated diploid cell clones presenting interaction. The next step is now to identify polypeptides
10 involved in the selected interactions.

EXAMPLE 3: Identification of positive clones

3.A. PCR on yeast colonies

15 Introduction

PCR amplification of fragments of plasmid DNA directly on yeast colonies is a quick and efficient procedure to identify sequences cloned into this plasmid. It is directly derived from a published protocol (Wang H. et al., Analytical Biochemistry, 237, 145-146, 1996). However,
20 it is not a standardized protocol: in our hands it varies from strain to strain, and is dependent on experimental conditions (number of cells, Taq polymerase source, etc). This protocol should be optimized to specific local conditions.

25 MATERIALS

- For 1 well, PCR mix composition is:
32.5 μ l water,
5 μ l 10X PCR buffer (Pharmacia),
30 1 μ l dNTP 10 mM,

DA.
Le August 2 2000

0,5 µl Taq polymerase (85µ/µl -Pharmacia),
 0,5 µl oligonucleotide ABS1 10 pmole/µl:5'-
 GCGTTTGAATCACTACAGG-3',
 0,5 µl oligonucleotide ABS2 10 pmole/µl:5'-
 5 CACGATGCACGTTGAAGTG-3'.
 - 1N NaOH.

Experiment

10 Grow positive colonies overnight at 30°C on a 96 well cell culture cluster (Costar), containing 150 µl DO-Leu-Trp-His+Tetracyclin with shaking. Resuspend culture and transfer immediately 100 µl on a Thermowell 96 (Costar).

Centrifuge 5 minutes at 4000 rpm at room temperature.

15 Remove supernatant. Dispense 5 µl NaOH in each well, shake 1 minute.

Place the Thermowell in the thermocycler (GeneAmp 9700, Perkin Elmer) 5 minutes at 99,9°C and then 10 minutes at 4°C.

In each well, add PCR mix, shake well.

20 Set up the PCR program as followed:

94°C 3 minutes

94°C 30 seconds

53°C 1 minute 30 seconds x 35 cycles

72°C 3 minutes

25 72°C 5 minutes

15°C ∞

Check the quality, the quantity and the length of the PCR fragment on agarose gel.

The length of the cloned fragment is the estimated length of the PCR fragment minus 300 base pairs that correspond to the amplified flanking plasmid sequences.

5 **3.B Plasmids rescue from yeast by electroporation**

Introduction

10 The previous protocol of PCR on yeast cell may not be successful, in such a case, we rescue plasmids from yeast by electroporation. This experiment allows the recovery of prey plasmids from yeast cells by transformation of *E.coli* with a yeast cellular extract. We can then amplify the prey plasmid and sequence the cloned fragment.

15

Material

Plasmid rescue

Glass beads 425-600 μ m (Sigma)

20 Phenol/chloroform (1/1) premixed with isoamyl alcohol (Amresco)

Extraction buffer: 2% Triton X100, 1% SDS, 100 mM NaCl, 10 mM TrisHCl pH 8,0, 1 mM EDTA pH 8.0.

Mix ethanol/ NH_4Ac : 6 volumes ethanol with 7.5 M NH_4 Acetate, 70% Ethanol and yeast cells in patches on plates.

25

Electroporation

SOC medium

M9 medium

Selective plates: M9-Leu+Ampicillin

30

2 mm electroporation cuvettes (Eurogentec)

DA.

Le August 2 2000

Experiment

Plasmid rescue

- 5 Prepare cell patch on DO-Leu-Trp-His with cell culture of section 2.C.

Scrape the cell of each patch in Eppendorf tube, add 300 µl of glass beads in each tube, then add 200 µl extraction buffer and add 200µl phenol: chloroform:isoamyl alcohol (25:24:1).

- 10 Centrifuge tubes 10 minutes at 15000 rpm.

Transfer 180 µl supernatant to a sterile Eppendorf tube and add to each 500 µl ethanol/NH₄Ac, vortex.

Centrifuge tubes 15 minutes, 15000 rpm at 4°C.

Wash pellet with 200 µl 70% ethanol, remove ethanol and dry pellet,

- 15 Resuspend pellet in 10 µl water. Store extracts at -20°C.

Electroporation

Material: Electrocompetent MC1066 cells prepared according to standard protocols (Maniatis).

- 20 Add 1 µl of yeast plasmid DNA-extract to pre-chilled Eppendorf tube, and keep on ice.

Mix 1 µl plasmid yeast DNA-extract sample, add 20 µl electrocompetent cells and transfer in a cold electroporation cuvette.

Set the Biorad electroporator on 200 ohms resistance, 25 µF capacity;

- 25 2.5 kV. Place cuvette in the cuvette holder and electroporate.

Add 1 ml SOC into the cuvette and transfer the cell-mix into sterile Eppendorf tube.

Let cells recover for 30 minutes at 37°C, spin the cells down 1 minute, 4000x g and pour off supernatant. Keep about 100 µl medium and use it

to resuspend the cells and spread them on selective plates (e.g. M9-Leu plates).

Incubate plates for 36 hours at 37°C.

Grow one colony and extract plasmids. Check presence and size of
5 insert through enzymatic digestion and agarose gel. Sequence insert.

EXAMPLE 4: Protein-protein interaction.

For each bait, the previously protocol leads to the identification
10 of prey polynucleotide sequences. Using a suitable software program (eg Blastwun, available on the Internet site of the University of Washington: <http://bioweb.pasteur.fr/seqanal/interfaces/blastwu.html>) the region of the HCV genome is encoded by the prey fragment may be determined and whether the fusion proteins encoded are in the same open reading frame
15 of translation as the HCV polyprotein or not.

EXAMPLE 5 : Identification of SID®

The presence of contiguous polypeptides in the HCV genome
20 and the high complexity of the prey library used prevents the determination of SID®s by previous means since prey fragments can overlap multiple polypeptides. The high complexity of the prey library used relative to the small genome size also prevented such a simple analysis since prey fragments can overlap multiple interacting domains.
25 It was also necessary to overcome the problems caused by protein preys encoded by out-of-frame fusions of regions of the HCV genome.

In order to determine the SID®s for a particular bait protein, it was therefore necessary to devise a suitable algorithm which would take into account all these problems:

DA.
Le August 2 2000

5.1. The prey fragments are initially sorted according to which reading frame of the polypeptide sequence they correspond to. This enables the separation of physiologically relevant prey protein from out-of-frame fusions which bind in the two-hybrid assay.

5 5.2. Each prey fragment is compared pairwise with other prey fragments and two fragments are clustered together if they overlap by more than 30% of their lengths (see fig. 8). Further fragments are assigned to the cluster if, and only if, overlap all the fragments in the cluster by more than 30% of their length.

10 5.3 For each cluster of fragments thus produced, a pre-SID is defined as the intersection of all the fragments present in the cluster defined in 5.2 (figure 9).

5.4. The pre-SIDs defined in 5.3 are then analysed pairwise and if the region of intersection between two pre-SIDs is greater than 30 bp
15 then a SID® is defined as this region of intersection. If the non-intersecting region of a pre-SID is of more than 30 bp in length and this non-intersecting region represents more than 30% of the length of one of the fragments that comprises this region, then this non-intersecting region is also defined as a SID@s (figure 10).

20 5.5 The number of fragments contributing to each SID defined in 5.4 is counted. In the case of overlapping SIDs®, the SID® which contains the most fragments is identified, and all the fragments which contribute to this SID® are removed from overlapping SIDs®. The inspection of the fragments which remain in these overlapping SIDs®
25 determines the final sequence of the SID® (figure 11).

TABLE 1
Summary of the protein-protein interactions
between the SID polypeptides of the invention
and H77 strain HCV polypeptides

Bait	SEQ ID N°(1)	begin(2)	end(2)	SEQ ID N°(3)	SID	SEQ ID N° (4)	begin (2)	end (2)	SEQ ID N° (5)
Core(87%)	114	302	614	77	Core(100%)	39	446	600	1
Core(100%)	115	342	683	78	NS3(100%)	40	4814	4922	2
Core(100%)	115	342	683	78	Core(100%)	41	380	616	3
E1(100%)	116	995	1342	79	E2(100%)	42	1871	1987	4
E1(4%)/E2(95 %)	117	1478	1756	80	NS3(100%)	43	4787	5242	5
E2(100%)	118	1745	2278	81	E2(100%)	44	1871	1958	6
E2(100%)	119	1799	2090	82	E2(100%)	45	1808	1890	8
NS2(12%)/NS3 (87%)	120	3312	4150	83	NS4A(59%)/ NS4B(40%)	46	5375	5542	9
NS3(100%)	121	3767	4244	84	NS3(100%)	47	4676	4801	10
NS3(100%)	122	3779	4571	85	NS3(100%)	48	4856	4945	11
NS3(100%)	123	3974	4559	86	NS3(100%)	49	4817	4903	12
NS3(100%)	124	4238	4857	87	NS5B(100%)	50	7979	8109	13
NS3(100%)	125	4298	4859	88	NS3(100%)	51	4031	4118	14
NS3(100%)	126	4691	5168	89	E2(100%)	52	1784	1888	15
NS3(100%)	127	4838	5230	90	E2(100%)	53	1871	1968	16
NS3(1%)/NS4 A(98%)	128	5310	5467	91	NS4B(100%)	54	5918	6154	17
NS4A(100%)	129	5342	5400	92	NS3(100%)	55	3512	3956	18
NS4B(86%)/N S5A(13%)	130	5717	6344	93	NS4B(53%)/ NS5A(46%)	56	6197	6310	19
NS4B(70%)/N S5A(29%)	131	5819	6444	94	E2(100%)	57	1844	1933	20
NS4B(55%)/N S5A(44%)	132	5882	6562	95	NS5B(100%)	58	9083	9222	21
NS4B(82%)/N S5A(17%)	133	5897	6335	96	NS4B(100%)	59	5819	6080	22
NS4B(100%)	134	6011	6177	97	E2(100%)	60	1823	1955	23

5

DA.
 Le August 2 2000

TABLE 1 (continued)

Bait	SEQ ID N°(1)	begin(2)	end(2)	SEQ ID N°(3)	SID	SEQ ID N°(4)	begin(2)	end(2)	SEQ ID N°(5)
NS4B(30%)/NS5A(69%)	135	6107	6605	98	NS4B(100%)	61	5879	6072	24
NS4B(12%)/NS5A(87%)	136	6141	7069	99	E2(100%)	62	1784	1875	25
NS4B(8%)/NS5A(91%)	137	6182	7034	100	E1(100%)	63	1226	1458	26
NS4B(9%)/NS5A(90%)	138	6188	6939	101	NS4B(70%)/NS5A(28%)	64	6176	6291	27
NS5A(100%)	139	6317	6576	102	NS3(100%)	65	4784	4928	28
NS5A(100%)	140	6440	6727	103	NS5A(100%)	66	6557	6721	29
NS5A(100%)	141	7019	7249	104	NS3(100%)	67	4451	4790	30
NS5A(100%)	142	7274	7549	105	NS4B(100%)	68	6029	6194	31
NS5B(100%)	143	7613	8027	106	NS5B(100%)	69	8354	8665	32
NS5B(100%)	144	7838	8743	107	NS5B(100%)	70	7769	8011	33
NS5B(100%)	145	7856	8458	108	NS3(100%)	71	4715	4901	34
NS5B(100%)	146	7976	8759	109	NS5B(100%)	72	7775	8011	35
NS5B(100%)	147	8564	8948	110	E2(100%)	73	1805	1887	36
NS5B(100%)	148	8708	8978	111	E2(100%)	74	1751	1865	37
NS5B(100%)	149	8996	9220	112	NS4B(57%)/NS5A(41%)	75	6194	6303	38
NS5B(100%)	150	9032	9226	113	NS4B(63%)/NS5A(35%)	76	6206	6286	39

- 5 (1) Nucleic acid sequence encoding the polypeptide from the H77 strain of HCV which binds to the SID polypeptide (4) described in the same line.
- (2) 5'-end and 3'-end nucleotide positions of the sequence SEQ ID (1) in reference to the nomenclature disclosed by Yanagi et al. (1997)
- 10 (3) Aminoacid sequence of the polypeptide from the H77 strain of HCV which binds to the SID polypeptide (4) described in the same line.
- (4) Nucleic acid sequence encoding the SID polypeptide which binds to the polypeptide of the aminoacid sequence (3) described in the same line.
- 15 (5) Aminoacid sequence of the SID polypeptide which binds to the polypeptide of the aminoacid sequence (3) described in the same line.

REFERENCES

- Aguzzi F et al. , Estratto Dal. Boll. 1 st Sieroter; Milanese, 1977, vol.56: 212-216.
- 5 • Brigham KL et al., 1993, Am. J. Respir Cell Mol. Biol.8(2):209-213
- BARTENSCHLAGER R. et al., 1995, J. Virol., 69 (12): 7519-7528.
- Curiel et al. Gene Transfer to Respiratory Epithelial Cells via the Receptor Mediated Endocytosis Pathway, Am. J. Respir. Cell Mol. Biol. 6 (1992) 247-252
- 10 • Curiel et al. Adenovirus Enhancement of Transferrin-Polylysine-Mediated Gene Delivery, Proc. Natl. Acad. Sci. 88 (1991) 8850-8854
- Chalfie et al., (1994), Science, vol. 263: 802-805.
- Chen F-T.A. et al. , 1991, Clin. Chem., vol.77: 14-19.
- 15 • Davis et al., (1995), Development Biology, vol.170: 726-729
- Delagrave et al., (1995), Biotechnology, vol.13: 151-154
- DI MARCO et al., 2000, The Journal of Biological Chemistry, vol.275 (10):7152-7157
- 20 • DUBUISSON J., 1994, J.Viral. vol.68:6147-6160
- Drumm, M. L. et al., Cell 62:1227-1233 (1990).
- Fromont-Racine M et al. (1997), Nature Genetics, vol.16 (3):277-282.
- FLAJOLET M. et al., 2000 Gene, vol.242:369-379.
- 25 • EDWARDS and LEARTHERBARROW, 1997, Analytical Biochemistry, 246:1-6.
- Goding et al. J.W., (1986), In: Monoclonal antibodies: Principles and practice-production and application of monoclonal antibodies in cell biology, biochemistry and immunology, Acad. Press, London, pages 255-280.
- 30 •
- GALLINARY P et al., 1999, Biochemistry, vol.38:5620-5632
- GRAKOU I A., 1993, J.Viral., vol.67:1385-1395.

DA.

Le August 2 2000

- HOUGHTON, M (1996), Hepatitis C virus, fields editors.
- HIGIKATA M., 1993, J.Viral., vol.67:4665-4675
- Heim et al., (1994), Proc. Natl. Acad. Sci., volume 91: 12501-12.504.
- 5 • Hu and Cheng, (1995), Febs. Letters, vol.369: 331-334.
- Ichinose N et al., (1991), In: Fluorometric analysis in biomedical chemistry, vol.10, page 110, Chemical analysis, Winefordner JD et al. Eds., John Wiley and Sons, New York.
- KARIMOVA et al., 1998, Proc. Natl. Acad. Sci., USA, 95:5752-5756.
- 10 • Keegan et al. (1986), Science, vol.231 (4739): 699-704.
- KOCH Y, 1977, Biochem. Biophys. Res. Commun, vol.74:488-491
- Kohler and Milstein, 1975, Nature, 256: 495
- Kozbor et al., 1983, Hybridoma, 2(1):7-16.
- KEEGAN et al., 1986, Science, Vol.231:689-407
- 15 • Kaether and Gerdes, (1995), Febs. Letters, vol.369:267-271.
- Leger et al., 1997, Hum. Antibodies, 8(1):3-16
- Martineau et al., 1998, J. Mol. Biol., 280(1): 117-127
- Muzyczka, N., Curr. Top. Micro. Immuno. 158:97-129 (1992)
- Merlini G et al. , 1983, J. Clin. Chem. Biochem. , vol.21: 841-844,
- 20 • Maggio ET, " Enzyme-immuno assay ", 1980, CRC Press Incorporated , Boca Raton, Fla.
- MA and PATSHNE, 1987, Cell, vol. 48: 847-853
- MIN et al., 1999, virus genes, vol.19 (1):33-43
- Nielsen et al., 1991, J. Chromatogr., vol.539: 177
- 25 • PATEL J. et al. 1999, Journal of General Virology, vol.80:1681-1690.
- Pontiroli et al., 1987, Diabet. Metab., vol.13:441-443.
- Ridder et al., 1995, Biotechnology (NY), 13(3):255-260
- Reinmann et al., 1997, AIDS Res. Hum. Retroviruses, 13(II): 933-943
- 30 • Rosenfeld, M. A. et al., Cell 68:143-155 (1992)

- Rizzuto et al. , (1995), Current Biology, vol.5: 635-142.
- ROUGEOT C et al., 1994, Eur. J. Biochem., vol.219(3):765-773
- Shattil SJ et al., (1987) , Blood, vol.70: 307.
- Shattil et al. SJ(1985), J. Biol. Chem., vol.260:11.107.
- 5 • Smith *et al.*, 1988, Gene 67:31-40.
- Schofield, Brit. Microencapsulated. Bull., 51(1):56-71 (1995)
- Behr, Bioconjugate Chem., 5, 382-389 (1994)
- SZABO A. et al., 1995, Curr. Opin. Struct. Biol., 5(5): 699-705.
- Trubetskoy, V. S. et al., Biochem. Biophys. Acta 1131:311-313
- 10 (1993))
- UTKIEWICZ NJ et al., 2000, vol. 267: 278-282
- URBANI A et al., 1999, Biochemistry, vol.38:5206-5215
- Wu et al., 1992, J. Biol. Chem. 267:963-967.
- Wu and Wu, 1988, J. Biol. Chem. 263:14621-14624.
- 15 • Wilson, J. M. et al., 1992, Endocrinology, 130(5):2947-2954
- White Wa et al. , 1986, Biochem. Clin. vol.10:571-574.
- Yanagi et al., Proc. Nat. Acad. Sci USA, 1997, 94 :8738-8743

Claims

1. A nucleic acid which encodes a polypeptide selected from the group consisting of the amino acid sequences SEQ ID N° 1 to 38 or a variant thereof, and a sequence complementary thereto.
5
2. A nucleic acid according to claim 1 which encodes a polypeptide having at least 95% aminoacid identity with a polypeptide selected from the group consisting of the aminoacid sequences SEQ ID N°1 to 38, and a sequence complementary thereto.
10
3. A nucleic acid according to claim 1 which is selected from the group consisting of the sequences SEQ ID N°39 to 76 or a sequence complementary thereto.
15
4. A nucleic acid according to claim 1 which possesses at least 95% nucleic acid identity with a nucleic acid selected from the group consisting of the sequences SEQ ID N°39 to 76.
- 20 5. A nucleic acid according to claim 1 encoding a polypeptide having an aminoacid sequence selected from the group consisting of the sequences consisting of at least:
 - 45 consecutive aminoacids of SEQ ID N°1;
 - 30 consecutive aminoacidss of SEQ ID N°2;
 - 25 - 65 consecutive aminoacids of SEQ ID N°3;
 - 30 consecutive aminoacids of SEQ ID N°4;
 - 130 consecutive aminoacids of SEQ ID N°5;
 - 25 consecutive aminoacids of SEQ ID N°6;
 - 23 consecutive aminoacids of SEQ ID N°7.
 - 30 - 48 consecutive aminoacids of SEQ ID N°8;
 - 36 consecutive aminoacids of SEQ ID N°9;
 - 25 consecutive aminoacids of SEQ ID N°10;

- 24 consecutive aminoacids of SEQ ID N°11;
- 37 consecutive aminoacids of SEQ ID N°12;
- 25 consecutive aminoacids of SEQ ID N°13;
- 30 consecutive aminoacids of SEQ ID N°14;
- 5 - 27 consecutive aminoacids of SEQ ID N°15;
- 69 consecutive aminoacids of SEQ ID N°16;
- 130 consecutive aminoacids of SEQ ID N°17;
- 33 consecutive aminoacids of SEQ ID N°18;
- 25 consecutive aminoacids of SEQ ID N°19;
- 10 - 40 consecutive aminoacids of SEQ ID N°20;
- 78 consecutive aminoacids of SEQ ID N°21;
- 39 consecutive aminoacids of SEQ ID N°22;
- 57 consecutive aminoacids of SEQ ID N°23;
- 26 consecutive aminoacids of SEQ ID N°24;
- 15 - 68 consecutive aminoacids of SEQ ID N°25;
- 34 consecutive aminoacids of SEQ ID N°26;
- 42 consecutive aminoacids of SEQ ID N°27;
- 48 consecutive aminoacids of SEQ ID N°28.
- 102 consecutive aminoacids of SEQ ID N°29:
- 20 - 49 consecutive aminoacids of SEQ ID N°30;
- 92 consecutive aminoacids of SEQ ID N° 31;
- 49 consecutive aminoacids of SEQ ID N°30;
- 92 consecutive aminoacids of SEQ ID N°31;
- 71 consecutive aminoacids of SEQ ID N°32;
- 25 - 55 consecutive aminoacids of SEQ ID N°33;
- 69 consecutive aminoacids of SEQ ID N°34;
- 23 consecutive aminoacids of SEQ ID N°35;
- 33 consecutive aminoacids of SEQ ID N°36;
- 32 consecutive aminoacids of SEQ ID N°37;
- 30 and
- 22 consecutive aminoacids of SEQ ID N°38.

6. A nucleic acid according to claim 1 encoding a polypeptide having an
 35 amino acid sequence comprising from one to three substitutions,

DA.

Le August 2 2000

additions or deletions of one amino acid as regards a polypeptide selected from the group consisting of the amino acid sequences SEQ ID N°1 to 38 or as regards a polypeptide according to claim 5 and a sequence complementary thereto.

5

7. A polypeptide selected from the group consisting of the amino acid sequences SEQ ID N°1 to 38 or a variant thereof.

10 8. A polypeptide according to claim 7 having at least 95% aminoacid identity with a polypeptide selected from the group consisting of the aminoacid sequences SEQ ID N°1 to 38.

15 9. A polypeptide according to claim 7 having an aminoacid sequence selected from the group consisting of the sequences consisting of at least:

- 45 consecutive aminoacids of SEQ ID N°1;
- 30 consecutive aminoacidss of SEQ ID N°2;
- 65 consecutive aminoacids of SEQ ID N°3;
- 20 - 30 consecutive aminoacids of SEQ ID N°4;
- 130 consecutive aminoacids of SEQ ID N°5;
- 25 consecutive aminoacids of SEQ ID N°6;
- 23 consecutive aminoacids of SEQ ID N°7.
- 48 consecutive aminoacids of SEQ ID N°8;
- 25 - 36 consecutive aminoacids of SEQ ID N°9;
- 25 consecutive aminoacids of SEQ ID N°10;
- 24 consecutive aminoacids of SEQ ID N°11;
- 37 consecutive aminoacids of SEQ ID N°12;
- 25 consecutive aminoacids of SEQ ID N°13;
- 30 - 30 consecutive aminoacids of SEQ ID N°14;
- 27 consecutive aminoacids of SEQ ID N°15;
- 69 consecutive aminoacids of SEQ ID N°16;
- 130 consecutive aminoacids of SEQ ID N°17;

- 33 consecutive aminoacids of SEQ ID N°18;
 - 25 consecutive aminoacids of SEQ ID N°19;
 - 40 consecutive aminoacids of SEQ ID N°20;
 - 78 consecutive aminoacids of SEQ ID N°21;
 - 5 - 39 consecutive aminoacids of SEQ ID N°22;
 - 57 consecutive aminoacids of SEQ ID N°23;
 - 26 consecutive aminoacids of SEQ ID N°24;
 - 68 consecutive aminoacids of SEQ ID N°25;
 - 34 consecutive aminoacids of SEQ ID N°26;
 - 10 - 42 consecutive aminoacids of SEQ ID N°27;
 - 48 consecutive aminoacids of SEQ ID N°28.
 - 102 consecutive aminoacids of SEQ ID N°29:
 - 49 consecutive aminoacids of SEQ ID N°30;
 - 92 consecutive aminoacids of SEQ ID N° 31;
 - 15 - 49 consecutive aminoacids of SEQ ID N°30;
 - 92 consecutive aminoacids of SEQ ID N°31;
 - 71 consecutive aminoacids of SEQ ID N°32;
 - 55 consecutive aminoacids of SEQ ID N°33;
 - 69 consecutive aminoacids of SEQ ID N°34;
 - 20 - 23 consecutive aminoacids of SEQ ID N°35;
 - 33 consecutive aminoacids of SEQ ID N°36;
 - 32 consecutive aminoacids of SEQ ID N°37;
- and
- 22 consecutive aminoacids of SEQ ID N°38.

25

10. A polypeptide according to claim 7 having an amino acid sequence comprising from one to three substitutions, additions or deletions of one amino acid as regards a polypeptide selected from the group consisting of the amino acid sequences SEQ ID N°1 to 38 or as regards a
- 30 polypeptide according to claim 9.

11. An antibody directed against a polypeptide according to any one of claims 7 to 10.

DA.

Le August 2 2000

12. A recombinant vector containing inserted therein a nucleic acid according to any one of claims 1 to 5.

13. The recombinant vector according to claim 12 which is selected from
5 the group consisting of the plasmids pACT11st and pAS2ΔΔ.

14. The recombinant vector according to claim 12 which is selected from the group consisting of pT25, pKT25, pUT18 and pUT18C.

10 15. The recombinant vector according to claim 12 which is selected from the group consisting of pP6 and pB5.

16. A cell host transformed with a vector according to any one of claims 12 to 15 or with a nucleic acid according to anyone of claims 1 to 5.

15

17. A method for producing a polypeptide according to any one of claims 7 to 10, wherein said method comprises the step of :

- a) cultivating a cell host according to claim 18 in an appropriate culture medium;
- 20 b) recovering the recombinant polypeptide from the culture supernatant or from the cell lysate.

18. A yeast two-hybrid system method for selecting a recombinant cell clone containing a vector comprising a nucleic acid insert encoding a
25 prey polypeptide which binds with a SID® polypeptide, wherein said method comprises the steps of :

- a) mating at least one first recombinant yeast cell clone of a collection of recombinant yeast cell clones transformed with a plasmid containing the prey polynucleotide to be assayed with a second haploid recombinant
30 *Saccharomyces cerevisiae* cell clone transformed with a plasmid

containing a bait polynucleotide encoding a SID® polypeptide according to any one of claims 7 to 10;

b) cultivating diploid cells obtained in step a) on a selective medium; and

c) selecting recombinant cell clones which grow on said selective
5 medium.

19. The yeast two-hybrid method of claim 18 which further comprises the step of :

d) characterizing the prey polynucleotide contained in each recombinant
10 cell clone selected in step c).

20. A bacterial two-hybrid method for identifying a recombinant cell clone containing a prey polynucleotide encoding a prey polypeptide which binds with a SID® polypeptide, wherein said method comprises the steps
15 of :

a) transforming bacterial cell clones with a plasmid containing a SID® polynucleotide encoding a SID® polypeptide according to any one of claims 7 to 10;

b) rescuing prey plasmids containing prey polynucleotides wherein each
20 prey polynucleotide is a DNA fragment from the genome of a desired organism and wherein each prey plasmid is contained in one recombinant yeast cell clone of a collection of recombinant yeast cell clones;

c) transforming the recombinant bacterial cell clones obtained in step a)
25 with the plasmids rescued in step b);

d) cultivating bacterial recombinant cells obtained in step c) on a selective medium; and

e) selecting recombinant cell clones which grow on said selective
30 medium.

DA.

Le August 2 2000

21. The bacterial two-hybrid method of claim 20, wherein said method further comprises the step of f) characterising the prey polynucleotide contained in each recombinant cell clone selected at step e).

5 22. The method according to any one of claims 18 to 21, wherein the prey polypeptide is a human polypeptide.

23. The method according to any one of claims 18 to 21, wherein the prey polypeptide is an HCV polypeptide.

10

24. The method of claim 23, wherein the prey polypeptide is encoded by a strain of HCV which is pathogenic for human.

25. A set of two nucleic acids consisting of :

- 15 i) a first nucleic acid encoding a SID® polypeptide according to any one of claims 7 to 10; and
ii) a second nucleic acid encoding a prey polypeptide which binds specifically with the SID® polypeptide defined in i).

20 26. A set of two nucleic acids which is selected from the group consisting of the following sets:

SEQ ID N°77/SEQ ID N°1; SEQ ID N°78/SEQ ID N°2; SEQ ID N°78/SEQ ID N°3; SEQ ID N°79/SEQ ID N°4; SEQ ID N°80/SEQ ID N°5;
25 SEQ ID N°81/SEQ ID N°6; SEQ ID N°82/SEQ ID N°7; SEQ ID N°83/SEQ ID N°8; SEQ ID N°84/SEQ ID N°9; SEQ ID N°85/SEQ ID N°10; SEQ ID N°86/SEQ ID N°11; SEQ ID N°87/SEQ ID N°12; SEQ ID N°88/SEQ ID N°13; SEQ ID N°89/SEQ ID N°14; SEQ ID N°90/SEQ ID N°15; SEQ ID N°91/SEQ ID N°16; SEQ ID N°92/SEQ ID N°17; SEQ ID
30 N°93/SEQ ID N°18; SEQ ID N°94/SEQ ID N°19; SEQ ID N°95/SEQ ID N°20; SEQ ID N°96/SEQ ID N°21; SEQ ID N°97/SEQ ID N°22; SEQ ID

N°98/SEQ ID N°23; SEQ ID N°99/SEQ ID N°24; SEQ ID N°100/SEQ ID N°25. SEQ ID N°101/SEQ ID N°26. SEQ ID N°102/SEQ ID N°27; SEQ ID N°103/SEQ ID N°28. SEQ ID N°104/SEQ ID N°29; SEQ ID N°105/SEQ ID N°30; SEQ ID N°106/SEQ ID N°31; SEQ ID N°107/SEQ ID N°32; SEQ ID N°108/SEQ ID N°33; SEQ ID N°109/SEQ ID N°34; SEQ ID N°110/SEQ ID N°35; SEQ ID N°111/SEQ ID N°36; SEQ ID N°112/SEQ ID N°37; and SEQ ID N°113/SEQ ID N°38.

27. A set of two polypeptides consisting of :

- i) a first polypeptide consisting of a SID@ polypeptide according to any one of claims 7 to 10; and
- ii) a second polypeptide, also termed prey polypeptide, which binds specifically with the first polypeptide.

28. A set of two polypeptides which is selected from the group consisting of the following sets:

SEQ ID N°114/SEQ ID N°39; SEQ ID N°115/SEQ ID N°40; SEQ ID N°115/SEQ ID N°41; SEQ ID N°116/SEQ ID N°42; SEQ ID N°117/SEQ ID N°43; SEQ ID N°118/SEQ ID N°44; SEQ ID N°119/SEQ ID N°45; SEQ ID N°120/SEQ ID N°46; SEQ ID N°121/SEQ ID N°47; SEQ ID N°122/SEQ ID N°48; SEQ ID N°123/SEQ ID N°49; SEQ ID N°124/SEQ ID N°50; SEQ ID N°125/SEQ ID N°51; SEQ ID N°126/SEQ ID N°52; SEQ ID N°127/SEQ ID N°53; SEQ ID N°128/SEQ ID N°54; SEQ ID N°129/SEQ ID N°55; SEQ ID N°130/SEQ ID N°56; SEQ ID N°131/SEQ ID N°57; SEQ ID N°132/SEQ ID N°58; SEQ ID N°133/SEQ ID N°59; SEQ ID N°134/SEQ ID N°60; SEQ ID N°135/SEQ ID N°61; SEQ ID N°136/SEQ ID N°62; SEQ ID N°137/SEQ ID N°63; SEQ ID N°138/SEQ ID N°64; SEQ ID N°139/SEQ ID N°65; SEQ ID N°140/SEQ ID N°66; SEQ ID N°141/SEQ ID N°67; SEQ ID N°142/SEQ ID N°68; SEQ ID N°143/SEQ ID N°69; SEQ ID N°144/SEQ ID N°70. SEQ ID N°145/SEQ ID N°71; SEQ ID N°146/SEQ ID N°72. SEQ ID N°147/SEQ

DA.

Le August 2 2000

ID N°73; SEQ ID N°148/SEQ ID N°74; SEQ ID N°149/SEQ ID N°75 and SEQ ID N°150/SEQ ID N°76.

296. A complex formed between the two polypeptides of claim 29 or 30.

5

30. A method for selecting a molecule which inhibits the binding between a set of two polypeptides according to claim 27 or 28, wherein said method comprises the steps of :

a) cultivating a recombinant host cell containing a reporter gene the
10 expression of which is toxic for said recombinant host cell , said host cell being transformed with two vectors wherein :

i) the first vector contains a nucleic acid comprising a polynucleotide encoding a first hybrid polypeptide containing one of said two polypeptides and a DNA binding domain;

15 ii) the second vector contains a nucleic acid comprising a polynucleotide encoding a second hybrid polypeptide containing the second of said two polypeptides and an activating domain capable of activating said toxic reporter gene when the first and the second hybrid polypeptides are interacting;

20 on a selective medium containing the molecule to be tested and allowing the growth of said recombinant host cell when the toxic reporter gene is not activated; and

b) selecting the molecule which inhibits the growth of the recombinant host cell defined in step a).

25

31. A method for selecting a molecule which inhibits the protein-protein interaction of a set of two polypeptides according to claim 29 or 30, wherein said method comprises the steps of :

a) cultivating a recombinant host cell containing a reporter gene the
30 expression of which is toxic for said recombinant host cell, said host cell being transformed with two vectors wherein :

- i) the first vector contains a nucleic acid comprising a polynucleotide encoding a first hybrid polypeptide containing one of said set of two polypeptides and the first domain of an enzyme;
- ii) the second vector contains a nucleic acid comprising a polynucleotide encoding a second hybrid polypeptide containing the second of said two polypeptides and the second part of said enzyme capable of activating said toxic reporter gene when the first and the second hybrid polypeptides are interacting, said interaction recovering the catalytic activity of the enzyme;
- on a selective medium containing the molecule to be tested and allowing the growth of said recombinant host cell when the toxic gene is not activated; and
- b) selecting the molecule which inhibits the growth of the recombinant host cell defined in step a).

15

32. A kit for the screening of a molecule which inhibits the protein-protein interaction of a set of two polypeptides according to claim 27 or 28, wherein said kit comprises a recombinant cell host containing a reporter gene the expression of which is toxic for said recombinant cell host, said cell host being transformed with two vectors wherein :

20

- i) the first vector contains a nucleic acid comprising a polynucleotide encoding a first hybrid polypeptide containing one of said two polypeptides and a DNA binding domain;
- ii) the second vector contains a nucleic acid comprising a polynucleotide encoding a second hybrid polypeptide containing the second of said two polypeptides and an activating domain capable of activating said toxic reporter gene when the first and the second hybrid polypeptides are interacting

25

33 A kit for the screening of a molecule which inhibits the protein-protein interaction of a set of two polypeptides according to claim 27 or 28,

30

DA.

Le August 2 2000

wherein said kit comprises a recombinant host cell containing a reporter gene the expression of which is toxic for said recombinant host cell, said host cell being transformed with two plasmids wherein :

- i) the first vector contains a nucleic acid comprising a polynucleotide
5 encoding a first hybrid polypeptide containing one of said two polypeptides and the first domain of an enzyme;
- ii) the second plasmid contains a nucleic acid comprising a polynucleotide encoding a second hybrid polypeptide containing the second of said two polypeptides and the second part of said enzyme
10 capable of activating said toxic reporter gene when the first and the second hybrid polypeptides are interacting, said interaction recovering the catalytic activity of the enzyme.

34. A marker compound wherein said compound comprises :

- 15 a) a Selected Interacting Domain (SID®) polypeptide according to any one of claims 7 to 10 or a variant thereof; and
- b) a detectable molecule bound thereto.

35. The marker compound of claim 34, wherein the detectable molecule
20 consists of a fluorescent protein.

36. The marker compound of claim 35, wherein the detectable protein is selected from the group consisting of GFP and YFP.

25 37. The marker compound of claim 35, wherein the detectable molecule is endowed with a catalytic activity.

38. The marker compound of claim 37, wherein the detectable molecule is selected from the group consisting of a hydrolase, a transferase, a
30 lyase, an isomerase, a ligase, a synthetase and a oxidoreductase.

39. The marker compound of claim 34, wherein the detectable molecule is radioactive.

40. The marker compound of claim 34, wherein the detectable molecule is chemiluminescent.

5 41. The marker compound according to any one of claims 34 to 40, wherein the detectable molecule is covalently bound to the Selected Interacting Domain (SID®) polypeptide or a variant thereof.

10 42. The marker compound according to any one of claims 34 to 40, wherein the detectable molecule is non covalently bound to the Selected Interacting Domain (SID®) polypeptide or a variant thereof.

15 43. The marker compound of claim 42, wherein the detectable molecule is an antibody directed specifically against the Selected Interacting Domain (SID®) polypeptide.

44 The marker compound of claim 43, wherein said antibody is labelled radioactively or non radioactively.

20 45. The marker compound according to claim 34, wherein :
a) the Selected Interacting Domain (SID®) polypeptide or a variant thereof is covalently bound to a first ligand.; and
b) the detectable molecule comprises a second ligand which binds specifically to the first ligand.

25 46. The marker compound according to claim 45, wherein the first ligand is biotin and the second ligand is streptavidin.

30 47 A nucleic acid encoding a marker compound according to any one of claims 34 to 41.

48. A nucleic acid encoding the Selected Interacting Domain (SID®) polypeptide or a variant thereof onto which is covalently bound a first ligand defined in claims 45 and 46.

35

DA.
Le August 2 2000

496. A recombinant vector comprising inserted therein a nucleic acid according to any one of claims 47 and 48.

50. The recombinant vector according to claim 48, which is selected from
5 the group consisting of pACT11st, pASΔΔ, pT25, pKT25, pUT18, pUT18C, pP6 and pB5..

51. A recombinant host cell which has been transfected with a nucleic acid according to any one of claims 47 and 48 or a recombinant vector
10 according to any one of claims 43 and 44.

52. The recombinant host cell according to claim 51 which is of prokaryotic origin.

15 53. The recombinant host cell according to claim 51 which is of eukaryotic origin.

54. The recombinant host cell according to claim 52 which is a mammalian host cell.

20 53 A method for detecting a polypeptide of interest within a sample, which comprises the steps of :

- a) contacting a marker compound or a plurality of marker compounds according to any one of claims 34 to 46 with the sample;
- 25 b) detecting the complexes formed between said marker compound or said plurality of marker compounds and said polypeptide of interest.

56. A kit for detecting a polypeptide of interest within a sample, which comprises a marker compound according to any one of claims 34 to 46.
30

57. A method for detecting a polypeptide of interest within a prokaryotic or an eukaryotic host cell, said method comprising the steps of :
a) providing a cell host to be assayed;

- b) transfecting said host cell with a nucleic acid according to any one of claims 41 and 42 or with a recombinant vector according to any one of claims 496 and 50;
- c) detecting the complexes formed between the marker compound
5 expressed by the transfected cell host and the polypeptide of interest.

58. A kit for detecting a polypeptide of interest within a prokaryotic or an eukaryotic host cell which comprises a nucleic acid according to any one of claims 47 and 48 or a recombinant vector according to any one of
10 claims 496 and 50.

59. A method for detecting a polypeptide of interest within a prokaryotic or eukaryotic host cell, said method comprising the steps of :
- a) providing a cell host to be assayed;
 - 15 b) introducing a marker compound according to any one of claims 36 to 48 within said cell host;
 - c) detecting the complexes formed between the marker compound and the polypeptide of interest within the cell.

20 60. A kit for detecting a polypeptide of interest within a prokaryotic or eukaryotic host cell comprising a marker compound according to any one of claims 34 to 46.

61. A method for detecting a polypeptide or a plurality of polypeptides of
25 interest within a sample, wherein said method comprises the steps of :
- a) providing a substrate onto which a Selected Interacting Domain (SID®) polypeptide according to any one of claims 7 to 10 or a variant thereof, or a plurality of Selected Interacting Domain (SID®) polypeptides according to any one of claims 7 to 10 or variants thereof
30 is (are) immobilised;
 - b) bringing into contact the substrate defined in a) with the sample to be assayed;
 - c) detecting the complexes formed between the Selected Interacting Domain (SID®) polypeptides or variants thereof or a variant thereof, or

the plurality of Selected Interacting Domain (SID®) polypeptide and a molecule or a plurality of molecules initially contained in the sample;

62. The method of claim 61, wherein a plurality of Selected Interacting
5 Domain (SID®) polypeptides or variants thereof are immobilised on the substrate in an ordered manner.

63. The method of claim 61, wherein the Selected Interacting Domain
10 (SID®) polypeptide or a variant thereof, or the plurality of Selected Interacting Domain (SID®) polypeptides or variants thereof are covalently bound to the substrate.

64. The method of claim 61, wherein the Selected Interacting Domain
15 (SID®) polypeptide or a variant thereof, or the plurality of Selected Interacting Domain (SID®) polypeptides or a variant thereof are non-covalently bound to the substrate.

65. The method of claim 61, wherein the Selected Interacting Domain
20 (SID®) polypeptide or a variant thereof, or the plurality of Selected Interacting Domain (SID®) polypeptides or variants thereof are covalently bound to a first ligand and wherein the substrate is coated with a second ligand which specifically binds to the first ligand.

66. The method of claim 61, wherein the first ligand is biotin and the
25 second ligand is streptavidin.

67. The method according to any one of claims 61 to 66, wherein the
30 Selected Interacting Domain (SID®) polypeptide or a variant thereof, or the plurality of Selected Interacting Domain (SID®) polypeptides or variants thereof are covalently linked to a spacer and wherein said spacer is covalently bound to the substrate in order to immobilise the Selected Interacting Domain (SID®) polypeptide, or a variant thereof or the plurality of Selected Interacting Domain (SID®) polypeptides.

68.. The method according to any one of claims 61 to 66 wherein the detection step c) consists of detecting changes in optical characteristics of the substrate.

5 69. A device for the detection of a polypeptide or a plurality of polypeptides of interest within a sample, wherein said device comprises a substrate onto which a Selected Interacting Domain (SID®) polypeptide according to any one of claims 7 to 10 or a variant thereof or a plurality of Selected Interacting Domain (SID®) polypeptides according to any one of claims 7 to 10 or variants thereof is (are) immobilised.

70. A pharmaceutical composition comprising a pharmaceutically effective amount of a nucleic acid comprising a polynucleotide encoding a Selected Interacting Domain (SID®) polypeptide according to any one of claims 7 to 10 or a variant thereof.

71. A method for preventing or curing a viral infection by a Hepatitis C virus in a human or an animal, wherein said method comprises a step of administering to the human or animal body a pharmaceutically effective amount of a Selected Interacting Domain (SID®) polypeptide according to any one of claims 7 to 10.

72. A method for preventing or curing a viral infection by a Hepatitis C virus in a human or an animal, wherein said method comprises a step of administering to the human or animal body a pharmaceutically effective amount of a nucleic acid comprising a polynucleotide encoding a Selected Interacting Domain (SID®) polypeptide according to any one of claims 7 to 10, and wherein said polynucleotide is placed under the control of regulatory sequence which is functional in said human or said animal.

73. A method for preventing or curing a viral or a bacterial infection in a human or an animal, wherein said method comprises a step of administering to the human or animal body a pharmaceutically effective amount of a recombinant expression vector comprising a polynucleotide

DA.

Le August 2 2000

encoding a Selected Interacting Domain (SID®) polypeptide according to any one of claims 7 to 10.

74. A method for selecting a SID® polypeptide comprising the steps of :

5 1) Selecting a collection of nucleic acids (prey nucleic acids) which bind specifically to a given bait polypeptide of interest; and

 2) determining the nucleic acid sequences which encode for SID® polypeptides after having generated sets of polynucleotides from
10 the collection of nucleic acids selected at step 1).

75. The method of claim 74, wherein step 1) consists of a yeast two-hybrid method or a bacterial two-hybrid method.

15 76. The method of claim 74, wherein step 2) comprises the following steps of :

 a) selecting from the collection of prey polynucleotides obtained at the end of step 1) all prey polynucleotides encoding a prey polypeptide capable of interacting with said bait polypeptide and containing a
20 common nucleic acid fragment;

 b) aligning the nucleotide sequences of the prey polynucleotides selected at step a) and gathering in one set or in a plurality of sets of sequences those nucleotide sequences which have sequences that overlap for more than 30% of their respective nucleic
25 acid length, wherein each common overlapping nucleotide sequence in one set of sequences defines a sequence encoding a pre-SID® polypeptide ; and

 c) aligning two sequences encoding two respective pre-SID® polypeptides, and:

30 (i) defining an overlapping nucleic acid sequence between the sequences encoding the two respective pre-SID® polypeptides as a sequence encoding a SID® polypeptide, provided that the overlapping sequence is of at least 30 nucleotides in length;

 ii) defining a non-overlapping nucleic acid sequence between
35 the sequences encoding the two respective pre-SID® polypeptides as a

sequence encoding a SID® polypeptide, provided that (1) said non-overlapping sequence has more than 30 nucleotides in length and (2) said non-overlapping sequence represents at least 30% in length of any one of the polynucleotides contained in the set of prey polynucleotides used for defining the sequence encoding each pre-SID® polypeptide.

77. The method of claim 76 wherein step 2) further comprises the steps of :

d) counting the number of overlapping prey polynucleotides contained in a first set of polynucleotides defining a sequence encoding a first SID® polypeptide;

e) counting the number of overlapping prey polynucleotides contained in a second set of polynucleotides defining a sequence encoding a second SID® polypeptide which overlaps with the sequence encoding the first SID® polypeptide;

f) determining which sequence among those encoding respectively the first SID® polypeptide and the second SID® polypeptide has been defined with the largest number of prey polynucleotides and selecting this set of prey sequences;

g) adding to the set of prey sequences selected at step f) those sequences that were contained in the set of prey sequences used for defining the sequence encoding the SID® polypeptide with the smallest number of prey sequences and which overlap with the sequence encoding the SID® polypeptide with the largest number of prey sequences;

h) aligning the prey sequences added at step g) with the sequences already contained in the set of prey sequences which defined the sequence encoding the SID® polypeptide with the largest number of prey sequences;

i) defining an overlapping sequence between the whole sequences which were aligned in step h), wherein said overlapping sequence consists of a sequence encoding a SID® polypeptide.

DA.
Le August 2 2000

78. The method according to any one of claims 74 to 76, wherein the collection of prey nucleic acids is prepared starting from the genomic DNA of an organism containing contiguous Open Reading Frames.

5 79. The method according to claim 78, wherein said organism is a virus.

80. The method according to claim 79, wherein the virus consists of the Hepatitis C virus.

10 81. The method according to claim 80, wherein the Hepatitis C virus is pathogenic for a mammal, including human.

82. A SID® nucleic acid selected according to the method of any one of claims 74 to 81.

15

83. A SID® polypeptide encoded by a nucleic acid according to claim 82.

20

THIS PAGE BLANK (USPTO)

ABSTRACT

**SID® NUCLEIC ACIDS AND POLYPEPTIDES
SELECTED FROM A PATHOGENIC STRAIN OF
HEPATITIS C VIRUS AND APPLICATIONS
THEREOF**

The present invention relates to nucleic acids encoding SID® polypeptides which bind selectively to a polypeptide encoded by a pathogenic strain of the hepatitis C virus, as well as to the SID® polypeptides which are encoded by said nucleic acids.

The invention also concerns vectors comprising a nucleic acid encoding a SID® polypeptide as well as host cells transformed with such vectors.

The invention is also directed to two-hybrid methods which make use of the nucleic acids encoding a SID® polypeptide selected from a pathogenic strain of the hepatitis C virus as well as to methods for selecting molecules which inhibit the binding between a SID® polypeptide and a polypeptide which specifically binds thereto.

THIS PAGE BLANK (USPTO)

1/11




FIGURE 1

2/11




FIGURE 2

3/11





FIGURE 3

4/11



Lac Z⁺

ATG ACC ATG ATT ACG CCA AGC TTG CAT GCC TGC AGC TCG ACT CTA GAG GAT CCC CGG GTA -CCG AGC TCG AAT TCA
TAC TGG TAC TAA TGC GGT TCG AAC GTA CCG AGC TCC AGC TCA GAT CTC CTA GCG GGC CAT GGC TCG AGC TTA ACT
MET THR MET ILE THR PRO SER LEU HIS ALA CYS ARG SER THR LEU GLU ASP PRO ARG VAL PRO SER SER ASN SER






FIGURE 4

PstI SalI XbaI BamHI SmaI KpnI SacI EcoRI ClaI

GCC TGC AGG TCG ACT CTA GAG GAT CCC GCG GTA CCG AGC TCG AAT TCA TGC ATA TAA
CCG ACG TCC AGC TCA GAT CTC CTA GCG GGC CAT GGC TCG AGC TTA ACT ACG TAT ATT
ATA CYS ARG SER THR LEU GLU ASP PRO ARG VAL PRO SER SER ASN SER ILE STOP


FIGURE 5

5/11



(*) Restriction sites are not unique

FIGURE 6



(*) Restriction site is not unique

FIGURE 7

6/11




FIGURE 8

7/11




FIGURE 9




FIGURE 10

9/11




FIGURE 11

10/11




FIGURE 12

11/11




FIGURE 13

THIS PAGE BLANK (USPTO)

SEQUENCE LISTING

<110> HYBRIGENICS S.A.

<120> SID nucleic acids and polypeptides selected from a
pathogenic strain of the hepatitis C virus and
applications

<130> Hybrigenics - SID HCV

<140>

<141>

<160> 156

<170> PatentIn Ver. 2.1

<210> 1

<211> 50

<212> PRT

<213> Hepatitis C virus

<400> 1

Leu Leu Pro Arg Arg Gly Pro Arg Leu Gly Val Arg Ala Thr Arg Lys
1 5 10 15

Thr Ser Glu Arg Ser Gln Pro Arg Gly Arg Arg Gln Pro Ile Pro Lys
20 25 30

Ala Arg Arg Pro Glu Gly Arg Thr Trp Ala Gln Pro Gly Tyr Pro Trp
35 40 45

Pro Leu
50

<210> 2

<211> 35

<212> PRT

<213> Hepatitis C virus

<400> 2

Gly Arg Gly Lys Pro Gly Ile Tyr Arg Phe Val Ala Pro Gly Glu Arg
1 5 10 15

Pro Ser Gly Met Phe Asp Ser Ser Val Leu Cys Glu Cys Tyr Asp Ala
20 25 30

Gly Cys Ala
35

<210> 3
 <211> 77
 <212> PRT
 <213> Hepatitis C virus

<400> 3
 Asn Thr Asn Arg Arg Pro Gln Asp Val Lys Phe Pro Gly Gly Gly Gln
 1 5 10 15
 Ile Val Gly Gly Val Tyr Leu Leu Pro Arg Arg Gly Pro Arg Leu Gly
 20 25 30
 Val Arg Ala Thr Arg Lys Thr Ser Glu Arg Ser Gln Pro Arg Gly Arg
 35 40 45
 Arg Gln Pro Ile Pro Lys Ala Arg Arg Pro Glu Gly Arg Thr Trp Ala
 50 55 60
 Gln Pro Gly Tyr Pro Trp Pro Leu Tyr Gly Asn Glu Gly
 65 70 75

<210> 4
 <211> 37
 <212> PRT
 <213> Hepatitis C virus

<400> 4
 Pro Ser Pro Val Val Val Gly Thr Thr Asp Arg Ser Gly Ala Pro Thr
 1 5 10 15
 Tyr Ser Trp Gly Ala Asn Asp Thr Asp Val Phe Val Leu Asn Asn Thr
 20 25 30
 Arg Pro Pro Leu Gly
 35

<210> 5
 <211> 150
 <212> PRT
 <213> Hepatitis C virus

<400> 5
 Ser Arg Thr Gln Arg Arg Gly Arg Thr Gly Arg Gly Lys Pro Gly Ile
 1 5 10 15
 Tyr Arg Phe Val Ala Pro Gly Glu Arg Pro Ser Gly Met Phe Asp Ser
 20 25 30
 Ser Val Leu Cys Glu Cys Tyr Asp Ala Gly Cys Ala Trp Tyr Glu Leu
 35 40 45

Thr Pro Ala Glu Thr Thr Val Arg Leu Arg Ala Tyr Met Asn Thr Pro
 50 55 60
 Gly Leu Pro Val Cys Gln Asp His Leu Glu Phe Trp Glu Gly Val Phe
 65 70 75 80
 Thr Gly Leu Thr His Ile Asp Ala His Phe Leu Ser Gln Thr Lys Gln
 85 90 95
 Ser Gly Glu Asn Phe Pro Tyr Leu Val Ala Tyr Gln Ala Thr Val Cys
 100 105 110
 Ala Arg Ala Gln Ala Pro Pro Pro Ser Trp Asp Gln Met Trp Lys Cys
 115 120 125
 Leu Ile Arg Leu Lys Pro Thr Leu His Gly Pro Thr Pro Leu Leu Tyr
 130 135 140
 Arg Leu Gly Ala Val Gln
 145 150

<210> 6
 <211> 28
 <212> PRT
 <213> Hepatitis C virus

<400> 6
 Pro Ser Pro Val Val Val Gly Thr Thr Asp Arg Ser Gly Ala Pro Thr
 1 5 10 15
 Tyr Ser Trp Gly Ala Asn Asp Thr Asp Val Phe Val
 20 25

<210> 7
 <211> 26
 <212> PRT
 <213> Hepatitis C virus

<400> 7
 Pro Pro Arg Pro Cys Gly Ile Val Pro Ala Lys Ser Val Cys Gly Pro
 1 5 10 15
 Val Tyr Cys Phe Thr Pro Ser Pro Val Val
 20 25

<210> 8
 <211> 54

<212> PRT

<213> Hepatitis C virus

<400> 8

Cys Val Val Ile Val Gly Arg Ile Val Leu Ser Gly Lys Pro Ala Ile
1 5 10 15

Ile Pro Asp Arg Glu Val Leu Tyr Gln Glu Phe Asp Glu Met Glu Glu
20 25 30

Cys Ser Gln His Leu Pro Tyr Ile Glu Gln Gly Met Met Leu Ala Glu
35 40 45

Gln Phe Lys Gln Lys Ala
50

<210> 9

<211> 40

<212> PRT

<213> Hepatitis C virus

<400> 9

Gly Asp Phe Asp Ser Val Ile Asp Cys Asn Thr Cys Val Thr Gln Thr
1 5 10 15

Val Asp Phe Ser Leu Asp Pro Thr Phe Thr Ile Glu Thr Thr Thr Leu
20 25 30

Pro Gln Asp Ala Val Ser Arg Thr
35 40

<210> 10

<211> 28

<212> PRT

<213> Hepatitis C virus

<400> 10

Glu Arg Pro Ser Gly Met Phe Asp Ser Ser Val Leu Cys Glu Cys Tyr
1 5 10 15

Asp Ala Gly Cys Ala Trp Tyr Glu Leu Thr Pro Ala
20 25

<210> 11

<211> 27

<212> PRT

<213> Hepatitis C virus

<400> 11

Arg Gly Lys Pro Gly Ile Tyr Arg Phe Val Ala Pro Gly Glu Arg Pro
1 5 10 15

Ser Gly Met Phe Asp Ser Ser Val Leu Cys Glu
20 25

<210> 12

<211> 42

<212> PRT

<213> Hepatitis C virus

<400> 12

Leu Glu Asp Ser Val Thr Pro Ile Asp Thr Thr Ile Met Ala Lys Asn
1 5 10 15

Glu Val Phe Cys Val Gln Pro Glu Lys Gly Gly Arg Lys Pro Ala Arg
20 25 30

Leu Ile Val Phe Pro Asp Leu Gly Val Arg
35 40

<210> 13

<211> 28

<212> PRT

<213> Hepatitis C virus

<400> 13

Pro Thr Gly Ser Gly Lys Ser Thr Lys Val Pro Ala Ala Tyr Ala Ala
1 5 10 15

Gln Gly Tyr Lys Val Leu Val Leu Asn Pro Ser Val
20 25

<210> 14

<211> 33

<212> PRT

<213> Hepatitis C virus

<400> 14

Glu Arg Pro Tyr Cys Trp His Tyr Pro Pro Arg Pro Cys Gly Ile Val
1 5 10 15

Pro Ala Lys Ser Val Cys Gly Pro Val Tyr Cys Phe Thr Pro Ser Pro
20 25 30

Val

<210> 15
 <211> 31
 <212> PRT
 <213> Hepatitis C virus

<400> 15
 Pro Ser Pro Val Val Val Gly Thr Thr Asp Arg Ser Gly Ala Pro Thr
 1 5 10 15
 Tyr Ser Trp Gly Ala Asn Asp Thr Asp Val Phe Val Leu Asn Asn
 20 25 30

<210> 16
 <211> 77
 <212> PRT
 <213> Hepatitis C virus

<400> 16
 Ala Gly Ala Leu Val Ala Phe Lys Ile Met Ser Gly Glu Val Pro Ser
 1 5 10 15
 Thr Glu Asp Leu Val Asn Leu Leu Pro Ala Ile Leu Ser Pro Gly Ala
 20 25 30
 Leu Val Val Gly Val Val Cys Ala Ala Ile Leu Arg Arg His Val Gly
 35 40 45
 Pro Gly Glu Gly Ala Val Gln Trp Met Asn Arg Leu Ile Ala Phe Ala
 50 55 60
 Ser Arg Gly Asn His Val Ser Pro Thr His Tyr Val Pro
 65 70 75

<210> 17
 <211> 147
 <212> PRT
 <213> Hepatitis C virus

<400> 17
 Glu Val Gln Ile Val Ser Thr Ala Thr Gln Thr Phe Leu Ala Thr Cys
 1 5 10 15
 Ile Asn Gly Val Cys Trp Thr Val Tyr His Gly Ala Gly Thr Arg Thr
 20 25 30
 Ile Ala Ser Pro Lys Gly Pro Val Ile Gln Met Tyr Thr Asn Val Asp
 35 40 45

Gln Asp Leu Val Gly Trp Pro Ala Pro Gln Gly Ser Arg Ser Leu Thr
 50 55 60
 Pro Cys Thr Cys Gly Ser Ser Asp Leu Tyr Leu Val Thr Arg His Ala
 65 70 75 80
 Asp Val Ile Pro Val Arg Arg Arg Gly Asp Ser Arg Gly Ser Leu Leu
 85 90 95
 Ser Pro Arg Pro Ile Ser Tyr Leu Lys Gly Ser Ser Gly Gly Pro Leu
 100 105 110
 Leu Cys Pro Ala Gly His Ala Val Gly Leu Phe Arg Ala Ala Val Cys
 115 120 125
 Thr Arg Gly Val Ala Lys Ala Val Asp Phe Ile Pro Val Glu Asn Leu
 130 135 140
 Gly Thr Thr
 145

<210> 18
 <211> 36
 <212> PRT
 <213> Hepatitis C virus

<400> 18
 Val Thr Gln Leu Leu Arg Arg Leu His Gln Trp Ile Ser Ser Glu Cys
 1 5 10 15
 Thr Thr Pro Cys Ser Gly Ser Trp Leu Arg Asp Ile Trp Asp Trp Ile
 20 25 30
 Cys Glu Val Leu
 35

<210> 19
 <211> 28
 <212> PRT
 <213> Hepatitis C virus

<400> 19
 Val Cys Gly Pro Val Tyr Cys Phe Thr Pro Ser Pro Val Val Val Gly
 1 5 10 15
 Thr Thr Asp Arg Ser Gly Ala Pro Thr Tyr Ser Trp
 20 25

<210> 20
 <211> 45
 <212> PRT
 <213> Hepatitis C virus

<400> 20
 Pro Pro Leu Arg Ala Trp Arg His Arg Ala Arg Ser Val Arg Ala Arg
 1 5 10 15
 Leu Leu Ser Arg Gly Gly Arg Ala Ala Ile Cys Gly Lys Tyr Leu Phe
 20 25 30
 Asn Trp Ala Val Arg Thr Lys Leu Lys Leu Thr Pro Ile
 35 40 45

<210> 21
 <211> 86
 <212> PRT
 <213> Hepatitis C virus

<400> 21
 Thr Ala Phe Val Gly Ala Gly Leu Ala Gly Ala Ala Ile Gly Ser Val
 1 5 10 15
 Gly Leu Gly Lys Val Leu Val Asp Ile Leu Ala Gly Tyr Gly Ala Gly
 20 25 30
 Val Ala Gly Ala Leu Val Ala Phe Lys Ile Met Ser Gly Glu Val Pro
 35 40 45
 Ser Thr Glu Asp Leu Val Asn Leu Leu Pro Ala Ile Leu Ser Pro Gly
 50 55 60
 Ala Leu Val Val Gly Val Val Cys Ala Ala Ile Leu Arg Arg His Val
 65 70 75 80
 Gly Pro Gly Glu Gly Ala
 85

<210> 22
 <211> 43
 <212> PRT
 <213> Hepatitis C virus

<400> 22
 Gly Ile Val Pro Ala Lys Ser Val Cys Gly Pro Val Tyr Cys Phe Thr
 1 5 10 15
 Pro Ser Pro Val Val Val Gly Thr Thr Asp Arg Ser Gly Ala Pro Thr

20 25 30
 Tyr Ser Trp Gly Ala Asn Asp Thr Asp Val Phe
 35 40

<210> 23
 <211> 63
 <212> PRT
 <213> Hepatitis C virus

<400> 23
 Val Leu Val Asp Ile Leu Ala Gly Tyr Gly Ala Gly Val Ala Gly Ala
 1 5 10 15
 Leu Val Ala Phe Lys Ile Met Ser Gly Glu Val Pro Ser Thr Glu Asp
 20 25 30
 Leu Val Asn Leu Leu Pro Ala Ile Leu Ser Pro Gly Ala Leu Val Val
 35 40 45
 Gly Val Val Cys Ala Ala Ile Leu Arg Arg His Val Gly Pro Gly
 50 55 60

<210> 24
 <211> 29
 <212> PRT
 <213> Hepatitis C virus

<400> 24
 Glu Arg Pro Tyr Cys Trp His Tyr Pro Pro Arg Pro Cys Gly Ile Val
 1 5 10 15
 Pro Ala Lys Ser Val Cys Gly Pro Val Tyr Cys Phe Thr
 20 25

<210> 25
 <211> 76
 <212> PRT
 <213> Hepatitis C virus

<400> 25
 Arg Arg His Trp Thr Thr Gln Asp Cys Asn Cys Ser Ile Tyr Pro Gly
 1 5 10 15
 His Ile Thr Gly His Arg Met Ala Trp Asp Met Met Met Asn Trp Ser
 20 25 30
 Pro Thr Ala Ala Leu Val Val Ala Gln Leu Leu Arg Ile Pro Gln Ala

10

35 40 45
Ile Met Asp Met Ile Ala Gly Ala His Trp Gly Val Leu Ala Gly Ile
50 55 60
Ala Tyr Phe Ser Met Val Gly Asn Trp Ala Lys Val
65 70 75

<210> 26
<211> 37
<212> PRT
<213> Hepatitis C virus

<400> 26
Ala Ile Leu Ser Ser Leu Thr Val Thr Gln Leu Leu Arg Arg Leu His
1 5 10 15
Gln Trp Ile Ser Ser Glu Cys Thr Thr Pro Cys Ser Gly Ser Trp Leu
20 25 30
Arg Asp Ile Trp Asp
35

<210> 27
<211> 47
<212> PRT
<213> Hepatitis C virus

<400> 27
Val Ser Arg Thr Gln Arg Arg Gly Arg Thr Gly Arg Gly Lys Pro Gly
1 5 10 15
Ile Tyr Arg Phe Val Ala Pro Gly Glu Arg Pro Ser Gly Met Phe Asp
20 25 30
Ser Ser Val Leu Cys Glu Cys Tyr Asp Ala Gly Cys Ala Trp Tyr
35 40 45

<210> 28
<211> 53
<212> PRT
<213> Hepatitis C virus

<400> 28
Leu Pro Ala Pro Asn Tyr Lys Phe Ala Leu Trp Arg Val Ser Ala Glu
1 5 10 15
Glu Tyr Val Glu Ile Arg Arg Val Gly Asp Phe His Tyr Val Ser Gly

20 25 30
 Met Thr Thr Asp Asn Leu Lys Cys Pro Cys Gln Ile Pro Ser Pro Glu
 35 40 45

Phe Phe Thr Glu Leu
 50

<210> 29
 <211> 112
 <212> PRT
 <213> Hepatitis C virus

<400> 29
 Gly Glu Ile Pro Phe Tyr Gly Lys Ala Ile Pro Leu Glu Val Ile Lys
 1 5 10 15

Gly Gly Arg His Leu Ile Phe Cys His Ser Lys Lys Lys Cys Asp Glu
 20 25 30

Leu Ala Ala Lys Leu Val Ala Leu Gly Ile Asn Ala Val Ala Tyr Tyr
 35 40 45

Arg Gly Leu Asp Val Ser Val Ile Pro Thr Ser Gly Asp Val Val Val
 50 55 60

Val Ser Thr Asp Ala Leu Met Thr Gly Phe Thr Gly Asp Phe Asp Ser
 65 70 75 80

Val Ile Asp Cys Asn Thr Cys Val Thr Gln Thr Val Asp Phe Ser Leu
 85 90 95

Asp Pro Thr Phe Thr Ile Glu Thr Thr Thr Leu Pro Gln Asp Ala Val
 100 105 110

<210> 30
 <211> 54
 <212> PRT
 <213> Hepatitis C virus

<400> 30
 Val Cys Ala Ala Ile Leu Arg Arg His Val Gly Pro Gly Glu Gly Ala
 1 5 10 15

Val Gln Trp Met Asn Arg Leu Ile Ala Phe Ala Ser Arg Gly Asn His
 20 25 30

Val Ser Pro Thr His Tyr Val Pro Glu Ser Asp Ala Ala Ala Arg Val
 35 40 45

Thr Ala Ile Leu Ser Ser
 50

<210> 31
 <211> 102
 <212> PRT
 <213> Hepatitis C virus

<400> 31
 Ala Ile Lys Ser Leu Thr Glu Arg Leu Tyr Val Gly Gly Pro Leu Thr
 1 5 10 15

Asn Ser Arg Gly Glu Asn Cys Gly Tyr Arg Arg Cys Arg Ala Ser Gly
 20 25 30

Val Leu Thr Thr Ser Cys Gly Asn Thr Leu Thr Cys Tyr Ile Lys Ala
 35 40 45

Arg Ala Ala Cys Arg Ala Ala Gly Leu Gln Asp Cys Thr Met Leu Val
 50 55 60

Cys Gly Asp Asp Leu Val Val Ile Cys Glu Ser Ala Gly Val Gln Glu
 65 70 75 80

Asp Ala Ala Ser Leu Arg Ala Phe Thr Glu Ala Met Thr Arg Tyr Ser
 85 90 95

Ala Pro Pro Gly Asp Pro
 100

<210> 32
 <211> 79
 <212> PRT
 <213> Hepatitis C virus

<400> 32
 Leu Gln Val Leu Asp Ser His Tyr Gln Asp Val Leu Lys Glu Val Lys
 1 5 10 15

Ala Ala Ala Ser Lys Val Lys Ala Asn Leu Leu Ser Val Glu Glu Ala
 20 25 30

Cys Ser Leu Thr Pro Pro His Ser Ala Lys Ser Lys Phe Gly Tyr Gly
 35 40 45

Ala Lys Asp Val Arg Cys His Ala Arg Lys Ala Val Ala His Ile Asn
 50 55 60

Ser Val Trp Lys Asp Leu Leu Glu Asp Ser Val Thr Pro Ile Asp
 65 70 75

<210> 33
 <211> 61
 <212> PRT
 <213> Hepatitis C virus

<400> 33
 Thr Gln Thr Val Asp Phe Ser Leu Asp Pro Thr Phe Thr Ile Glu Thr
 1 5 10 15
 Thr Thr Leu Pro Gln Asp Ala Val Ser Arg Thr Gln Arg Arg Gly Arg
 20 25 30
 Thr Gly Arg Gly Lys Pro Gly Ile Tyr Arg Phe Val Ala Pro Gly Glu
 35 40 45
 Arg Pro Ser Gly Met Phe Asp Ser Ser Val Leu Cys Glu
 50 55 60

<210> 34
 <211> 77
 <212> PRT
 <213> Hepatitis C virus

<400> 34
 Val Leu Asp Ser His Tyr Gln Asp Val Leu Lys Glu Val Lys Ala Ala
 1 5 10 15
 Ala Ser Lys Val Lys Ala Asn Leu Leu Ser Val Glu Glu Ala Cys Ser
 20 25 30
 Leu Thr Pro Pro His Ser Ala Lys Ser Lys Phe Gly Tyr Gly Ala Lys
 35 40 45
 Asp Val Arg Cys His Ala Arg Lys Ala Val Ala His Ile Asn Ser Val
 50 55 60
 Trp Lys Asp Leu Leu Glu Asp Ser Val Thr Pro Ile Asp
 65 70 75

<210> 35
 <211> 26
 <212> PRT
 <213> Hepatitis C virus

<400> 35

Tyr Pro Pro Arg Pro Cys Gly Ile Val Pro Ala Lys Ser Val Cys Gly
 1 5 10 15

Pro Val Tyr Cys Phe Thr Pro Ser Pro Val
 20 25

<210> 36

<211> 37

<212> PRT

<213> Hepatitis C virus

<400> 36

Pro Ile Ser Tyr Ala Asn Gly Ser Gly Leu Asp Glu Arg Pro Tyr Cys
 1 5 10 15

Trp His Tyr Pro Pro Arg Pro Cys Gly Ile Val Pro Ala Lys Ser Val
 20 25 30

Cys Gly Pro Val Tyr
 35

<210> 37

<211> 35

<212> PRT

<213> Hepatitis C virus

<400> 37

Thr Val Thr Gln Leu Leu Arg Arg Leu His Gln Trp Ile Ser Ser Glu
 1 5 10 15

Cys Thr Thr Pro Cys Ser Gly Ser Trp Leu Arg Asp Ile Trp Asp Trp
 20 25 30

Ile Cys Glu
 35

<210> 38

<211> 25

<212> PRT

<213> Hepatitis C virus

<400> 38

Leu Leu Arg Arg Leu His Gln Trp Ile Ser Ser Glu Cys Thr Thr Pro
 1 5 10 15

Cys Ser Gly Ser Trp Leu Arg Asp Ile
 20 25

<210> 39
<211> 152
<212> DNA
<213> Hepatitis C virus

<400> 39
cttgttgccg cgcaggggcc ctagattggg tgtgcgcgcg acgaggaaga cttccgagcg 60
gtcgcaacct cgaggtagac gtcagcctat cccaaggca cgtcggcccg agggcaggac 120
ctgggctcag ccgggtacc cttggccct ct 152

<210> 40
<211> 106
<212> DNA
<213> Hepatitis C virus

<400> 40
tggcaggggg aagccaggca tctatagatt tgtggcacgc ggggagcgcc cctccggcat 60
gttcgactcg tccgtcctct gtgagtgcga tgacgcgggc tgtgct 106

<210> 41
<211> 234
<212> DNA
<213> Hepatitis C virus

<400> 41
taacaccaac cgtcgccac aggacgtcaa gttcccgggt ggcggtcaga tcgttggtgg 60
agtttacttg ttgccgcgca ggggccctag attgggtgtg cgcgcgacga ggaagacttc 120
cgagcggtcg caacctcgag gtagacgtca gcctatcccc aaggcacgtc ggcccgaggg 180
caggacctgg gtcagcccg ggtacccttg gccctctat ggcaatgagg gttg 234

<210> 42
<211> 114
<212> DNA
<213> Hepatitis C virus

<400> 42
tcccagcccc gtggtggtgg gaacgaccga caggtcgggc gcgcctacct acagctgggg 60
tgcaaatgat acggatgtct tcgtccttaa caacaccagg ccaccgctgg gcaa 114

<210> 43
<211> 453
<212> DNA
<213> Hepatitis C virus

<400> 43
ctccaggact caacgccggg gcaggactgg cagggggaag ccaggcatct atagatttgt 60
ggcaccgggg gagcgccct cggcatgtt cgactcgcc gtccctctgt agtgctatga 120

```

cgcgggctgt gcttggtatg agctcacgcc cgccgagact acagttaggc tacgagcgta 180
catgaacacc ccgggggttc ccgtgtgcca ggaccatott gaattttggg agggcgcttt 240
tacgggcctc actcatatag atgcccactt tttatcccag acaaagcaga gtggggagaa 300
ctttccttac ctggtagcgt accaagccac cgtgtgcgct agggctcaag cccctcccc 360
atcgtgggac cagatgtgga agtgtttgat ccgccttaaa cccaccctcc atgggccaac 420
accctgcta tacagactgg gcgctgttca gaa 453

```

```

<210> 44
<211> 85
<212> DNA
<213> Hepatitis C virus

```

```

<400> 44
tcccagcccc gtggtggtgg gaacgaccga caggtcgggc gcgcctacct acagctgggg 60
tgcaaatgat acgatgtct tcgtc 85

```

```

<210> 45
<211> 80
<212> DNA
<213> Hepatitis C virus

```

```

<400> 45
ccctccaaga ccttgtggca ttgtgccgc aaagagcgtg tgtggcccg tatattgctt 60
cactcccagc ccgtggtgg 80

```

```

<210> 46
<211> 165
<212> DNA
<213> Hepatitis C virus

```

```

<400> 46
ctgcgtggtc atagtgggca ggatcgtctt gtccgggaag ccggcaatta tacctgacag 60
ggaggttctc taccaggagt tcgatgagat ggaagagtgc tctcagcact taccgtacat 120
cgagcaaggg atgatgctcg ctgagcagtt caagcagaag gccct 165

```

```

<210> 47
<211> 123
<212> DNA
<213> Hepatitis C virus

```

```

<400> 47
cggcgacttc gactctgtga tagactgcaa cacgtgtgtc actcagacag tcgatttcag 60
ccttgaccct acctttacca ttgagacaac cacgtcccc caggatgctg tctccaggac 120
tca 123

```

```

<210> 48
<211> 87
<212> DNA
<213> Hepatitis C virus

```

<400> 48

ggagcgcccc tccggcatgt tcgactcgtc cgtcctctgt gagtgcctatg acgcgggctg 60
tgcttggtat gagctcacgc ccgccga 87

<210> 49

<211> 84

<212> DNA

<213> Hepatitis C virus

<400> 49

cagggggaag ccaggcatct atagatttgt ggcaccgggg gagcgccct cggcatgtt 60
cgactcgtcc gtcctctgtg agtg 84

<210> 50

<211> 128

<212> DNA

<213> Hepatitis C virus

<400> 50

tctggaagac agtgtaacac caatagacac taccatcatg gccagaacg aggttttctg 60
cgttcagcct gagaaggggg gtcgtaagcc agctcgtctc atcgtgttcc ccgacctggg 120
cgtgcgcg 128

<210> 51

<211> 85

<212> DNA

<213> Hepatitis C virus

<400> 51

tcccaccggc agcggtaaga gcaccaaggt cccggctgcg tacgcagccc agggctacaa 60
ggtgttggtg ctcaaccct ctgtt 85

<210> 52

<211> 102

<212> DNA

<213> Hepatitis C virus

<400> 52

cgaacgcccc tactgctggc actaccctcc aagaccttgt ggcattgtgc ccgcaaagag 60
cgtgtgtggc ccggtatatt gcttcaactcc cagccccgtg gt 102

<210> 53

<211> 95

<212> DNA

<213> Hepatitis C virus

<400> 53

tcccagcccc gtggtggtgg gaacgaccga caggtcgggc gcgcctacct acagctgggg 60

tgcaaatgat acggatgtct tcgtccttaa caaca

95

<210> 54

<211> 234

<212> DNA

<213> Hepatitis C virus

<400> 54

ggcgggagct cttgtagcat tcaagatcat gagcggtag gteccctcca cggaggacct 60
 ggtcaatctg ctgcccacca tectctcgcc tggagccctt gtagtcgggtg tggctcgcgc 120
 agcaatactg cgccggcacg ttggcccggg cgaggggggca gtgcaatgga tgaaccggct 180
 aatagccttc gcctcccggg ggaaccatgt ttccccacg cactacgtgc cgga 234

<210> 55

<211> 442

<212> DNA

<213> Hepatitis C virus

<400> 55

tgagggtccag atcgtgtcaa ctgctaccca aaccttcctg gcaacgtgca tcaatgggggt 60
 atgctggact gtctaccacg gggccggaac gaggaccatc gcatcaccca agggtcctgt 120
 catccagatg tataccaatg tggaccaaga ccttgtgggc tggcccgctc ctcaaggctc 180
 ccgctcattg acaccctgta cctgcggctc ctcgacactt tacctggtca cgaggcacgc 240
 cgatgtcatt cccgtgcgce ggcgagggtg tagcaggggt agcctgcttt cgccccggcc 300
 catttcttac ttgaaaggct cctcgggggg tccgctgttg tgccccggcg gacacgccgt 360
 gggcctattc agggccgchg tgtgcacccg tggagtggct aaagcgggtg actttatccc 420
 tgtgggagaac ctagggacaa cc 442

<210> 56

<211> 111

<212> DNA

<213> Hepatitis C virus

<400> 56

tgtaaccag ctctgagggc gactgcatca gtggataagc tcggagtgtg ccaactccatg 60
 ctccgggtcc tggctaaggg acatctggga ctggatatgc gaggtgctga g 111

<210> 57

<211> 87

<212> DNA

<213> Hepatitis C virus

<400> 57

cgtgtgtggc ccggtatatt gcttcactcc cagccccgtg gtgggtgggaa cgaccgacag 60
 gtcgggcgcg cctacctaca gctgggg 87

<210> 58

<211> 137

<212> DNA

<213> Hepatitis C virus

<400> 58

```
cccgcccttg cgagcttgga gacaccgggc cggagcgtc cgcgctaggc ttctgtccag 60
aggaggcagg gctgccatat gtggcaagta cctcttcaac tgggcagtaa gaacaaagct 120
caaactcact ccaatag                                     137
```

<210> 59

<211> 259

<212> DNA

<213> Hepatitis C virus

<400> 59

```
tactgccttt gtgggtgctg gcctagctgg cgcgcccatc ggcagcgctg gactggggaa 60
ggtcctcgtg gacattcttg cagggtatgg cgcgggcgtg gcgggagctc ttgtagcatt 120
caagatcatg agcggtgagg tcccctccac ggaggacctg gtcaatctgc tgcccgccat 180
cctctcgctt ggagcccttg tagtcggtgt ggtctgcgca gcaatactgc gccggcacgt 240
tggcccgggc gagggggca                                     259
```

<210> 60

<211> 130

<212> DNA

<213> Hepatitis C virus

<400> 60

```
tggcattgtg cccgcaaaga gcgtgtgtgg cccggtatat tgcttcactc ccagccccgt 60
gggtgtggga acgaccgaca ggtcgggcgc gcctacctac agctgggggtg caaatgatac 120
ggatgtcttc                                     130
```

<210> 61

<211> 191

<212> DNA

<213> Hepatitis C virus

<400> 61

```
ggtcctcgtg gacattcttg cagggtatgg cgcgggcgtg gcgggagctc ttgtagcatt 60
caagatcatg agcggtgagg tcccctccac ggaggacctg gtcaatctgc tgcccgccat 120
cctctcgctt ggagcccttg tagtcggtgt ggtctgcgca gcaatactgc gccggcacgt 180
tggcccgggc g                                     191
```

<210> 62

<211> 89

<212> DNA

<213> Hepatitis C virus

<400> 62

```
cgaacgcccc tactgctggc actaccctcc aagaccttgt ggcattgtgc ccgcaaagag 60
cgtgtgtggc ccggtatatt gcttcactc                                     89
```

<210> 63
 <211> 230
 <212> DNA
 <213> Hepatitis C virus

<400> 63
 caggcgccac tggacgacgc aagactgcaa ttgttctatc tatcccggcc atataacggg 60
 tcatcgcatg gcatgggata tgatgatgaa ctgggtcccct acggcagcgt tgggtggtagc 120
 tcagctgctc cggatcccac aagccatcat ggacatgac gctggtgctc actggggagt 180
 cctggcgggc atagcgtatt tctccatggg ggggaactgg gcgaaggtec 230

<210> 64
 <211> 113
 <212> DNA
 <213> Hepatitis C virus

<400> 64
 tgccatactc agcagcctca ctgtaaccca gctcctgagg cgactgcac agtgggataag 60
 ctcgagagtgt accactccat gctccgggtc ctggctaagg gacatctggg act 113

<210> 65
 <211> 142
 <212> DNA
 <213> Hepatitis C virus

<400> 65
 tgtctccagg actcaacgcc ggggcaggac tggcaggggg aagccaggca tctatagatt 60
 tgtggcaccg ggggagcgcc cctccggcat gttcgactcg tccgtcctct gtgagtgcta 120
 tgacgcgggc tgtgcttggg at 142

<210> 66
 <211> 162
 <212> DNA
 <213> Hepatitis C virus

<400> 66
 ccttcctgcg ccgaactata agttcgcgct gtggaggggtg tctgcagagg aatacgtgga 60
 gataaggcgg gtgggggact tccactacgt atcgggtatg actactgaca atcttaaattg 120
 cccgtgccag atcccatcgc ccgaattttt cacagaattg ga 162

<210> 67
 <211> 337
 <212> DNA
 <213> Hepatitis C virus

<400> 67
 cggagagatc cccttttacg gcaaggctat cccctcagag gtgatcaagg ggggaagaca 60
 tctcatcttc tgccactcaa agaagaagtg cgacgagctc gccgcgaagc tgggtcgatt 120
 gggcatcaat gccgtggcct actaccggcg tcttgacgtg tctgtcatcc cgaccagcgg 180
 cgatgttgtc gtcgtgtcga ccgatgctct catgactggc tttaccggcg acttcgactc 240

tgtgatagac tgcaacacgt gtgtcactca gacagtcgat ttcagccttg accctacctt 300
taccattgag acaaccacgc tccccagga tgctgtc 337

<210> 68

<211> 163

<212> DNA

<213> Hepatitis C virus

<400> 68

ggtctgcgca gcaataactgc gccggcacgt tggcccgggc gagggggcag tgcaatggat 60
gaaccggcta atagccttcg cctcccgggg gaaccatgtt tccccacgc actacgtgcc 120
ggagagcgat gcagccgccc gcgtcactgc catactcagc agc 163

<210> 69

<211> 309

<212> DNA

<213> Hepatitis C virus

<400> 69

ggccatcaag tccctcactg agaggcttta tgttgggggc cctcttacca attcaagggg 60
ggaaaactgc ggctaccgca ggtgccgcgc gagcggcgta ctgacaacta gctgtggtaa 120
caccctcact tgctacatca aggcccgggc agcctgtcga gccgcagggc tccaggactg 180
caccatgctc gtgtgtggcg acgacttagt cgttatctgt gaaagtgcgg ggggtccagga 240
ggacgcggcg agcctgagag ccttcacgga ggctatgacc aggtactcgg ccccccccg 300
ggaccccc 309

<210> 70

<211> 240

<212> DNA

<213> Hepatitis C virus

<400> 70

actgcaagtt ctggacagcc attaccagga cgtgctcaag gaggtcaaag cagcggcgtc 60
aaaagtgaag gctaacttgc tatccgtaga ggaagcttgc agcctgacgc cccacattc 120
agccaaatcc aagtttggct atggggcaaa agacgtccgt tgccatgcca gaaaggccgt 180
agccacatc aactccgtgt ggaaagacct tctggaagac agtctaacac caatagacac 240

<210> 71

<211> 184

<212> DNA

<213> Hepatitis C virus

<400> 71

cactcagaca gtcgatttca gccttgaccc tacctttacc attgagacaa ccacgctccc 60
ccaggatgct gtctccagga ctcaacgcgc gggcaggact ggcaggggga agccaggcat 120
ctatagattt gtggcaccgg gggagcgccc ctccggcatg ttcgactcgt ccgtcctctg 180
tgag 184

<210> 72

<211> 234
<212> DNA
<213> Hepatitis C virus

<400> 72
agttctggac agccattacc aggacgtgct caaggaggtc aaagcagcgg cgtcaaaagt 60
gaaggctaac ttgctatccg tagaggaagc ttgcagcctg acgccccac attcagccaa 120
atccaagttt ggctatgggg caaaagacgt ccgttgccat gccagaaagg ccgtagccca 180
catcaactcc gtgtggaaag accttctgga agacagtgtg acaccaatag acac 234

<210> 73
<211> 80
<212> DNA
<213> Hepatitis C virus

<400> 73
ctaccctcca agaccttgtg gcattgtgcc cgcaaagagc gtgtgtggcc cggatatattg 60
cttcactccc agccccgtgg 80

<210> 74
<211> 112
<212> DNA
<213> Hepatitis C virus

<400> 74
tcctatcagt tatgccaacg gaagcggcct cgacgaacgc ccctactgct ggcaactacc 60
tccaagacct tgtggcattg tgcccgcaaa gagcgtgtgt ggcccggat at 112

<210> 75
<211> 107
<212> DNA
<213> Hepatitis C virus

<400> 75
cactgtaacc cagctcctga ggcgactgca tcagtggata agctcggagt gtaccactcc 60
atgctccggg tcctggctaa gggacatctg ggactggata tgcgagg 107

<210> 76
<211> 78
<212> DNA
<213> Hepatitis C virus

<400> 76
gtccctgagg cgactgcac agtggataag ctccggagtgt accactccat gctccggttc 60
ctggctaagg gacatctg 78

<210> 77
<211> 103
<212> PRT

<213> Hepatitis C virus

<400> 77

Ala Cys Glu Cys Pro Gly Arg Ser Arg Arg Pro Cys Thr Met Ser Thr
 1 5 10 15

Asn Pro Lys Pro Gln Arg Lys Thr Lys Arg Asn Thr Asn Arg Arg Pro
 20 25 30

Gln Asp Val Lys Phe Pro Gly Gly Gly Gln Ile Val Gly Gly Val Tyr
 35 40 45

Leu Leu Pro Arg Arg Gly Pro Arg Leu Gly Val Arg Ala Thr Arg Lys
 50 55 60

Thr Ser Glu Arg Ser Gln Pro Arg Gly Arg Arg Gln Pro Ile Pro Lys
 65 70 75 80

Ala Arg Arg Pro Glu Gly Arg Thr Trp Ala Gln Pro Gly Tyr Pro Trp
 85 90 95

Pro Leu Tyr Gly Asn Glu Gly
 100

<210> 78

<211> 113

<212> PRT

<213> Hepatitis C virus

<400> 78

Met Ser Thr Asn Pro Lys Pro Gln Arg Lys Thr Lys Arg Asn Thr Asn
 1 5 10 15

Arg Arg Pro Gln Asp Val Lys Phe Pro Gly Gly Gly Gln Ile Val Gly
 20 25 30

Gly Val Tyr Leu Leu Pro Arg Arg Gly Pro Arg Leu Gly Val Arg Ala
 35 40 45

Thr Arg Lys Thr Ser Glu Arg Ser Gln Pro Arg Gly Arg Arg Gln Pro
 50 55 60

Ile Pro Lys Ala Arg Arg Pro Glu Gly Arg Thr Trp Ala Gln Pro Gly
 65 70 75 80

Tyr Pro Trp Pro Leu Tyr Gly Asn Glu Gly Cys Gly Trp Ala Gly Trp
 85 90 95

Leu Leu Ser Pro Arg Gly Ser Arg Pro Ser Trp Gly Pro Thr Asp Pro
 100 105 110

Arg

<210> 79
 <211> 114
 <212> PRT
 <213> Hepatitis C virus

<400> 79
 Ala Ile Leu His Thr Pro Gly Cys Val Pro Cys Val Arg Glu Gly Asn
 1 5 10 15
 Ala Ser Arg Cys Trp Val Ala Val Thr Pro Thr Val Ala Thr Arg Asp
 20 25 30
 Gly Lys Leu Pro Thr Thr Gln Leu Arg Arg His Ile Asp Leu Leu Val
 35 40 45
 Gly Ser Ala Thr Leu Cys Ser Ala Leu Tyr Val Gly Asp Leu Cys Gly
 50 55 60
 Ser Val Phe Leu Val Gly Gln Leu Phe Thr Phe Ser Pro Arg Arg His
 65 70 75 80
 Trp Thr Thr Gln Asp Cys Asn Cys Ser Ile Tyr Pro Gly His Ile Thr
 85 90 95
 Gly His Arg Met Ala Trp Asp Met Met Met Asn Trp Ser Pro Thr Ala
 100 105 110
 Ala Leu

<210> 80
 <211> 91
 <212> PRT
 <213> Hepatitis C virus

<400> 80
 Gly Val Asp Ala Glu Thr His Val Thr Gly Gly Asn Ala Gly Arg Thr
 1 5 10 15
 Thr Ala Gly Leu Val Gly Leu Leu Thr Pro Gly Ala Lys Gln Asn Ile
 20 25 30
 Gln Leu Ile Asn Thr Asn Gly Ser Trp His Ile Asn Ser Thr Ala Leu
 35 40 45
 Asn Cys Asn Glu Ser Leu Asn Thr Gly Trp Leu Ala Gly Leu Phe Tyr
 50 55 60

25

Gln His Lys Phe Asn Ser Ser Gly Cys Pro Glu Arg Leu Ala Ser Cys
 65 70 75 80

Arg Arg Leu Thr Asp Phe Ala Gln Gly Trp Gly
 85 90

<210> 81

<211> 176

<212> PRT

<213> Hepatitis C virus

<400> 81

Trp Gly Pro Ile Ser Tyr Ala Asn Gly Ser Gly Leu Asp Glu Arg Pro
 1 5 10 15

Tyr Cys Trp His Tyr Pro Pro Arg Pro Cys Gly Ile Val Pro Ala Lys
 20 25 30

Ser Val Cys Gly Pro Val Tyr Cys Phe Thr Pro Ser Pro Val Val Val
 35 40 45

Gly Thr Thr Asp Arg Ser Gly Ala Pro Thr Tyr Ser Trp Gly Ala Asn
 50 55 60

Asp Thr Asp Val Phe Val Leu Asn Asn Thr Arg Pro Pro Leu Gly Asn
 65 70 75 80

Trp Phe Gly Cys Thr Trp Met Asn Ser Thr Gly Phe Thr Lys Val Cys
 85 90 95

Gly Ala Pro Pro Cys Val Ile Gly Gly Val Gly Asn Asn Thr Leu Leu
 100 105 110

Cys Pro Thr Asp Cys Phe Arg Lys His Pro Glu Ala Thr Tyr Ser Arg
 115 120 125

Cys Gly Ser Gly Pro Trp Ile Thr Pro Arg Cys Met Val Asp Tyr Pro
 130 135 140

Tyr Arg Leu Trp His Tyr Pro Cys Thr Ile Asn Tyr Thr Ile Phe Lys
 145 150 155 160

Val Arg Met Tyr Val Gly Gly Val Glu His Arg Leu Glu Ala Ala Cys
 165 170 175

<210> 82

<211> 96

<212> PRT

<213> Hepatitis C virus

<400> 82

Trp	His	Tyr	Pro	Pro	Arg	Pro	Cys	Gly	Ile	Val	Pro	Ala	Lys	Ser	Val
1				5					10					15	
Cys	Gly	Pro	Val	Tyr	Cys	Phe	Thr	Pro	Ser	Pro	Val	Val	Val	Gly	Thr
			20					25					30		
Thr	Asp	Arg	Ser	Gly	Ala	Pro	Thr	Tyr	Ser	Trp	Gly	Ala	Asn	Asp	Thr
		35					40					45			
Asp	Val	Phe	Val	Leu	Asn	Asn	Thr	Arg	Pro	Pro	Leu	Gly	Asn	Trp	Phe
	50					55					60				
Gly	Cys	Thr	Trp	Met	Asn	Ser	Thr	Gly	Phe	Thr	Lys	Val	Cys	Gly	Ala
65					70					75					80
Pro	Pro	Cys	Val	Ile	Gly	Gly	Val	Gly	Asn	Asn	Thr	Leu	Leu	Cys	Pro
				85					90					95	

<210> 83

<211> 278

<212> PRT

<213> Hepatitis C virus

<400> 83

Ala	Ala	Cys	Gly	Asp	Ile	Ile	Asn	Gly	Leu	Pro	Val	Ser	Ala	Arg	Arg
1				5					10					15	
Gly	Gln	Glu	Ile	Leu	Leu	Gly	Pro	Ala	Asp	Gly	Met	Val	Ser	Lys	Gly
			20					25					30		
Trp	Arg	Leu	Leu	Ala	Pro	Ile	Thr	Ala	Tyr	Ala	Gln	Gln	Thr	Arg	Gly
		35					40					45			
Leu	Leu	Gly	Cys	Ile	Ile	Thr	Ser	Leu	Thr	Gly	Arg	Asp	Lys	Asn	Gln
	50					55					60				
Val	Glu	Gly	Glu	Val	Gln	Ile	Val	Ser	Thr	Ala	Thr	Gln	Thr	Phe	Leu
65					70					75					80
Ala	Thr	Cys	Ile	Asn	Gly	Val	Cys	Trp	Thr	Val	Tyr	His	Gly	Ala	Gly
				85					90					95	
Thr	Arg	Thr	Ile	Ala	Ser	Pro	Lys	Gly	Pro	Val	Ile	Gln	Met	Tyr	Thr
			100					105					110		

```

Asn Val Asp Gln Asp Leu Val Gly Trp Pro Ala Pro Gln Gly Ser Arg
    115                      120                      125

Ser Leu Thr Pro Cys Thr Cys Gly Ser Ser Asp Leu Tyr Leu Val Thr
    130                      135                      140

Arg His Ala Asp Val Ile Pro Val Arg Arg Arg Gly Asp Ser Arg Gly
    145                      150                      155                      160

Ser Leu Leu Ser Pro Arg Pro Ile Ser Tyr Leu Lys Gly Ser Ser Gly
                      165                      170                      175

Gly Pro Leu Leu Cys Pro Ala Gly His Ala Val Gly Leu Phe Arg Ala
    180                      185                      190

Ala Val Cys Thr Arg Gly Val Ala Lys Ala Val Asp Phe Ile Pro Val
    195                      200                      205

Glu Asn Leu Gly Thr Thr Met Arg Ser Pro Val Phe Thr Asp Asn Ser
    210                      215                      220

Ser Pro Pro Ala Val Pro Gln Ser Phe Gln Val Ala His Leu His Ala
    225                      230                      235                      240

Pro Thr Gly Ser Gly Lys Ser Thr Lys Val Pro Ala Ala Tyr Ala Ala
    245                      250                      255

Gln Gly Tyr Lys Val Leu Val Leu Asn Pro Ser Val Ala Ala Thr Leu
    260                      265                      270

Gly Phe Gly Ala Tyr Met
    275

```

```

<210> 84
<211> 158
<212> PRT
<213> Hepatitis C virus

```

```

<400> 84
Arg Arg Arg Gly Asp Ser Arg Gly Ser Leu Leu Ser Pro Arg Pro Ile
    1                      5                      10                      15

Ser Tyr Leu Lys Gly Ser Ser Gly Gly Pro Leu Leu Cys Pro Ala Gly
    20                      25                      30

His Ala Val Gly Leu Phe Arg Ala Ala Val Cys Thr Arg Gly Val Ala
    35                      40                      45

Lys Ala Val Asp Phe Ile Pro Val Glu Asn Leu Gly Thr Thr Met Arg
    50                      55                      60

Ser Pro Val Phe Thr Asp Asn Ser Ser Pro Pro Ala Val Pro Gln Ser

```

65		70		75		80									
Phe	Gln	Val	Ala	His	Leu	His	Ala	Pro	Thr	Gly	Ser	Gly	Lys	Ser	Thr
				85					90					95	
Lys	Val	Pro	Ala	Ala	Tyr	Ala	Ala	Gln	Gly	Tyr	Lys	Val	Leu	Val	Leu
			100					105					110		
Asn	Pro	Ser	Val	Ala	Ala	Thr	Leu	Gly	Phe	Gly	Ala	Tyr	Met	Ser	Lys
		115					120					125			
Ala	His	Gly	Val	Asp	Pro	Asn	Ile	Arg	Thr	Gly	Val	Arg	Thr	Ile	Thr
	130						135				140				
Thr	Gly	Ser	Pro	Ile	Thr	Tyr	Ser	Thr	Tyr	Gly	Lys	Phe	Leu		
145					150					155					

<210> 85
 <211> 263
 <212> PRT
 <213> Hepatitis C virus

<400> 85
 Asp Ser Arg Gly Ser Leu Leu Ser Pro Arg Pro Ile Ser Tyr Leu Lys
 1 5 10 15

Gly Ser Ser Gly Gly Pro Leu Leu Cys Pro Ala Gly His Ala Val Gly
 20 25 30

Leu Phe Arg Ala Ala Val Cys Thr Arg Gly Val Ala Lys Ala Val Asp
 35 40 45

Phe Ile Pro Val Glu Asn Leu Gly Thr Thr Met Arg Ser Pro Val Phe
 50 55 60

Thr Asp Asn Ser Ser Pro Pro Ala Val Pro Gln Ser Phe Gln Val Ala
 65 70 75 80

His Leu His Ala Pro Thr Gly Ser Gly Lys Ser Thr Lys Val Pro Ala
 85 90 95

Ala Tyr Ala Ala Gln Gly Tyr Lys Val Leu Val Leu Asn Pro Ser Val
 100 105 110

Ala Ala Thr Leu Gly Phe Gly Ala Tyr Met Ser Lys Ala His Gly Val
 115 120 125

Asp Pro Asn Ile Arg Thr Gly Val Arg Thr Ile Thr Thr Gly Ser Pro
 130 135 140

Ile Thr Tyr Ser Thr Tyr Gly Lys Phe Leu Ala Asp Gly Gly Cys Ser
 145 150 155 160

Gly Gly Ala Tyr Asp Ile Ile Ile Cys Asp Glu Cys His Ser Thr Asp
 165 170 175
 Ala Thr Ser Ile Leu Gly Ile Gly Thr Val Leu Asp Gln Ala Glu Thr
 180 185 190
 Ala Gly Ala Arg Leu Val Val Leu Ala Thr Ala Thr Pro Pro Gly Ser
 195 200 205
 Val Thr Val Ser His Pro Asn Ile Glu Glu Val Ala Leu Ser Thr Thr
 210 215 220
 Gly Glu Ile Pro Phe Tyr Gly Lys Ala Ile Pro Leu Glu Val Ile Lys
 225 230 235 240
 Gly Gly Arg His Leu Ile Phe Cys His Ser Lys Lys Lys Cys Asp Glu
 245 250 255
 Leu Ala Ala Lys Leu Val Ala
 260

<210> 86
 <211> 194
 <212> PRT
 <213> Hepatitis C virus

<400> 86
 Asp Asn Ser Ser Pro Pro Ala Val Pro Gln Ser Phe Gln Val Ala His
 1 5 10 15
 Leu His Ala Pro Thr Gly Ser Gly Lys Ser Thr Lys Val Pro Ala Ala
 20 25 30
 Tyr Ala Ala Gln Gly Tyr Lys Val Leu Val Leu Asn Pro Ser Val Ala
 35 40 45
 Ala Thr Leu Gly Phe Gly Ala Tyr Met Ser Lys Ala His Gly Val Asp
 50 55 60
 Pro Asn Ile Arg Thr Gly Val Arg Thr Ile Thr Thr Gly Ser Pro Ile
 65 70 75 80
 Thr Tyr Ser Thr Tyr Gly Lys Phe Leu Ala Asp Gly Gly Cys Ser Gly
 85 90 95
 Gly Ala Tyr Asp Ile Ile Ile Cys Asp Glu Cys His Ser Thr Asp Ala
 100 105 110
 Thr Ser Ile Leu Gly Ile Gly Thr Val Leu Asp Gln Ala Glu Thr Ala
 115 120 125

Gly Ala Arg Leu Val Val Leu Ala Thr Ala Thr Pro Pro Gly Ser Val
 130 135 140

Thr Val Ser His Pro Asn Ile Glu Glu Val Ala Leu Ser Thr Thr Gly
 145 150 155 160

Glu Ile Pro Phe Tyr Gly Lys Ala Ile Pro Leu Glu Val Ile Lys Gly
 165 170 175

Gly Arg His Leu Ile Phe Cys His Ser Lys Lys Lys Cys Asp Glu Leu
 180 185 190

Ala Ala

<210> 87
 <211> 205
 <212> PRT
 <213> Hepatitis C virus

<400> 87
 Leu Ala Asp Gly Gly Cys Ser Gly Gly Ala Tyr Asp Ile Ile Ile Cys
 1 5 10 15

Asp Glu Cys His Ser Thr Asp Ala Thr Ser Ile Leu Gly Ile Gly Thr
 20 25 30

Val Leu Asp Gln Ala Glu Thr Ala Gly Ala Arg Leu Val Val Leu Ala
 35 40 45

Thr Ala Thr Pro Pro Gly Ser Val Thr Val Ser His Pro Asn Ile Glu
 50 55 60

Glu Val Ala Leu Ser Thr Thr Gly Glu Ile Pro Phe Tyr Gly Lys Ala
 65 70 75 80

Ile Pro Leu Glu Val Ile Lys Gly Gly Arg His Leu Ile Phe Cys His
 85 90 95

Ser Lys Lys Lys Cys Asp Glu Leu Ala Ala Lys Leu Val Ala Leu Gly
 100 105 110

Ile Asn Ala Val Ala Tyr Tyr Arg Gly Leu Asp Val Ser Val Ile Pro
 115 120 125

Thr Ser Gly Asp Val Val Val Val Ser Thr Asp Ala Leu Met Thr Gly
 130 135 140

Phe Thr Gly Asp Phe Asp Ser Val Ile Asp Cys Asn Thr Cys Val Thr
 145 150 155 160

Gln Thr Val Asp Phe Ser Leu Asp Pro Thr Phe Thr Ile Glu Thr Thr


```

                165                170                175
Thr Leu Pro Gln Asp Ala Val Ser Arg Thr Gln Arg Arg Gly Arg Thr
                180                185                190

Gly Arg Gly Lys Pro Gly Ile Tyr Arg Phe Val Ala Pro
                195                200                205

<210> 88
<211> 186
<212> PRT
<213> Hepatitis C virus

<400> 88
Ser Thr Asp Ala Thr Ser Ile Leu Gly Ile Gly Thr Val Leu Asp Gln
  1                5                10                15

Ala Glu Thr Ala Gly Ala Arg Leu Val Val Leu Ala Thr Ala Thr Pro
                20                25                30

Pro Gly Ser Val Thr Val Ser His Pro Asn Ile Glu Glu Val Ala Leu
                35                40                45

Ser Thr Thr Gly Glu Ile Pro Phe Tyr Gly Lys Ala Ile Pro Leu Glu
  50                55                60

Val Ile Lys Gly Gly Arg His Leu Ile Phe Cys His Ser Lys Lys Lys
  65                70                75                80

Cys Asp Glu Leu Ala Ala Lys Leu Val Ala Leu Gly Ile Asn Ala Val
                85                90                95

Ala Tyr Tyr Arg Gly Leu Asp Val Ser Val Ile Pro Thr Ser Gly Asp
                100                105                110

Val Val Val Val Ser Thr Asp Ala Leu Met Thr Gly Phe Thr Gly Asp
                115                120                125

Phe Asp Ser Val Ile Asp Cys Asn Thr Cys Val Thr Gln Thr Val Asp
                130                135                140

Phe Ser Leu Asp Pro Thr Phe Thr Ile Glu Thr Thr Thr Leu Pro Gln
                145                150                155                160

Asp Ala Val Ser Arg Thr Gln Arg Arg Gly Arg Thr Gly Arg Gly Lys
                165                170                175

Pro Gly Ile Tyr Arg Phe Val Ala Pro Gly
                180                185

```

<210> 89
 <211> 158
 <212> PRT
 <213> Hepatitis C virus

<400> 89
 Val Ile Asp Cys Asn Thr Cys Val Thr Gln Thr Val Asp Phe Ser Leu
 1 5 10 15
 Asp Pro Thr Phe Thr Ile Glu Thr Thr Thr Leu Pro Gln Asp Ala Val
 20 25 30
 Ser Arg Thr Gln Arg Arg Gly Arg Thr Gly Arg Gly Lys Pro Gly Ile
 35 40 45
 Tyr Arg Phe Val Ala Pro Gly Glu Arg Pro Ser Gly Met Phe Asp Ser
 50 55 60
 Ser Val Leu Cys Glu Cys Tyr Asp Ala Gly Cys Ala Trp Tyr Glu Leu
 65 70 75 80
 Thr Pro Ala Glu Thr Thr Val Arg Leu Arg Ala Tyr Met Asn Thr Pro
 85 90 95
 Gly Leu Pro Val Cys Gln Asp His Leu Glu Phe Trp Glu Gly Val Phe
 100 105 110
 Thr Gly Leu Thr His Ile Asp Ala His Phe Leu Ser Gln Thr Lys Gln
 115 120 125
 Ser Gly Glu Asn Phe Pro Tyr Leu Val Ala Tyr Gln Ala Thr Val Cys
 130 135 140
 Ala Arg Ala Gln Ala Pro Pro Pro Ser Trp Asp Gln Met Trp
 145 150 155

<210> 90
 <211> 129
 <212> PRT
 <213> Hepatitis C virus

<400> 90
 Arg Phe Val Ala Pro Gly Glu Arg Pro Ser Gly Met Phe Asp Ser Ser
 1 5 10 15
 Val Leu Cys Glu Cys Tyr Asp Ala Gly Cys Ala Trp Tyr Glu Leu Thr
 20 25 30
 Pro Ala Glu Thr Thr Val Arg Leu Arg Ala Tyr Met Asn Thr Pro Gly
 35 40 45
 Leu Pro Val Cys Gln Asp His Leu Glu Phe Trp Glu Gly Val Phe Thr

33

50 55 60
 Gly Leu Thr His Ile Asp Ala His Phe Leu Ser Gln Thr Lys Gln Ser
 65 70 75 80
 Gly Glu Asn Phe Pro Tyr Leu Val Ala Tyr Gln Ala Thr Val Cys Ala
 85 90 95
 Arg Ala Gln Ala Pro Pro Pro Ser Trp Asp Gln Met Trp Lys Cys Leu
 100 105 110
 Ile Arg Leu Lys Pro Thr Leu His Gly Pro Thr Pro Leu Leu Tyr Arg
 115 120 125
 Leu

<210> 91
 <211> 51
 <212> PRT
 <213> Hepatitis C virus

<400> 91
 Thr Ser Thr Trp Val Leu Val Gly Gly Val Leu Ala Ala Leu Ala Ala
 1 5 10 15
 Tyr Cys Leu Ser Thr Gly Cys Val Val Ile Val Gly Arg Ile Val Leu
 20 25 30
 Ser Gly Lys Pro Ala Ile Ile Pro Asp Arg Glu Val Leu Tyr Gln Glu
 35 40 45
 Phe Asp Glu
 50

<210> 92
 <211> 18
 <212> PRT
 <213> Hepatitis C virus

<400> 92
 Ala Ala Leu Ala Ala Tyr Cys Leu Ser Thr Gly Cys Val Val Ile Val
 1 5 10 15
 Gly Arg

<210> 93

<211> 208
 <212> PRT
 <213> Hepatitis C virus

<400> 93

Phe	Thr	Ala	Ala	Val	Thr	Ser	Pro	Leu	Thr	Thr	Gly	Gln	Thr	Leu	Leu
1				5					10					15	
Phe	Asn	Ile	Leu	Gly	Gly	Trp	Val	Ala	Ala	Gln	Leu	Ala	Ala	Pro	Gly
			20					25						30	
Ala	Ala	Thr	Ala	Phe	Val	Gly	Ala	Gly	Leu	Ala	Gly	Ala	Ala	Ile	Gly
		35					40					45			
Ser	Val	Gly	Leu	Gly	Lys	Val	Leu	Val	Asp	Ile	Leu	Ala	Gly	Tyr	Gly
	50					55					60				
Ala	Gly	Val	Ala	Gly	Ala	Leu	Val	Ala	Phe	Lys	Ile	Met	Ser	Gly	Glu
65					70					75					80
Val	Pro	Ser	Thr	Glu	Asp	Leu	Val	Asn	Leu	Leu	Pro	Ala	Ile	Leu	Ser
				85					90					95	
Pro	Gly	Ala	Leu	Val	Val	Gly	Val	Val	Cys	Ala	Ala	Ile	Leu	Arg	Arg
		100						105					110		
His	Val	Gly	Pro	Gly	Glu	Gly	Ala	Val	Gln	Trp	Met	Asn	Arg	Leu	Ile
		115					120					125			
Ala	Phe	Ala	Ser	Arg	Gly	Asn	His	Val	Ser	Pro	Thr	His	Tyr	Val	Pro
	130					135					140				
Glu	Ser	Asp	Ala	Ala	Ala	Arg	Val	Thr	Ala	Ile	Leu	Ser	Ser	Leu	Thr
145					150					155					160
Val	Thr	Gln	Leu	Leu	Arg	Arg	Leu	His	Gln	Trp	Ile	Ser	Ser	Glu	Cys
			165						170					175	
Thr	Thr	Pro	Cys	Ser	Gly	Ser	Trp	Leu	Arg	Asp	Ile	Trp	Asp	Trp	Ile
			180					185					190		
Cys	Glu	Val	Leu	Ser	Asp	Phe	Lys	Thr	Trp	Leu	Lys	Ala	Lys	Leu	Met
		195					200					205			

<210> 94
 <211> 207
 <212> PRT
 <213> Hepatitis C virus

35

<400> 94

Thr Ala Phe Val Gly Ala Gly Leu Ala Gly Ala Ala Ile Gly Ser Val
 1 5 10 15

Gly Leu Gly Lys Val Leu Val Asp Ile Leu Ala Gly Tyr Gly Ala Gly
 20 25 30

Val Ala Gly Ala Leu Val Ala Phe Lys Ile Met Ser Gly Glu Val Pro
 35 40 45

Ser Thr Glu Asp Leu Val Asn Leu Leu Pro Ala Ile Leu Ser Pro Gly
 50 55 60

Ala Leu Val Val Gly Val Val Cys Ala Ala Ile Leu Arg Arg His Val
 65 70 75 80

Gly Pro Gly Glu Gly Ala Val Gln Trp Met Asn Arg Leu Ile Ala Phe
 85 90 95

Ala Ser Arg Gly Asn His Val Ser Pro Thr His Tyr Val Pro Glu Ser
 100 105 110

Asp Ala Ala Ala Arg Val Thr Ala Ile Leu Ser Ser Leu Thr Val Thr
 115 120 125

Gln Leu Leu Arg Arg Leu His Gln Trp Ile Ser Ser Glu Cys Thr Thr
 130 135 140

Pro Cys Ser Gly Ser Trp Leu Arg Asp Ile Trp Asp Trp Ile Cys Glu
 145 150 155 160

Val Leu Ser Asp Phe Lys Thr Trp Leu Lys Ala Lys Leu Met Pro Gln
 165 170 175

Leu Pro Gly Ile Pro Phe Val Ser Cys Gln Arg Gly Tyr Arg Gly Val
 180 185 190

Trp Arg Gly Asp Gly Ile Met His Thr Arg Cys His Cys Gly Ala
 195 200 205

<210> 95

<211> 225

<212> PRT

<213> Hepatitis C virus

<400> 95

Leu Val Asp Ile Leu Ala Gly Tyr Gly Ala Gly Val Ala Gly Ala Leu
 1 5 10 15

Val Ala Phe Lys Ile Met Ser Gly Glu Val Pro Ser Thr Glu Asp Leu
 20 25 30

[illegible]

```
<210> 96
<211> 145
<212> PRT
<213> Hepatitis C virus
```

```
<400> 96
Ala Gly Tyr Gly Ala Gly Val Ala Gly Ala Leu Val Ala Phe Lys Ile
  1             5             10             15
```

Met Ser Gly Glu Val Pro Ser Thr Glu Asp Leu Val Asn Leu Leu Pro
20 25 30

Ala Ile Leu Ser Pro Gly Ala Leu Val Val Gly Val Val Cys Ala Ala

35		40		45
Ile Leu Arg Arg His Val Gly Pro Gly Glu Gly Ala Val Gln Trp Met				
50		55		60
Asn Arg Leu Ile Ala Phe Ala Ser Arg Gly Asn His Val Ser Pro Thr				
65		70		75 80
His Tyr Val Pro Glu Ser Asp Ala Ala Ala Arg Val Thr Ala Ile Leu				
	85		90	95
Ser Ser Leu Thr Val Thr Gln Leu Leu Arg Arg Leu His Gln Trp Ile				
	100		105	110
Ser Ser Glu Cys Thr Thr Pro Cys Ser Gly Ser Trp Leu Arg Asp Ile				
	115		120	125
Trp Asp Trp Ile Cys Glu Val Leu Ser Asp Phe Lys Thr Trp Leu Lys				
	130		135	140
Ala				
145				

<210> 97
 <211> 54
 <212> PRT
 <213> Hepatitis C virus

<400> 97

Ala Leu Val Val Gly Val Val Cys Ala Ala Ile Leu Arg Arg His Val				
1		5		10 15
Gly Pro Gly Glu Gly Ala Val Gln Trp Met Asn Arg Leu Ile Ala Phe				
	20		25	30
Ala Ser Arg Gly Asn His Val Ser Pro Thr His Tyr Val Pro Glu Ser				
	35		40	45
Asp Ala Ala Ala Arg Val				
50				

<210> 98
 <211> 165
 <212> PRT
 <213> Hepatitis C virus

<400> 98

Ala Ser Arg Gly Asn His Val Ser Pro Thr His Tyr Val Pro Glu Ser				
1		5		10 15

Asp Ala Ala Ala Arg Val Thr Ala Ile Leu Ser Ser Leu Thr Val Thr
 20 25 30
 Gln Leu Leu Arg Arg Leu His Gln Trp Ile Ser Ser Glu Cys Thr Thr
 35 40 45
 Pro Cys Ser Gly Ser Trp Leu Arg Asp Ile Trp Asp Trp Ile Cys Glu
 50 55 60
 Val Leu Ser Asp Phe Lys Thr Trp Leu Lys Ala Lys Leu Met Pro Gln
 65 70 75 80
 Leu Pro Gly Ile Pro Phe Val Ser Cys Gln Arg Gly Tyr Arg Gly Val
 85 90 95
 Trp Arg Gly Asp Gly Ile Met His Thr Arg Cys His Cys Gly Ala Glu
 100 105 110
 Ile Thr Gly His Val Lys Asn Gly Thr Met Arg Ile Val Gly Pro Arg
 115 120 125
 Thr Cys Arg Asn Met Trp Ser Gly Thr Phe Pro Ile Asn Ala Tyr Thr
 130 135 140
 Thr Gly Pro Cys Thr Pro Leu Pro Ala Pro Asn Tyr Lys Phe Ala Leu
 145 150 155 160
 Trp Arg Val Ser Ala
 165

<210> 99
 <211> 308
 <212> PRT
 <213> Hepatitis C virus

<400> 99
 Tyr Val Pro Glu Ser Asp Ala Ala Ala Arg Val Thr Ala Ile Leu Ser
 1 5 10 15
 Ser Leu Thr Val Thr Gln Leu Leu Arg Arg Leu His Gln Trp Ile Ser
 20 25 30
 Ser Glu Cys Thr Thr Pro Cys Ser Gly Ser Trp Leu Arg Asp Ile Trp
 35 40 45
 Asp Trp Ile Cys Glu Val Leu Ser Asp Phe Lys Thr Trp Leu Lys Ala
 50 55 60
 Lys Leu Met Pro Gln Leu Pro Gly Ile Pro Phe Val Ser Cys Gln Arg
 65 70 75 80
 Gly Tyr Arg Gly Val Trp Arg Gly Asp Gly Ile Met His Thr Arg Cys

85								90				95			
His	Cys	Gly	Ala	Glu	Ile	Thr	Gly	His	Val	Lys	Asn	Gly	Thr	Met	Arg
100				105				110							
Ile	Val	Gly	Pro	Arg	Thr	Cys	Arg	Asn	Met	Trp	Ser	Gly	Thr	Phe	Pro
115				120				125							
Ile	Asn	Ala	Tyr	Thr	Thr	Gly	Pro	Cys	Thr	Pro	Leu	Pro	Ala	Pro	Asn
130				135				140							
Tyr	Lys	Phe	Ala	Leu	Trp	Arg	Val	Ser	Ala	Glu	Glu	Tyr	Val	Glu	Ile
145				150				155				160			
Arg	Arg	Val	Gly	Asp	Phe	His	Tyr	Val	Ser	Gly	Met	Thr	Thr	Asp	Asn
165				170				175							
Leu	Lys	Cys	Pro	Cys	Gln	Ile	Pro	Ser	Pro	Glu	Phe	Phe	Thr	Glu	Leu
180				185				190							
Asp	Gly	Val	Arg	Leu	His	Arg	Phe	Ala	Pro	Pro	Cys	Lys	Pro	Leu	Leu
195				200				205							
Arg	Glu	Glu	Val	Ser	Phe	Arg	Val	Gly	Leu	His	Glu	Tyr	Pro	Val	Gly
210				215				220							
Ser	Gln	Leu	Pro	Cys	Glu	Pro	Glu	Pro	Asp	Val	Ala	Val	Leu	Thr	Ser
225				230				235				240			
Met	Leu	Thr	Asp	Pro	Ser	His	Ile	Thr	Ala	Glu	Ala	Ala	Gly	Arg	Arg
245				250				255							
Leu	Ala	Arg	Gly	Ser	Pro	Pro	Ser	Met	Ala	Ser	Ser	Ser	Ala	Ser	Gln
260				265				270							
Leu	Ser	Ala	Pro	Ser	Leu	Lys	Ala	Thr	Cys	Thr	Ala	Asn	His	Asp	Ser
275				280				285							
Pro	Asp	Ala	Glu	Leu	Ile	Glu	Ala	Asn	Leu	Leu	Trp	Arg	Gln	Glu	Met
290				295				300							
Gly	Gly	Asn	Ile												
305															

```
<210> 100
<211> 283
<212> PRT
<213> Hepatitis C virus
```

<400> 100
Leu Ser Ser Leu Thr Val Thr Gln Leu Leu Arg Arg Leu His Gln Trp
1 5 10 15

```

Ile Ser Ser Glu Cys Thr Thr Pro Cys Ser Gly Ser Trp Leu Arg Asp
      20                      25                      30

Ile Trp Asp Trp Ile Cys Glu Val Leu Ser Asp Phe Lys Thr Trp Leu
      35                      40                      45

Lys Ala Lys Leu Met Pro Gln Leu Pro Gly Ile Pro Phe Val Ser Cys
      50                      55                      60

Gln Arg Gly Tyr Arg Gly Val Trp Arg Gly Asp Gly Ile Met His Thr
      65                      70                      75                      80

Arg Cys His Cys Gly Ala Glu Ile Thr Gly His Val Lys Asn Gly Thr
      85                      90                      95

Met Arg Ile Val Gly Pro Arg Thr Cys Arg Asn Met Trp Ser Gly Thr
      100                     105                     110

Phe Pro Ile Asn Ala Tyr Thr Thr Gly Pro Cys Thr Pro Leu Pro Ala
      115                     120                     125

Pro Asn Tyr Lys Phe Ala Leu Trp Arg Val Ser Ala Glu Glu Tyr Val
      130                     135                     140

Glu Ile Arg Arg Val Gly Asp Phe His Tyr Val Ser Gly Met Thr Thr
      145                     150                     155                     160

Asp Asn Leu Lys Cys Pro Cys Gln Ile Pro Ser Pro Glu Phe Phe Thr
      165                     170                     175

Glu Leu Asp Gly Val Arg Leu His Arg Phe Ala Pro Pro Cys Lys Pro
      180                     185                     190

Leu Leu Arg Glu Glu Val Ser Phe Arg Val Gly Leu His Glu Tyr Pro
      195                     200                     205

Val Gly Ser Gln Leu Pro Cys Glu Pro Glu Pro Asp Val Ala Val Leu
      210                     215                     220

Thr Ser Met Leu Thr Asp Pro Ser His Ile Thr Ala Glu Ala Ala Gly
      225                     230                     235                     240

Arg Arg Leu Ala Arg Gly Ser Pro Pro Ser Met Ala Ser Ser Ser Ala
      245                     250                     255

Ser Gln Leu Ser Ala Pro Ser Leu Lys Ala Thr Cys Thr Ala Asn His
      260                     265                     270

Asp Ser Pro Asp Ala Glu Leu Ile Glu Ala Asn
      275                     280

```

<210> 101
 <211> 249
 <212> PRT
 <213> Hepatitis C virus

<400> 101
 Ser Leu Thr Val Thr Gln Leu Leu Arg Arg Leu His Gln Trp Ile Ser
 1 5 10 15
 Ser Glu Cys Thr Thr Pro Cys Ser Gly Ser Trp Leu Arg Asp Ile Trp
 20 25 30
 Asp Trp Ile Cys Glu Val Leu Ser Asp Phe Lys Thr Trp Leu Lys Ala
 35 40 45
 Lys Leu Met Pro Gln Leu Pro Gly Ile Pro Phe Val Ser Cys Gln Arg
 50 55 60
 Gly Tyr Arg Gly Val Trp Arg Gly Asp Gly Ile Met His Thr Arg Cys
 65 70 75 80
 His Cys Gly Ala Glu Ile Thr Gly His Val Lys Asn Gly Thr Met Arg
 85 90 95
 Ile Val Gly Pro Arg Thr Cys Arg Asn Met Trp Ser Gly Thr Phe Pro
 100 105 110
 Ile Asn Ala Tyr Thr Thr Gly Pro Cys Thr Pro Leu Pro Ala Pro Asn
 115 120 125
 Tyr Lys Phe Ala Leu Trp Arg Val Ser Ala Glu Glu Tyr Val Glu Ile
 130 135 140
 Arg Arg Val Gly Asp Phe His Tyr Val Ser Gly Met Thr Thr Asp Asn
 145 150 155 160
 Leu Lys Cys Pro Cys Gln Ile Pro Ser Pro Glu Phe Phe Thr Glu Leu
 165 170 175
 Asp Gly Val Arg Leu His Arg Phe Ala Pro Pro Cys Lys Pro Leu Leu
 180 185 190
 Arg Glu Glu Val Ser Phe Arg Val Gly Leu His Glu Tyr Pro Val Gly
 195 200 205
 Ser Gln Leu Pro Cys Glu Pro Glu Pro Asp Val Ala Val Leu Thr Ser
 210 215 220
 Met Leu Thr Asp Pro Ser His Ile Thr Ala Glu Ala Ala Gly Arg Arg
 225 230 235 240
 Leu Ala Arg Gly Ser Pro Pro Ser Met
 245

<210> 102
 <211> 85
 <212> PRT
 <213> Hepatitis C virus

<400> 102
 Thr Trp Leu Lys Ala Lys Leu Met Pro Gln Leu Pro Gly Ile Pro Phe
 1 5 10 15
 Val Ser Cys Gln Arg Gly Tyr Arg Gly Val Trp Arg Gly Asp Gly Ile
 20 25 30
 Met His Thr Arg Cys His Cys Gly Ala Glu Ile Thr Gly His Val Lys
 35 40 45
 Asn Gly Thr Met Arg Ile Val Gly Pro Arg Thr Cys Arg Asn Met Trp
 50 55 60
 Ser Gly Thr Phe Pro Ile Asn Ala Tyr Thr Thr Gly Pro Cys Thr Pro
 65 70 75 80
 Leu Pro Ala Pro Asn
 85

<210> 103
 <211> 94
 <212> PRT
 <213> Hepatitis C virus

<400> 103
 Glu Ile Thr Gly His Val Lys Asn Gly Thr Met Arg Ile Val Gly Pro
 1 5 10 15
 Arg Thr Cys Arg Asn Met Trp Ser Gly Thr Phe Pro Ile Asn Ala Tyr
 20 25 30
 Thr Thr Gly Pro Cys Thr Pro Leu Pro Ala Pro Asn Tyr Lys Phe Ala
 35 40 45
 Leu Trp Arg Val Ser Ala Glu Glu Tyr Val Glu Ile Arg Arg Val Gly
 50 55 60
 Asp Phe His Tyr Val Ser Gly Met Thr Thr Asp Asn Leu Lys Cys Pro
 65 70 75 80
 Cys Gln Ile Pro Ser Pro Glu Phe Phe Thr Glu Leu Asp Gly
 85 90

<210> 104
 <211> 75
 <212> PRT
 <213> Hepatitis C virus

<400> 104
 Ile Glu Ala Asn Leu Leu Trp Arg Gln Glu Met Gly Gly Asn Ile Thr
 1 5 10 15
 Arg Val Glu Ser Glu Asn Lys Val Val Ile Leu Asp Ser Phe Asp Pro
 20 25 30
 Leu Val Ala Glu Glu Asp Glu Arg Glu Val Ser Val Pro Ala Glu Ile
 35 40 45
 Leu Arg Lys Ser Arg Arg Phe Ala Arg Ala Leu Pro Val Trp Ala Arg
 50 55 60
 Pro Asp Tyr Asn Pro Pro Leu Val Glu Thr Trp
 65 70 75

<210> 105
 <211> 90
 <212> PRT
 <213> Hepatitis C virus

<400> 105
 His Gly Cys Pro Leu Pro Pro Pro Arg Ser Pro Pro Val Pro Pro Pro
 1 5 10 15
 Arg Lys Lys Arg Thr Val Val Leu Thr Glu Ser Thr Leu Ser Thr Ala
 20 25 30
 Leu Ala Glu Leu Ala Thr Lys Ser Phe Gly Ser Ser Ser Thr Ser Gly
 35 40 45
 Ile Thr Gly Asp Asn Thr Thr Thr Ser Ser Glu Pro Ala Pro Ser Gly
 50 55 60
 Cys Pro Pro Asp Ser Asp Val Glu Ser Tyr Ser Ser Met Pro Pro Leu
 65 70 75 80
 Glu Gly Glu Pro Gly Asp Pro Asp Leu Ser
 85 90

<210> 106
 <211> 137
 <212> PRT
 <213> Hepatitis C virus

<400> 106

Ser Trp Thr Gly Ala Leu Val Thr Pro Cys Ala Ala Glu Glu Gln Lys
 1 5 10 15

Leu Pro Ile Asn Ala Leu Ser Asn Ser Leu Leu Arg His His Asn Leu
 20 25 30

Val Tyr Ser Thr Thr Ser Arg Ser Ala Cys Gln Arg Gln Lys Lys Val
 35 40 45

Thr Phe Asp Arg Leu Gln Val Leu Asp Ser His Tyr Gln Asp Val Leu
 50 55 60

Lys Glu Val Lys Ala Ala Ala Ser Lys Val Lys Ala Asn Leu Leu Ser
 65 70 75 80

Val Glu Glu Ala Cys Ser Leu Thr Pro Pro His Ser Ala Lys Ser Lys
 85 90 95

Phe Gly Tyr Gly Ala Lys Asp Val Arg Cys His Ala Arg Lys Ala Val
 100 105 110

Ala His Ile Asn Ser Val Trp Lys Asp Leu Leu Glu Asp Ser Val Thr
 115 120 125

Pro Ile Asp Thr Thr Ile Met Ala Lys
 130 135

<210> 107

<211> 300

<212> PRT

<213> Hepatitis C virus

<400> 107

Ala Asn Leu Leu Ser Val Glu Glu Ala Cys Ser Leu Thr Pro Pro His
 1 5 10 15

Ser Ala Lys Ser Lys Phe Gly Tyr Gly Ala Lys Asp Val Arg Cys His
 20 25 30

Ala Arg Lys Ala Val Ala His Ile Asn Ser Val Trp Lys Asp Leu Leu
 35 40 45

Glu Asp Ser Val Thr Pro Ile Asp Thr Thr Ile Met Ala Lys Asn Glu
 50 55 60

Val Phe Cys Val Gln Pro Glu Lys Gly Gly Arg Lys Pro Ala Arg Leu
 65 70 75 80

Ile Val Phe Pro Asp Leu Gly Val Arg Val Cys Glu Lys Met Ala Leu
 85 90 95

```

Tyr Asp Val Val Ser Lys Leu Pro Leu Ala Val Met Gly Ser Ser Tyr
      100                      105                      110

Gly Phe Gln Tyr Ser Pro Gly Gln Arg Val Glu Phe Leu Val Gln Ala
      115                      120                      125

Trp Lys Ser Lys Lys Thr Pro Met Gly Phe Ser Tyr Asp Thr Arg Cys
      130                      135                      140

Phe Asp Ser Thr Val Thr Glu Ser Asp Ile Arg Thr Glu Glu Ala Ile
      145                      150                      155                      160

Tyr Gln Cys Cys Asp Leu Asp Pro Gln Ala Arg Val Ala Ile Lys Ser
      165                      170                      175

Leu Thr Glu Arg Leu Tyr Val Gly Gly Pro Leu Thr Asn Ser Arg Gly
      180                      185                      190

Glu Asn Cys Gly Tyr Arg Arg Cys Arg Ala Ser Gly Val Leu Thr Thr
      195                      200                      205

Ser Cys Gly Asn Thr Leu Thr Cys Tyr Ile Lys Ala Arg Ala Ala Cys
      210                      215                      220

Arg Ala Ala Gly Leu Gln Asp Cys Thr Met Leu Val Cys Gly Asp Asp
      225                      230                      235                      240

Leu Val Val Ile Cys Glu Ser Ala Gly Val Gln Glu Asp Ala Ala Ser
      245                      250                      255

Leu Arg Ala Phe Thr Glu Ala Met Thr Arg Tyr Ser Ala Pro Pro Gly
      260                      265                      270

Asp Pro Pro Gln Pro Glu Tyr Asp Leu Glu Leu Ile Thr Ser Cys Ser
      275                      280                      285

Ser Asn Val Ser Val Ala His Asp Gly Ala Gly Lys
      290                      295                      300

```

<210> 108

<211> 199

<212> PRT

<213> Hepatitis C virus

<400> 108

```

Glu Glu Ala Cys Ser Leu Thr Pro Pro His Ser Ala Lys Ser Lys Phe
  1              5              10              15

```

```

Gly Tyr Gly Ala Lys Asp Val Arg Cys His Ala Arg Lys Ala Val Ala
      20              25              30

```

```

His Ile Asn Ser Val Trp Lys Asp Leu Leu Glu Asp Ser Val Thr Pro

```

35	40	45
Ile Asp Thr Thr Ile Met Ala Lys Asn Glu Val Phe Cys Val Gln Pro		
50	55	60
Glu Lys Gly Gly Arg Lys Pro Ala Arg Leu Ile Val Phe Pro Asp Leu		
65	70	75
Gly Val Arg Val Cys Glu Lys Met Ala Leu Tyr Asp Val Val Ser Lys		
	85	90
		95
Leu Pro Leu Ala Val Met Gly Ser Ser Tyr Gly Phe Gln Tyr Ser Pro		
	100	105
		110
Gly Gln Arg Val Glu Phe Leu Val Gln Ala Trp Lys Ser Lys Lys Thr		
	115	120
		125
Pro Met Gly Phe Ser Tyr Asp Thr Arg Cys Phe Asp Ser Thr Val Thr		
	130	135
		140
Glu Ser Asp Ile Arg Thr Glu Glu Ala Ile Tyr Gln Cys Cys Asp Leu		
145	150	155
		160
Asp Pro Gln Ala Arg Val Ala Ile Lys Ser Leu Thr Glu Arg Leu Tyr		
	165	170
		175
Val Gly Gly Pro Leu Thr Asn Ser Arg Gly Glu Asn Cys Gly Tyr Arg		
	180	185
		190
Arg Cys Arg Ala Ser Gly Val		
195		

<210> 109

<211> 260

<212> PRT

<213> Hepatitis C virus

<400> 109

Leu Leu Glu Asp Ser Val Thr Pro Ile Asp Thr Thr Ile Met Ala Lys
1 5 10 15

Asn Glu Val Phe Cys Val Gln Pro Glu Lys Gly Gly Arg Lys Pro Ala
20 25 30

Arg Leu Ile Val Phe Pro Asp Leu Gly Val Arg Val Cys Glu Lys Met
35 40 45

Ala Leu Tyr Asp Val Val Ser Lys Leu Pro Leu Ala Val Met Gly Ser
50 55 60

Ser Tyr Gly Phe Gln Tyr Ser Pro Gly Gln Arg Val Glu Phe Leu Val
65 70 75 80

Gln Ala Trp Lys Ser Lys Lys Thr Pro Met Gly Phe Ser Tyr Asp Thr
 85 90 95
 Arg Cys Phe Asp Ser Thr Val Thr Glu Ser Asp Ile Arg Thr Glu Glu
 100 105 110
 Ala Ile Tyr Gln Cys Cys Asp Leu Asp Pro Gln Ala Arg Val Ala Ile
 115 120 125
 Lys Ser Leu Thr Glu Arg Leu Tyr Val Gly Gly Pro Leu Thr Asn Ser
 130 135 140
 Arg Gly Glu Asn Cys Gly Tyr Arg Arg Cys Arg Ala Ser Gly Val Leu
 145 150 155 160
 Thr Thr Ser Cys Gly Asn Thr Leu Thr Cys Tyr Ile Lys Ala Arg Ala
 165 170 175
 Ala Cys Arg Ala Ala Gly Leu Gln Asp Cys Thr Met Leu Val Cys Gly
 180 185 190
 Asp Asp Leu Val Val Ile Cys Glu Ser Ala Gly Val Gln Glu Asp Ala
 195 200 205
 Ala Ser Leu Arg Ala Phe Thr Glu Ala Met Thr Arg Tyr Ser Ala Pro
 210 215 220
 Pro Gly Asp Pro Pro Gln Pro Glu Tyr Asp Leu Glu Leu Ile Thr Ser
 225 230 235 240
 Cys Ser Ser Asn Val Ser Val Ala His Asp Gly Ala Gly Lys Arg Val
 245 250 255
 Tyr Tyr Leu Thr
 260

<210> 110
 <211> 127
 <212> PRT
 <213> Hepatitis C virus

<400> 110
 Val Ile Cys Glu Ser Ala Gly Val Gln Glu Asp Ala Ala Ser Leu Arg
 1 5 10 15
 Ala Phe Thr Glu Ala Met Thr Arg Tyr Ser Ala Pro Pro Gly Asp Pro
 20 25 30
 Pro Gln Pro Glu Tyr Asp Leu Glu Leu Ile Thr Ser Cys Ser Ser Asn
 35 40 45

Val Ser Val Ala His Asp Gly Ala Gly Lys Arg Val Tyr Tyr Leu Thr
 50 55 60

Arg Asp Pro Thr Thr Pro Leu Ala Arg Ala Ala Trp Glu Thr Ala Arg
 65 70 75 80

His Thr Pro Val Asn Ser Trp Leu Gly Asn Ile Ile Met Phe Ala Pro
 85 90 95

Thr Leu Trp Ala Arg Met Ile Leu Met Thr His Phe Phe Ser Val Leu
 100 105 110

Ile Ala Arg Asp Gln Leu Glu Gln Ala Leu Asn Cys Glu Ile Tyr
 115 120 125

<210> 111
 <211> 89
 <212> PRT
 <213> Hepatitis C virus

<400> 111
 Val Ser Val Ala His Asp Gly Ala Gly Lys Arg Val Tyr Tyr Leu Thr
 1 5 10 15

Arg Asp Pro Thr Thr Pro Leu Ala Arg Ala Ala Trp Glu Thr Ala Arg
 20 25 30

His Thr Pro Val Asn Ser Trp Leu Gly Asn Ile Ile Met Phe Ala Pro
 35 40 45

Thr Leu Trp Ala Arg Met Ile Leu Met Thr His Phe Phe Ser Val Leu
 50 55 60

Ile Ala Arg Asp Gln Leu Glu Gln Ala Leu Asn Cys Glu Ile Tyr Gly
 65 70 75 80

Ala Cys Tyr Ser Ile Glu Pro Leu Asp
 85

<210> 112
 <211> 73
 <212> PRT
 <213> Hepatitis C virus

<400> 112
 Leu His Gly Leu Ser Ala Phe Ser Leu His Ser Tyr Ser Pro Gly Glu
 1 5 10 15

Ile Asn Arg Val Ala Ala Cys Leu Arg Lys Leu Gly Val Pro Pro Leu
 20 25 30

Arg Ala Trp Arg His Arg Ala Arg Ser Val Arg Ala Arg Leu Leu Ser
35 40 45

Arg Gly Gly Arg Ala Ala Ile Cys Gly Lys Tyr Leu Phe Asn Trp Ala
50 55 60

Val Arg Thr Lys Leu Lys Leu Thr Pro
65 70

<210> 113

<211> 63

<212> PRT

<213> Hepatitis C virus

<400> 113

Ser Pro Gly Glu Ile Asn Arg Val Ala Ala Cys Leu Arg Lys Leu Gly
1 5 10 15

Val Pro Pro Leu Arg Ala Trp Arg His Arg Ala Arg Ser Val Arg Ala
20 25 30

Arg Leu Leu Ser Arg Gly Gly Arg Ala Ala Ile Cys Gly Lys Tyr Leu
35 40 45

Phe Asn Trp Ala Val Arg Thr Lys Leu Lys Leu Thr Pro Ile Ala
50 55 60

<210> 114

<211> 310

<212> DNA

<213> Hepatitis C virus

<400> 114

tgcttgcgag tgccccggga ggtctcgtag accgtgcacc atgagcacga atcctaaacc 60
tcaaagaaaa accaaacgta acaccaaccg tcgcccacag gacgtcaagt tcccgggtgg 120
cggtcagatc gttggtggag tttacttggt gccgcgcagg ggccctagat tgggtgtgcg 180
cgcgacgagg aagacttccg agcgggtcgca acctcgaggt agacgtcagc ctatcccca 240
ggcacgtcgg cccgagggca ggacctgggc tcagcccggg tacccttggc ccctctatgg 300
caatgagggt 310

<210> 115

<211> 339

<212> DNA

<213> Hepatitis C virus

<400> 115

atgagcacga atcctaaacc tcaaagaaaa accaaacgta acaccaaccg tcgcccacag 60
gacgtcaagt tcccgggtgg cggtcagatc gttggtggag tttacttggt gccgcgcagg 120

```

ggccctagat tgggtgtgcg cgcgacgagg aagacttccg agcggtcgca acctcgaggt 180
agacgtcagc ctatcccca ggacgtcgcg cccgagggca ggacctgggc tcagcccggg 240
tacccttggc ccctctatgg caatgagggt tgcgggtggg cgggatggct cctgtctccc 300
cgtggctctc ggccctagctg gggcccccaca gacccccgg 339

```

```

<210> 116
<211> 345
<212> DNA
<213> Hepatitis C virus

```

```

<400> 116
tgccatcctg cacactccgg ggtgtgtccc ttgcgttcgc gagggtaacg cctcgaggtg 60
ttgggtggcg gtgaccccca cgggtggccac cagggacggc aaactcccca caacgcagct 120
tcgacgtcat atcgatctgc ttgtcgggag cgccaccctc tgctcggccc totacgtggg 180
ggacctgtgc ggggtctgtct ttcttgttgg tcaactgttt accttctctc ccaggcgcca 240
ctggacgacg caagactgca attgttctat ctatcccggc catataacgg gtcacgcgat 300
ggcatgggat atgatgatga actgggtccc taaggcagcg ttggt 345

```

```

<210> 117
<211> 276
<212> DNA
<213> Hepatitis C virus

```

```

<400> 117
cggcgtcgac ggggaaaccc acgtcacggg gggaaatgcc ggccgcacca cgggtggggt 60
tgttgggtctc cttacaccag ggcgcaagca gaacatocaa ctgatcaaca ccaacggcag 120
ttggcacatc aatagcacgg ccttgaattg caatgaaagc cttaacaccg gctggttagc 180
agggctcttc tatcaacaca aattcaactc ttcaggctgt cctgagaggt tggccagctg 240
ccgacgcctt accgattttg cccagggtcg ggggtcc 276

```

```

<210> 118
<211> 531
<212> DNA
<213> Hepatitis C virus

```

```

<400> 118
ctggggtcct atcagttatg ccaacggaag cggcctcgac gaacgcccct actgctggca 60
ctacctcca agaccttgtg gcatttgtgc cgcaaagagc gtgtgtggcc cggatatattg 120
cttcactccc agccccgtgg tgggtgggaac gaccgacagg tcgggcgcgc ctacctacag 180
ctgggggtgca aatgatacgg atgtcttcgt ccttaacaac accaggccac cgctgggcaa 240
ttggttcggg tgtacctgga tgaactcaac tggattcacc aaagtgtgcg gagegcccc 300
ttgtgtcatc ggaggggtgg gcaacaacac cttgctctgc ccactgatt gcttccgcaa 360
acatccggaa gccacatact ctcggtgcgg ctccgggtccc tggattacac ccagggtgat 420
ggtcgactac ccgtataggc tttggcacta tccttgtacc atcaattaca ccatattcaa 480
agtcaggatg tacgtgggag gggtcgagca caggctggaa gcggcctgca a 531

```

```

<210> 119
<211> 289
<212> DNA
<213> Hepatitis C virus

```

<400> 119

```

ctggcactac cctccaagac cttgtggcat tgtgcccgca aagagcgtgt gtggcccgggt 60
atattgcttc actcccagcc ccgtggtgggt gggaaacgacc gacaggtcgg gcgcgcctac 120
ctacagctgg ggtgcaaattg atacggatgt cttcgtcctt aacaacacca ggccaccgct 180
gggcaatttg ttcggttgta cctggatgaa ctcaactgga ttcaccaaag tgtgcggagc 240
gcccccttgt gtcacgagag ggggtgggcaa caacaccttg ctctgcccc 289

```

<210> 120

<211> 836

<212> DNA

<213> Hepatitis C virus

<400> 120

```

gccgcgtgcg gtgacatcat caacggcttg cccgtctctg cccgtagggg ccaggagata 60
ctgcttgggc cagccgacgg aatggctctc aaggggtgga ggttgctggc gcccatcacg 120
gcgtacgccc agcagacgag aggcctccta ggggtgataa tcaccagcct gactggccgg 180
gacaaaaacc aagtggaggg tgaggtccag atcgtgtcaa ctgctacca aaccttcctg 240
gcaacgtgca tcaatggggg atgctggact gtctaccacg gggccggaac gaggaccatc 300
gcatcaccca agggtcctgt catccagatg tataccaatg tggaccaaga ccttggtggc 360
tggcccgcct ctcaagggtc ccgctcattg acaccctgta cctgcggctc ctccgacctt 420
tacctggtca cgaggcacgc cgatgtcatt cccgtgcgcc gccgaggtga tagcaggggt 480
agcctgcttt cgcgccggcc catttcctac ttgaaaggct cctcgggggg tccgctgttg 540
tgcccccgcg gacacgcgct gggcctattc agggccgcgg tgtgcacccg tggagtggct 600
aaagcgggtg actttatccc tgtggagaac ctagggacaa ccatgagatc cccggtgttc 660
acggacaact cctctccacc agcagtgcgc cagagcttcc aggtggccca cctgcatgct 720
cccaccggca gcggttaagag caccaaggct ccggctgcgt acgcagccca gggctacaag 780
gtgttggtgc tcaacccctc tgttgctgca acgctgggct ttggtgctta catgtc 836

```

<210> 121

<211> 475

<212> DNA

<213> Hepatitis C virus

<400> 121

```

gcgcggcgca ggtgatagca ggggtagcct gctttcgccc cggcccatth cctacttgaa 60
aggctcctcg ggggggtccg tgttggtgcc cgcgggacac gccgtgggoc tattcagggc 120
cgcggtgtgc acccgtggag tggctaaagc ggtggacttt atccctgtgg agaaccctagg 180
gacaaccatg agatccccgg tgttcacgga caactcctct ccaccagcag tgccccagag 240
cttccaggtg gccacacctg atgctcccac cggcagcggg aagagcacca aggtcccggc 300
tgcgtagcga gccacgggct acaagggtgt ggtgctcaac ccctctgttg ctgcaacgct 360
gggctttggt gcttacatgt ccaaggccca tggggttgat cctaatatca ggaccggggg 420
gagaacaatt accactggca gcccacatcac gtactccacc tacggcaagt tcctt 475

```

<210> 122

<211> 790

<212> DNA

<213> Hepatitis C virus

<400> 122

```

tgatagcagg ggtagcctgc tttcgccccg gccatttcc tacttgaaag gtcctcggg 60

```

```

gggtccgctg ttgtgccccg cgggacacgc cgtgggccta ttcagggccg cgggtgtgcac 120
ccgtggagtg gctaaagcgg tggactttat ccctgtggag aacctaggga caaccatgag 180
atccccggtg ttcacggaca actcctctcc accagcagtg cccagagct tccaggtggc 240
ccacctgcat gctcccaccg gcagcggtaa gagcaccaag gtcccggctg cgtacgcagc 300
ccagggctac aagggtgttg tgc tcaaccc ctctgttgct gcaacgctgg gctttgggtgc 360
ttacatgtcc aaggcccatg gggttgatcc taatatcagg accgggggtga gaacaattac 420
cactggcagc cccatcacgt actccaccta cggcaagttc cttgccgacg gcgggtgctc 480
aggaggtgct tatgacataa taatttgtga cgagtgccac tccacggatg ccacatccat 540
cttgggcatc ggcaactgtc ttgaccaagc agagactgcg ggggcgagac tggttgtgct 600
cgccactgct acccctccgg gctccgtcac tgtgtcccat cctaaccatcg aggaggttgc 660
tctgtccacc accggagaga tcccccttta cggcaaggct atccccctcg aggtgatcaa 720
ggggggaaga catctcatct tctgccactc aaagaagaag tgcgacgagc tcgccgcgaa 780
gctggctcgca                                     790

```

<210> 123

<211> 583

<212> DNA

<213> Hepatitis C virus

<400> 123

```

ggacaactcc tctccaccag cagtgcocca gagcttccag gtggcccacc tgcattgctcc 60
caccggcagc ggtaagagca ccaaggtccc ggctgcgtac gcagcccagg gctacaaggt 120
gttgggtgct aacccctctg ttgctgcaac gctgggcttt ggtgcttaca tgtccaaggc 180
ccatgggggtt gatcctaata tcaggaccgg ggtgagaaca attaccactg gcagcccat 240
cacgtactcc acctacggca agttccttgc cgacggcggg tgctcaggag gtgcttatga 300
cataataatt tgtgacgagt gccactccac ggatgccaca tccatcttgg gcacgggac 360
tgtccttgac caagcagaga ctgcgggggc gagactgggt gtgctcgcca ctgctacccc 420
tccgggctcc gtcactgtgt cccatcctaa catcgaggag gttgctctgt ccaccaccgg 480
agagatcccc ttttacggca aggttatccc cctcgagggt atcaaggggg gaagacatct 540
catcttctgc cactcaaaga agaagtgcga cgagctcgcc gcg                                     583

```

<210> 124

<211> 617

<212> DNA

<213> Hepatitis C virus

<400> 124

```

ccttgccgac ggcggtgct caggaggtgc ttatgacata ataatttgtg acgagtgcc 60
ctccacggat gccacatcca tcttgggcat cggcactgtc cttgaccaag cagagactgc 120
gggggcgaga ctggttgtgc tcgccactgc taccctccg ggctccgtca ctgtgtccca 180
tcctaacatc gaggaggttg ctctgtccac caccggagag atcccccttt acggcaaggc 240
tatccccctc gagtgatca aggggggaag acatctcatc ttctgccact caaagaagaa 300
gtgcgacgag ctgcgcgca agctggctgc attgggcatc aatgcctgg cctactaccg 360
cggctcttgac gtgtctgtca tcccgaaccag cggcgatgtt gtctgtgtgt cgaccgatgc 420
tctcatgact ggctttaccg gcgacttcga ctctgtgata gactgcaaca cgtgtgtcac 480
tcagacagtc gatttcagcc ttgacctac ctttaccatt gagacaacca cgctcccca 540
ggatgctgtc tccaggactc aacgccgggg caggactggc aggggggaagc caggcatcta 600
tagatttgtg gcaccgg                                     617

```

<210> 125

<211> 559

<212> DNA

<213> Hepatitis C virus

<400> 125

```

ctccacggat gccacatcca tcttgggcat cggcactgtc cttgaccaag cagagactgc 60
gggggcgaga ctggttgtgc tcgccactgc taccctccg ggctccgtca ctgtgtccca 120
tcctaacatc gaggagggtg ctctgtccac caccggagag atcccccttt acggcaaggc 180
tatccccctc gaggtgatca aggggggaag acatctcatc ttctgccact caaagaagaa 240
gtgcgacgag ctgcgccgca agctggtcgc attgggcata aatgccgtgg cctactaccg 300
cggctcttgac gtgtctgtca tcccgaccag cggcgatgtt gtcgtcgtgt cgaccgatgc 360
tctcatgact ggctttaccg gcgacttoga ctctgtgata gactgcaaca cgtgtgtcac 420
tcagacagtc gatttcagcc ttgaccctac ctttaccatt gagacaacca cgctcccca 480
ggatgctgtc tccaggactc aacgccgggg caggactggc agggggaagc caggcatcta 540
tagatttgtg gcaccgggg 559

```

<210> 126

<211> 475

<212> DNA

<213> Hepatitis C virus

<400> 126

```

tgtgatagac tgcaacacgt gtgtcactca gacagtcat ttcagccttg accctacctt 60
taccattgag acaaccacgc tccccagga tgctgtctcc aggactcaac gccggggcag 120
gactggcagg ggggaagccag gcatctatag atttgtggca ccgggggagc gcccctccg 180
catgttcgac tcgtccgtcc tctgtgagtg ctatgacgag ggctgtgctt ggtatgagct 240
cacgcccgcc gagactacag ttaggctacg agcgtacatg aacaccccg ggcttcccg 300
gtgccaggac catcttgaat tttgggaggg cgtctttacg ggctcactc atatagatgc 360
ccacttttta tcccagacaa agcagagtgg ggagaacttt cttacctgg tagctacca 420
agccaccgtg tgcgctaggg ctcaagcccc tccccatcg tgggaccaga tgtgg 475

```

<210> 127

<211> 390

<212> DNA

<213> Hepatitis C virus

<400> 127

```

tagatttgtg gcaccggggg agcgccctc cggcatgttc gactcgtccg tcctctgtga 60
gtgctatgac gcgggctgtg cttgggtatga gctcacgcc gccgagacta cagttaggct 120
acgagcgtac atgaacaccc cggggcttcc cgtgtgccag gaccatcttg aattttggga 180
gggcgtcttt acgggcctca ctcatataga tgccacttt ttatcccaga caaagcagag 240
tggggagaaac tttccttacc tggtagcgta ccaagccacc gtgtgcgcta gggctcaagc 300
ccctcccca tcgtgggacc agatgtggaa gtgtttgatc cgccttaaac ccaccctcca 360
tgggccaaca cccctgctat acagactggg 390

```

<210> 128

<211> 155

<212> DNA

<213> Hepatitis C virus

<400> 128

```

acgagcacct ggggtgctcgt tggcggcgtc ctggctgtc tggccgcgta ttgcctgtca 60

```

```
acaggctgcg tggatcatagt gggcaggatc gtcttgtccg ggaagccggc aattatacct 120
gacagggagg ttctctacca ggagttcgat gagat 155
```

```
<210> 129
<211> 56
<212> DNA
<213> Hepatitis C virus
```

```
<400> 129
ggctgctctg gccgcgtatt gcctgtcaac aggctgcgtg gtcatagtgg gcagga 56
```

```
<210> 130
<211> 625
<212> DNA
<213> Hepatitis C virus
```

```
<400> 130
ttttacagct gccgtcacca gccactaac cactggccaa accctcctct tcaacatatt 60
ggggggggtgg gtggtgccc agctcgccgc ccccggtgcc gctactgcct ttgtgggtgc 120
tggcctagct ggcgcgccca tcggcagcgt tggactgggg aaggctcctg tggacattct 180
tgcagggtat ggcgcggggc tggcgggagc tcttgtagca ttcaagatca tgagcgggtga 240
ggtcccctcc acggaggacc tggatcaatct gctgcccgcc atcctctcgc ctggagccct 300
tgtagtcggt gtggtctgcg cagcaatact gcgcgggcac gttggcccg ggcagggggc 360
agtgcgaatgg atgaaccggc taatagcctt cgcctcccg ggggaacctg tttccccccac 420
gcactacgtg ccggagagcg atgcagccgc ccgcgtcact gccatactca gcagcctcac 480
tgtaaccagg ctccctgaggc gactgcatca gtggataagc tcggagtgtg ccactccatg 540
ctccggttcc tggctaaggg acatctggga ctggatatgc gaggtgctga gcgactttta 600
gacctggctg aaagccaagc tcatg 625
```

```
<210> 131
<211> 623
<212> DNA
<213> Hepatitis C virus
```

```
<400> 131
tactgccttt gtgggtgctg gcctagctgg cgcgcgccatc ggcagcgttg gactggggaa 60
ggtcctcgtg gacattcttg cagggtatgg cgcgggcgtg gcgggagctc ttgtagcatt 120
caagatcatg agcgggtgagg tcccctccac ggaggacctg gtcaatctgc tgcccgccat 180
cctctcgcct ggagcccttg tagtcggtgt ggtctgcgca gcaatactgc gccggcacgt 240
tggcccgggc gagggggcag tgcaatggat gaaccggcta atagccttcg cctcccgggg 300
gaacctgtt tccccacgc actacgtgcc ggagagcgat gcagccgcc gcgtcactgc 360
catactcagc agcctcactg taaccagct cctgaggcga ctgcatcagt ggataagctc 420
ggagtgtacc actccatgct ccggttcttg gctaaggagc atctgggact ggatatgcga 480
ggtgctgagc gactttaaga cctggctgaa agccaagctc atgccacaac tgccctgggat 540
tccctttgtg tccctgccgc gcgggtatag gggggtctgg cgaggagacg gcattatgca 600
cactcgtgc cactgtggag ctg 623
```

```
<210> 132
<211> 678
<212> DNA
```


<213> Hepatitis C virus

<400> 132

```
cctcgtggac attcttgcag ggtatggcgc gggcgtggcg ggagctcttg tagcattcaa 60
gatcatgagc ggtgaggtcc cctccacgga ggacctggtc aatctgctgc ccgccatcct 120
ctcgcctgga gcccttgtag tcggtgtggt ctgcgagca atactgcgcc ggcacgttgg 180
cccgggcgag ggggcagtgc aatggatgaa ccggctaata gccttcgcct cccgggggaa 240
ccatgtttcc cccacgcact acgtgccgga gagcgatgca gccgcccgcg tactgccat 300
actcagcagc ctcaactgtaa cccagctcct gaggcgactg catcagtgga taagctcgga 360
gtgtaccact ccatgctccg gttcctggct aagggacatc tgggactgga tatgagaggt 420
gctgagcgac ttttaagacct ggctgaaagc caagctcatg ccacaactgc ctgggattcc 480
ctttgtgtcc tgccagcgcg ggtatagggg ggtctggcga ggagacggca ttatgcacac 540
tcgctgccac tgtggagctg agatcactgg acatgtcaaa aacgggacga tgaggatcgt 600
cggtcctagg acctgcagga acatgtggag tgggacgttc cccattaacg cctacaccac 660
gggcccctgt actccct 678
```

<210> 133

<211> 436

<212> DNA

<213> Hepatitis C virus

<400> 133

```
tgcagggtat ggcgcggggcg tggcgggagc tcttgtagca ttcaagatca tgagcgggtga 60
gggtcccctcc acggaggacc tggatcaatct gctgcccgcc atcctctcgc ctggagccct 120
tgtagtccgt gtggtctgcg cagcaatact gcgcgggcac gttggcccgg gcgagggggc 180
agtgcattgg atgaaccggc taatagcctt cgcctcccgg ggggaacctg tttccccac 240
gcactacgtg ccggagagcg atgcagccgc ccgcgtcact gccatactca gcagcctcac 300
tgtaaccag ctctgagggc gactgcatca gtggataagc tcggagtgtg cactccatg 360
ctccggttcc tggctaaggg acatctggga ctggatatgc gaggtgctga gcgactttaa 420
gacctggctg aaagcc 436
```

<210> 134

<211> 164

<212> DNA

<213> Hepatitis C virus

<400> 134

```
agcccttgta gtccgtgtgg tctgcgcagc aatactgcgc cggcacgttg gcccgggcga 60
gggggcagtg caatggatga accggctaata agccttcgcc tcccggggga accatgtttc 120
ccccacgcac tacgtgccgg agagcgatgc agccgcccgc gtca 164
```

<210> 135

<211> 496

<212> DNA

<213> Hepatitis C virus

<400> 135

```
cgcctcccgg ggggaacctg tttccccac gcactacgtg ccggagagcg atgcagccgc 60
ccgcgtcact gccatactca gcagcctcac tgtaaccag ctctgagggc gactgcatca 120
gtggataagc tcggagtgtg cactccatg ctccggttcc tggctaaggg acatctggga 180
ctggatatgc gaggtgctga gcgactttaa gacctggctg aaagccaagc tcatgccaca 240
```

```

actgcctggg attccctttg tgtcctgcc a gcgcgggtat aggggggtct ggcgaggaga 300
cggcattatg cacactcgct gccactgtgg agctgagatc actggacatg tcaaaaaacgg 360
gacgatgagg atcgtcggtc ctaggacctg caggaacatg tggagtggga cgttccccat 420
taacgcctac accacggggc cctgtactcc ccttctcgcg ccgaactata agttcgcgct 480
gtggaggggtg tctgca                                     496

```

<210> 136

<211> 926

<212> DNA

<213> Hepatitis C virus

<400> 136

```

tacgtgccgg agagcgatgc agccgcccgc gtcactgcca tactcagcag cctcactgta 60
accagctccc tgaggcgact gcatcagtgg ataagctcgg agtgtaccac tccatgctcc 120
ggttcctggc taaggacat ctgggactgg atatgcgagg tgctgagcga ctttaagacc 180
tggctgaaag ccaagctcat gccacaactg cctgggattc cctttgtgtc ctgccagcgc 240
gggtataggg ggggtctggcg aggagacggc attatgcaca ctcgctgcca ctgtggagct 300
gagatcactg gacatgtcaa aaacgggacg atgaggatcg tcggtcctag gacctgcagg 360
aacatgtgga gtgggacgtt ccccattaac goctacacca cgggcccctg tactccccct 420
cctgcgccga actataagtt cgcgctgtgg aggggtgtctg cagaggaata cgtggagata 480
aggcgggtgg gggacttcca ctacgtatcg ggtatgacta ctgacaatct taaatgcccg 540
tgccagatcc catcgcccga atttttcaca gaattggacg ggggtgcgct acacagggtt 600
gcgccccctt gcaagccctt gctgcgggag gaggatatcat tcagagtagg actccacgag 660
taccgggtgg ggtcgcaatt accttgcgag cccgaaccgg acgtagccgt gttgacgtcc 720
atgctcactg atccctccca tataacagca gaggcgggccg ggagaagggt ggcgagaggg 780
tcacccccct ctatggccag ctccctcggt agccagctgt ccgctccatc totcaaggca 840
acttgcaccg ccaaccatga ctcccctgac gccgagctca tagaggctaa cctcctgtgg 900
aggcaggaga tgggcggcaa catcac                                     926

```

<210> 137

<211> 850

<212> DNA

<213> Hepatitis C virus

<400> 137

```

actcagcagc ctcactgtaa cccagctcct gaggcgactg catcagtgga taagctcgga 60
gtgtaccact ccatgctccg gttcctggct aaggacatc tgggactgga tatgcgaggt 120
gctgagcgac tttaagacct ggctgaaagc caagctcatg ccacaactgc ctgggattcc 180
ctttgtgtcc tgccagcgcg ggtatagggg ggtctggcga ggagacggca ttatgcacac 240
tcgctgccac tgtggagctg agatcactgg acatgtcaaa aacgggacga tgaggatcgt 300
cggtcctagg acctgcagga acatgtggag tgggacgttc cccatthaac cctacaccac 360
gggcccctgt actccccctt ctgcgccgaa ctataagttc gcgctgtgga ggggtgtctgc 420
agaggaatac gtggagataa ggcggtggg ggaacttcac tacgtatcgg gtatgactac 480
tgacaatctt aaatgcccgt gccagatccc atcgcccga tttttcacag aattggacgg 540
gggtgcgcta cacaggtttg cgcgcccttg caagcccttg ctgcgggagg aggtatcatt 600
cagagtagga ctccacgagt acccggtggg gtcgcaatta ccttgcgagc ccgaaccgga 660
cgtagccgtg ttgacgtcca tgetcactga tccctccat ataacagcag aggcggccgg 720
gagaagggtg gcgagagggt ccccccttc tatggccagc tcctcggcta gccagctgtc 780
cgctccatct ctcaaggcaa cttgcaccgc caaccatgac tcccctgacg ccgagctcat 840
agaggctaac                                     850

```

<210> 138
 <211> 749
 <212> DNA
 <213> Hepatitis C virus

<400> 138
 cagcctcact gtaaccacagc tcctgaggcg actgcatcag tggataagct cggagtgtac 60
 cactccatgc tccggttccg ggctaaggga catctgggac tggatatgcg aggtgctgag 120
 cgactttaag acctggctga aagccaagct catgccacaa ctgcctggga ttccctttgt 180
 gtccctgccag cgcgggtata ggggggtctg gcgaggagac ggcattatgc acactcgtg 240
 ccactgtgga gctgagatca ctggacatgt caaaaacggg acgatgagga tcgtcgggtcc 300
 taggacctgc aggaacatgt ggagtgggac gttccccatt aacgcctaca ccacgggccc 360
 ctgtactccc cttcctgcgc cgaactataa gttcgcgctg tggaggggtgt ctgcagagga 420
 atacgtggag ataaggcggg tgggggactt ccactacgta tcgggtatga ctactgacaa 480
 tcttaaatgc ccgtgccaga tcccatcgcc cgaatttttc acagaattgg acgggggtgcg 540
 cctacacagg tttgcgcccc cttgcaagcc cttgctgcgg gaggagggtat cattcagagt 600
 aggactccac gagtaccggg tggggtcgca attaccttgc gagcccgaac cggacgtagc 660
 cgtgttgacg tccatgctca ctgatccctc ccatataaca gcagaggcgg ccgggagaag 720
 gttggcgaga gggtcacccc cttctatgg 749

<210> 139
 <211> 257
 <212> DNA
 <213> Hepatitis C virus

<400> 139
 gacctggctg aaagccaagc tcatgccaca actgcctggg attccctttg tgtcctgcca 60
 gcgcgggtat aggggggtct ggcgaggaga cggcattatg cacactcgct gccactgtgg 120
 agctgagatc actggacatg tcaaaaacgg gacgatgagg atcgtcggtc ctaggacctg 180
 caggaacatg tggagtggga cgttcccat taacgcctac accacggggc cctgtactcc 240
 ccttctcgcg ccgaact 257

<210> 140
 <211> 285
 <212> DNA
 <213> Hepatitis C virus

<400> 140
 tgagatcact ggacatgtca aaaacgggac gatgaggatc gtcggtccta ggacctgcag 60
 gaacatgtgg agtgggacgt tccccattaa cgcctacacc acggggccct gtactcccct 120
 tcctgcgccg aactataagt tcgcgctgtg gagggtgtct gcagaggaat acgtggagat 180
 aaggcgggtg ggggacttcc actacgtatc gggatgact actgacaatc ttaaatgcc 240
 gtgccagatc ccatcgcccc aattttttcac agaattggac ggggt 285

<210> 141
 <211> 228
 <212> DNA
 <213> Hepatitis C virus

<400> 141
 catagaggct aacctcctgt ggaggcagga gatggggcggc aacatcacca gggttgagtc 60

```

agagaacaaa gtggtgattc tggactcctt cgatccgctt gtggcagagg aggatgagcg 120
ggaggtctcc gtacctgcag aaattctcg gaagtctcgg agattcgccc gggccctgcc 180
cgtctgggcy cggccggact acaaccccc gctagtagag acgtggaa 228

```

```

<210> 142
<211> 273
<212> DNA
<213> Hepatitis C virus

```

```

<400> 142
ccatggctgc ccgtaccac ctccacggtc cctcctgtg cctccgcctc ggaaaaagcg 60
tacgggtggc ctcaccgaat caaccctatc tactgccttg gccgagcttg ccaccaaag 120
ttttggcagc tcctcaactt ccggcattac gggcgacaat acgacaacat cctctgagcc 180
cgccccctct ggctgcccc ccgactccga cgttgagtcc tattcttcca tgccccccct 240
ggagggggag cctggggatc cggatctcag cga 273

```

```

<210> 143
<211> 412
<212> DNA
<213> Hepatitis C virus

```

```

<400> 143
ttcctggaca ggcgcactcg tcaccccgty cgtcgcgga gaacaaaaac tgcccatcaa 60
cgactgagc aactcgttgc tacgccatca caatctggtg tattccacca cttaocgag 120
tgcttgccaa aggcagaaga aagtcacatt tgacagactg caagttcttg acagccatta 180
ccaggacgtg ctcaaggagg tcaaagcagc ggcgtcaaaa gtgaaggcta acttgctatc 240
cgtagaggaa gcttgagcc tgacgcccc acattcagcc aaatccaagt ttggctatgg 300
ggcaaaagac gtccgttgcc atgccagaaa ggccgtagcc cacatcaact ccgtgtggaa 360
agaccttctg gaagacagt taacaccaat agacactacc atcatggcca ag 412

```

```

<210> 144
<211> 903
<212> DNA
<213> Hepatitis C virus

```

```

<400> 144
ggctaacttg ctatccgtag aggaagcttg cagcctgacg cccccacatt cagccaaatc 60
caagtttggc tatggggcaa aagacgtccg ttgccatgcc agaaaggccg tagcccacat 120
caactccgtg tggaaagacc ttctggaaga cagtgtaca ccaatagaca ctaccatcat 180
ggccaagaac gaggttttct gcggtcagcc tgagaagggg ggtcgtaagc cagctcgtct 240
catcgtgttc cccgacctgg gcggtgcgct gtgcgagaag atggccctgt acgacgtgg 300
tagcaagctc cccctggccg tgatgggaag ctctacgga ttccaatact caccaggaca 360
gcgggttgaa ttctcgtgc aagcgtggaa gtccaagaag acccgatgg ggttctcgta 420
tgataccgc tgttttgact ccacagtcac tgagagcgac atccgtacgg aggaggcaat 480
ttaccaatgt tgtgacctgg accccaagc ccgctggcc atcaagtccc tcaactgagag 540
gctttatgtt gggggccctc ttaccaattc aaggggggaa aactgoggct accgcaggtg 600
ccgcgcagc ggcgtactga caactagctg tggtaacacc ctcaactgct acatcaaggc 660
ccgggcagcc tgcgagccg cagggtcca ggactgcacc atgctcgtgt gtggcgagca 720
cttagtcgtt atctgtgaaa gtgcgggggt ccaggaggac gcggcgagcc tgagagcctt 780
cacggaggct atgaccaggt actccgcccc cccgggggac cccccacaac cagaatacga 840
cttgagctt ataacatcat gtcctccaa cgtgtcagtc gccacgacg gcgctggaaa 900

```

gag

903

<210> 145
 <211> 600
 <212> DNA
 <213> Hepatitis C virus

<400> 145
 agaggaagct tgcagcctga cgccccacaca ttcagccaaa tccaagtttg gctatggggc 60
 aaaagacgtc cgttgccatg ccagaaaggc cgtagcccac atcaactccg tgtggaaaga 120
 ccttctggaa gacagtgtaa caccaataga cactaccatc atggccaaga acgagggtttt 180
 ctgcgttcag cctgagaagg ggggtcgtaa gccagctcgt ctcatcgtgt tccccgacct 240
 gggcgtgcgc gtgtgcgaga agatggccct gtacgacgtg gttagcaagc tccccctggc 300
 cgtgatggga agtcctacg gattccaata ctaccagga cagcgggttg aattcctcgt 360
 gcaagcgtgg aagtccaaga agaccccgat ggggttctcg tatgataccc gctgttttga 420
 ctccacagtc actgagagcg acatccgtac ggaggaggca atttaccat gttgtgacct 480
 ggacccccaa gcccgctgg ccatcaagtc cctcactgag aggcctttatg ttggggggccc 540
 tcttaccat tcaagggggg aaaactgcgg ctaccgcagg tgccgcgcga gcggcggtact 600

<210> 146
 <211> 781
 <212> DNA
 <213> Hepatitis C virus

<400> 146
 ccttctggaa gacagtgtaa caccaataga cactaccatc atggccaaga acgagggtttt 60
 ctgcgttcag cctgagaagg ggggtcgtaa gccagctcgt ctcatcgtgt tccccgacct 120
 gggcgtgcgc gtgtgcgaga agatggccct gtacgacgtg gttagcaagc tccccctggc 180
 cgtgatggga agtcctacg gattccaata ctaccagga cagcgggttg aattcctcgt 240
 gcaagcgtgg aagtccaaga agaccccgat ggggttctcg tatgataccc gctgttttga 300
 ctccacagtc actgagagcg acatccgtac ggaggaggca atttaccat gttgtgacct 360
 ggacccccaa gcccgctgg ccatcaagtc cctcactgag aggcctttatg ttggggggccc 420
 tcttaccat tcaagggggg aaaactgcgg ctaccgcagg tgccgcgcga gcggcggtact 480
 gacaactagc tgtggttaaca cctcacttg ctacatcaag gcccgggcag cctgtcgagc 540
 cgcagggtc caggactgca ccatgctcgt gtgtggcgac gacttagtgc ttatctgtga 600
 aagtgcgggg gtccaggagg acgcggcgag cctgagagcc ttcacggagg ctatgaccag 660
 gtactccgcc cccccgggg accccccaca accagaatac gacttggagc ttataacatc 720
 atgctcctcc aacgtgtcag tgcgccacga cggcgctgga aagaggggtct actaccttac 780
 c 781

<210> 147
 <211> 382
 <212> DNA
 <213> Hepatitis C virus

<400> 147
 cgttatctgt gaaagtgcgg ggggtccagga ggacgcggcg agcctgagag ccttcacgga 60
 ggctatgacc aggtactccg ccccccccg ggacccccca caaccagaat acgacttgga 120
 gcttataaca tcatgctcct ccaacgtgtc agtcgcccac gacggcgctg gaaagagggg 180
 ctactacctt accogtgacc ctacaacccc cctcgcgaga gccgcgtggg agacagcaag 240
 acacactcca gtcaattcct ggctaggcaa cataatcatg tttgccccca cactgtggggc 300

gaggatgata ctgatgaccc atttcttttag cgtcctcata gccagggatc agcttgaaca 360
 ggctcttaac tgtgagatct ac 382

<210> 148
 <211> 268
 <212> DNA
 <213> Hepatitis C virus

<400> 148
 cgtgtcagtc gccacgacg gcgctggaaa gagggctctac taccttaccg gtgaccctac 60
 aacccccctc gcgagagccg cgtgggagac agcaagacac actccagtca attcctggct 120
 aggcaacata atcatgtttg cccccacact gtgggagagg atgatactga tgacccattt 180
 ctttagcgtc ctcatagcca gggatcagct tgaacaggct cttaactgtg agatctacgg 240
 agcctgctac tccatagaac cactggat 268

<210> 149
 <211> 222
 <212> DNA
 <213> Hepatitis C virus

<400> 149
 actccatggc ctcagcgcat ttctactcca cagttactct ccagggtgaaa tcaatagggt 60
 ggccgcatgc ctcagaaaac ttgggggtccc gcccttgoga gcttggagac accggggccc 120
 gagcgccgc gctaggcttc tgtccagagg aggcagggt gccatatgtg gcaagtacct 180
 cttcaactgg gcagtaagaa caaagctcaa actcactcca at 222

<210> 150
 <211> 192
 <212> DNA
 <213> Hepatitis C virus

<400> 150
 ctctccagggt gaaatcaata ggggtggcgc atgcctcaga aaacttgggg tcccgcctt 60
 gcgagcttgg agacaccggg cccggagcgt ccgcgctagg cttctgtcca gaggaggcag 120
 ggctgccata tgtggcaagt acctcttcaa ctgggcagta agaacaaagc tcaaactcac 180
 tccaatagcg gc 192

<210> 151
 <211> 10
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: linker
 sequence

<400> 151
 gggccacgaa 10

<210> 152
<211> 13
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: linker
sequence

<400> 152
ttcgtggccc ctg 13

<210> 153
<211> 138
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: pP6 vector
sequence

<400> 153
ctagccatgg ccgcaggggc cgcggccgca ctagtgggga tccttaatta aagggccact 60
ggggccccc gtaccggcgt ccccggcgcc ggcgtgatca cccctaggaa ttaatttccc 120
ggtgacccc ggggagct 138

<210> 154
<211> 128
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: pB5 vector
sequence

<400> 154
catggccgca ggggccgcgg ccgcactagt ggggatcctt aattaaaggc cactggggc 60
cccccgcggt ccccggcgcc ggcgtgatca cccctaggaa ttaatttccc ggtgacccc 120
ggggagct 128

<210> 155
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer

<400> 155
gcgtttggaa tcactacagg 20

<210> 156
<211> 19
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 156
cacgatgcac gttgaagtg

19